



Hessian Low Mountain Range

Full data report

Generated 2026-08-11

Hesse

NUTS-3 regions: DE721, DE722, DE723, DE724, DE725



Location within Germany



Basemap: © OpenStreetMap contributors © CARTO

Agriculture

Key figures


CROPLAND AREA	NUMBER OF FARMS	AVERAGE FARM SIZE
113,215 ha	3,832 count	54.32 ha
SHARE OF ORGANIC FARMING		
23.1 %		

Map



- Agriculture
- | | |
|----------------------|--|
| winter wheat | sugar beet |
| winter barley | rapeseed |
| winter rye | clover/alfalfa |
| other winter cereals | arable grass |
| spring wheat | permanent grassland |
| spring barley | vineyard |
| spring oat | fruit trees and other woody vegetation |
| maize | hops |
| legumes | other agricultural use |
| potato | |

Chart



Overview

Comparatively high livestock density. High proportion of grassland. High proportion of forest. Relatively small farm sizes and field sizes

Challenges

Feed production problematic (soil unsuitable for cultivation), therefore predominantly grassland farming. Decline in dairy farming (-6% dairy cows, -14% farms). Competition for land with energy production (wind power, photovoltaics)



Climate

Key figures



GROWING DEGREE DAYS, ANNUAL SUM (BASE 5 °C)	MEAN ANNUAL TEMPERATURE	ANNUAL PRECIPITATION
1,802 °C·d +1,091 °C·d by 2071-2100	8.7 °C +3.9 °C by 2071-2100	862 mm +2.9 % by 2071-2100
PRECIPITATION OF WARMEST QUARTER		
220 mm -8.6 % by 2071-2100		

Map



GDD
Baseline





GDD
2071-2100




Mean temp.
Baseline




Mean temp.
2071-2100





Precipitation
Baseline




Precipitation
2071-2100



Summer precip.
Baseline



Summer precip.
2071-2100




Heat accumulated above 5 °C over the year - a measure of how much growing season a crop gets.

Mean annual air temperature, averaged over the reference period.

Total precipitation summed over the year.

Precipitation falling in the warmest three-month quarter - relevant to summer drought stress.

Chart



Overview


Exposed to the prevailing wind direction, slope rainfall, lower temperatures compared to the plains.

Soil

Key figures


NITROGEN SURPLUS	PHOSPHORUS SURPLUS	GROUNDWATER ABSTRACTION (NON-PUBLIC SUPPLY)
No data available kg N/ha UAA	No data available kg P/ha UAA	2,244 1000 m3

Map



Legend shows the dominant semantic soil groups for this Living Lab; raw BUEK IDs stay in the data only as provenance.

Chart



Overview

Soil types are highly variable: Podzolic brown soils with low base saturation, red sandstone, brown soils with high to medium base saturation, pseudogley, parabraun soils from loess with high base saturation, rendzina on calcareous sites (locally limited).

Challenges


Soil erosion during heavy rainfall

Socio-economic


Key figures

TOTAL POPULATION	GDP PER CAPITA	UNEMPLOYMENT RATE
1,041,453 persons	40,568 EUR	5.34 %
DISPOSABLE HOUSEHOLD INCOME PER CAPITA		
26,575.6 EUR		

Map



Chart



Overview

Peripheral areas — long distances to regional centres. Population structure: decline, ageing. Comparatively low disposable household income.

Challenges


Sales market for products (milk, meat, etc.) located far away, lower direct marketing opportunities, Long distances to slaughterhouses and processing companies. Shortage of workers due to migration and ageing population.

Landscape


Key figures

FOREST AREA	NATURA 2000 AREA	NATURE RESERVES AREA
214,513 ha	134,837.9 ha	8,263.6 ha
SHARE OF SEALED SURFACE		
15 %		

Map



Chart



Overview

Low mountain regions 400-1000 m altitude with high relief energy.