"The color of a flower field changes with a butterfly's flight vector" is a metaphor for homochirality colorimetry via chiral nanostructure arrays

Zaid Haddadin¹, Anna My Nguyen², Paula Kirya², Lisa V. Poulikakos^{2,3}


- 1. Electrical & Computer Engineering, University of California San Diego, La Jolla, CA 92093
- 2. Mechanical & Aerospace Engineering, University of California San Diego, La Jolla, CA 92093
- 3. Program of Materials Science & Engineering, University of California San Diego, La Jolla, CA 92093

Introduction

Fibrous or filamentous biological structures, like microbial mats, can rotate a light wave's travel vector [1,2]. However, this rotation is too minimal to measure without meter-scale tools [3]. This work explores miniaturizing these tools to the nanometer level for in-situ applications.


One possibility is rotating a light vector into a circular path, either clockwise or counterclockwise [3]. Homochirality is when an orientation of circular light is favored over the other [3,4,5]. Detecting homochiral light is an agnostic biosignature of life [5].

We investigate how this biosignature could be detected using arrays of nanoscale particles. By using visible light, our optical arrays also act as colorimetric sensors of homochirality [3,6,7].



Materials and methods

We investigate how a cut to one corner of square- and rectangle-shaped nanostructures affects their sensitivities to differentiating between incident clockwise or counterclockwise oriented circular light.




Results • Differentiation of clockwise and counter-clockwise polarization states of circular light was achieved. • Squares: Figure 3 shows cuts to the square structure affect the reflectance response but result in non-differentiable outputs. Table 1 displays the non-differentiable colorimetry results. Figure 4 examines the near-field response: cuts simplify the problem from a quadrupole to a dipole, which dampens the reflectance response due to a decreased energy output. • Rectangles: Simplifying the problem in rectangles from four nodes to two nodes extinguishes the longer-wavelength reflectance peaks, as shown in Figure 5's "Medium Cut" group. This unequally dampens the shorter-wavelength peaks of the two polarization states. Both these observations contribute to the differentiable colorimetry results in Table 2. **Table 1:** Colorimetry results Figure 3: Applying cuts to a square-shaped nanoparticle from Figure I reflectances. No Cut **Small Cut Medium Cut** Clockwise Both Counter-Clockwise polarization clockwise Counter-clockwise states show No Cut #598C18 The prior peak The two peaks splits into two collapse into dampened Small Cut #428716 one, but the amplitude doesn't recover. Medium Cut #1C6117 0.0 ^{_} 380 536 550 550 380 Wavelength (nm) Wavelength (nm) Wavelength (nm) $\left|\left|\mathbf{E}_{0}\right|\right|_{2}^{2}$ Clockwise 200 100 Counter-50.0 Figure 5: Applying cuts to a rectangle-shaped nanoparticle **Table 2:** Colorimetry results from Figure III reflectances. No Cut Small Cut Medium Cut —— Clockwise Counter-Clockwise ····· Counter-clockwise clockwise 0.8 #484564 No Cut The longer-wavelength The shorterpeaks are extinguished. wavelength peak #424059 #433F62 The clockwise polarization Both polarization of clockwise peak is disproportionately states show the polarization slightly dampens. dampened. same response. Medium Cut #31352A #343146 0.0 ^{_} 380 646 380 440 Wavelength (nm) Wavelength (nm) Wavelength (nm)

Conclusions

Biological media with fibrous structures can alter light polarization, an agnostic indicator of life [1,2]. Designing nano-optical arrays enables miniaturized optical sensors for polarized light, which may be useful for *in-situ* investigations. Our study showed that nano-optical arrays can differentiate clockwise and counter-clockwise circularly polarized light, allowing colorimetry homochirality detection (see Figure 6).








However, the reason for this remains unclear. Figure 7 suggests that under clockwise light, the dipole charge aligns with the sideways flux, reducing light reflected towards the imaging apparatus. Under counter-clockwise light, the dipole charge opposes this flux. This could explain the disproportionate dampening in reflectance and the distinct colorimetry results.

Figure 7: Near-field of medium cut rectangle.

Electric field vectors (size is flux difference, not to scale)







Literature cited

- l. Ghosh N, Wood MFG, Li S hong, Weisel RD, Wilson BC, Li RK, et al. Journal of Biophotonics. 2009;2(3):145-56
- 2. Sparks WB, Parenteau MN, Blankenship RE, et al. Astrobiology. 2021 Feb;21(2):219-34.
- 3. Kwon J, Park KH, Choi WJ, et al. Acc Chem Res. 2023 Jun 20;56(12):1359–72.
- Avnir D. New Astronomy Reviews. 2021 Jun 1;92:101596
- Neveu M, Hays LE, Voytek MA, et al. Astrobiology. 2018;18(11):1375–402.
- Haddadin Z, Khan S, Poulikakos LV. ACS Photonics. 2024 Jan 17;11(1):187-195.
- 7. Fairchild MD. Color Appearance Models. John Wiley & Sons. 2005; 2nd Ed.

Acknowledgments

Research funding received through the UC San Diego MRSEC Seed Grant and from the Arnold and Mabel Beckman Foundation. Travel funding from AbGradCon24.

Thank you to Julia Holland for brainstorming the presentation's title, and to the rest of my coworkers in the Poulikakos Lab for the research discussions and continued support ©

Further information

I'm at **zalhadda@ucsd.edu** for further questions & discussions. Scan the below QR code for a PDF copy of this poster.





