Quantifying symmetry violations for optimising algorithmic designs


Zaid Haddadin¹, Dae Yong Kim², Jiuk Byun³, Lisa V. Poulikakos^{2,3}

1.Electrical & Computer Engineering, University of California San Diego 2.Mechanical & Aerospace Engineering, University of California San Diego 3. Materials Science & Engineering, University of California San Diego

Introduction Structures with sufficient asymmetry can yield unique light-matter interactions under differing polarisations of light (1,2). Light polarised to the norizontal & vertical responses (bottom) in rectangular nanograting structures (outlined in Axes of structures **lacking asymmetry** result in identical light-matter responses for varying polarisations of light (1,2).Light polarised to the diagonal states (top) show the same, but oppositely signed, electric field responses (bottom) in rectangular nanograting structures (outlined in black). Designing sufficiently asymmetric **structures is vital** for creating selective light-matter interactions across a wide arrangement of polarisations of light.

Materials & Methods


The degree of symmetry of a shape can **be quantified** by the distance of its projection onto a subspace of symmetric shapes (3).



The distance from the shape to a subspace of symmetric shapes is inversely proportional the shape's degree of symmetry.

Shape)))

Averaging the distance across a range of angles provides rotational and reflective symmetry scores:



Number of angles


Manipulating the symmetry operator specifies the type of symmetry scored: rotational or reflective.

Results

The symmetry scores corresponding to the degree of rotational or reflective symmetries were **calculated for various shapes.** These calculations transform symmetry from a binary concept – a shape is either symmetrical or asymmetrical – to a continuous spectrum: a shape has a certain degree of symmetry. These shapes were taken from an informations security hash visualisation (4) known as identicons (5).


The symmetry scores below take into account the shape (in white) in the context of the empty **space surrounding it (in black)** as light-matter interactions are affected by the empty space around a structure along with the geometry of the structure. The shapes are presented in order of decreasing rotational symmetry score from the left to the right.

Rotational & Reflective Symmetry Scores




First Steps Towards a Genetic Algorithm


The ability to determine the degree of symmetry of a shape can provide future insight towards engineering selective light-matter interactions across differing polarisations of light. To this extent, an evolutioninspired, genetic-based machine learning model is being constructed that takes advantage of symmetry scoring to construct an idealised asymmetric structure. Such an algorithm is envisioned to use the aforementioned shapes (among others) as elementary building blocks.



An M-by-N grid acts as a blueprint denoting the building block shape and a respective angle (0, 90, 180, or 270 degrees) for an asymmetric structure.



A decimal-to-binary transformation to create a binary-based "DNA analogue" that can evolve within a machine learning model's parameters.



Each decimal-containing grid square will refer to a building block shape. When put together, the building block shapes will result in a new asymmetric structure.

Conclusions

The asymmetry inherent in a geometric shape affects the expected light-matter interactions. Quantifying the degree of symmetry paves the path towards correlating the effects of changing specific symmetries to the observed light-matter interactions. This can open a future where the idealised asymmetry of a structure can be engineered for an intended application.

The ability to select for light-matter interactions through a geometric-tuning of structures can have implications for sensing applications in several fields such as astrobiology (6), geology (7), and medicine (1,2). Future steps of this work aim to realise the ideal combination of elementary building blocks to create maximal asymmetry.

Literature Cited

- 1. Poulikakos LV, Lawrence M, Barton DR, Jeffrey SS, Dionne JA. ACS Photonics. 2020 Nov 18;7(11):3216-
- 2. Haddadin, Z., Pike, T., Moses, J. and Poulikakos, L., 2021. Journal of Materials Chemistry C, 9(35), pp.11619-11639.
- 3. Dryzun, C. and Avnir, D., 2009. Physical Chemistry Chemical Physics, 11(42), p.9653.
- 4. Perrig, Adrian, and Dawn Song. International Workshop on Cryptographic Techniques and E-Commerce. Vol. 25. 1999.
- 5.Park, D., 2007. Available at:
- https://web.archive.org/web/20080703155519/http: //www.docuverse.com/blog/donpark/2007/01/18/ visual-security-9-block-ip-identification> [Accessed 29] June 2022].
- 6 .Committee on the Planetary Science and Astrobiology Decadal Survey Space Studies Board. 2022 [cited 2022 Jun 5]. Available from: https:// www.nap.edu/read/26522/chapter/1
- 7. Dash JG, Rempel AW, Wettlaufer JS. Rev Mod Phys. 2006 Jul 12;78(3):695-741.

Acknowledgements

Thank you to Samantha Bordy, Paula Kirya, Kaylani Patel, Anna My Nguyen, Martin Sanner, & Roxanna Sheets for being involved in the drafting process. Funding received through the MRSEC seed grant.

Further Information

Contact me at zalhadda@ucsd.edu if you have a question or comment. Our lab is at https://poulikakos.ucsd.edu



