

J1I-MC-GZQO Protocol v3.0 (synthetic excerpt)

1 Protocol Summary

Protocol J1I-MC-GZQO (synthetic excerpt). A Phase 3, randomized, double-blind, placebo-controlled study of once-weekly retatrutide in participants with gout and obesity or overweight. This document is a fictional excerpt produced for TrialBox testing; it is not the sponsor's protocol.

2 Objectives and Endpoints

Primary: proportion of participants with serum urate below 6.0 mg/dL at week 52. Key secondary: gout flare rate, change in body weight.

3 Study Design

Approximately 900 participants will be randomized 1:1:1 to retatrutide 4 mg, 8 mg or placebo for 52 weeks after a 4-week screening period.

5 Study Population

5.1 Inclusion Criteria

1. Male or female participants aged 18 years or older at screening.
2. Diagnosis of gout according to the 2015 ACR/EULAR gout classification criteria.
3. Body mass index (BMI) ≥ 27 kg/m² at screening.
4. Serum urate ≥ 6.8 mg/dL at screening, or ≥ 6.0 mg/dL if receiving urate-lowering therapy.
5. At least 2 gout flares within the 12 months before screening.
6. Receiving stable urate-lowering therapy for at least 3 months (continuous within the past 6 months, gaps ≤ 30 days).
7. Willing and able to self-inject study intervention once weekly.
8. Estimated GFR ≥ 30 mL/min/1.73 m² (CKD-EPI 2021) at screening.
9. Followed at the rheumatology or metabolism clinic with at least one visit in the past 12 months.

5.2 Exclusion Criteria

1. Type 1 diabetes mellitus.
2. History of acute or chronic pancreatitis.
3. Personal history of medullary thyroid carcinoma or multiple endocrine neoplasia type 2.
4. Use of a GLP-1 receptor agonist or GIP/GLP-1 receptor agonist within 90 days before screening.
5. Prior or planned bariatric surgery.
6. HbA1c $> 10\%$ at screening.
7. Malignancy within the past 5 years, except non-melanoma skin cancer.
8. Pregnant or breastfeeding, or positive pregnancy test.
9. Uncontrolled hypertension (systolic blood pressure ≥ 160 mmHg) at screening.

10. Myocardial infarction or stroke within 90 days before screening.
11. Acute gout flare ongoing at screening (within 14 days).
12. Planned surgery during the study period.
13. ALT >3× upper limit of normal (>120 U/L) at screening.
14. Participation in another interventional study within 30 days before screening.

6 Study Intervention

Retatrutide is administered subcutaneously once weekly.