



Critical transitions in chronic disease: transferring concepts from ecology to systems medicine

Christophe Trefois¹, Paul MA Antony¹, Jorge Goncalves², Alexander Skupin^{3,4} and Rudi Balling¹

Ecosystems and biological systems are known to be inherently complex and to exhibit nonlinear dynamics. Diseases such as microbiome dysregulation or depression can be seen as complex systems as well and were shown to exhibit patterns of nonlinearity in their response to perturbations. These nonlinearities can be revealed by a sudden shift in system states, for instance from health to disease. The identification and characterization of early warning signals which could predict upcoming critical transitions is of primordial interest as prevention of disease onset is a major aim in health care. In this review, we focus on recent evidence for critical transitions in diseases and discuss the potential of such studies for therapeutic applications.

Addresses

¹ Experimental Neurobiology Group, Luxembourg Centre for Systems Biomedicine (LCSB), University of Luxembourg, Campus Belval, 7 Avenue des Hauts-Fourneaux, L-4362 Esch-sur-Alzette, Luxembourg

² Systems Control Group, Luxembourg Centre for Systems Biomedicine (LCSB), University of Luxembourg, Campus Belval, 7 Avenue des Hauts-Fourneaux, L-4362 Esch-sur-Alzette, Luxembourg

³ National Center for Microscopy and Imaging Research, University of California San Diego, 9500 Gilman Drive, La Jolla, CA, United States

⁴ Integrative Cell Signalling Group, Luxembourg Centre for Systems Biomedicine (LCSB), University of Luxembourg, Campus Belval, 7 Avenue des Hauts-Fourneaux, L-4362 Esch-sur-Alzette, Luxembourg

Corresponding author: Balling, Rudi (rudi.balling@uni.lu)

Current Opinion in Biotechnology 2015, 34:48–55

This review comes from a themed issue on **Systems biology**

Edited by **Sarah Maria Fendt** and **Costas D Maranas**

For a complete overview see the [Issue](#) and the [Editorial](#)

Available online 10th December 2014

<http://dx.doi.org/10.1016/j.copbio.2014.11.020>

0958-1669/© 2014 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-SA license (<http://creativecommons.org/licenses/by-nc-sa/3.0/>).

Critical transitions in complex systems

Biological systems are complex, characterized by emerging behavior and often obey nonlinear dynamics (Box 1). In many cases this nonlinear behavior of biological systems leads to tipping points where the equilibrium state (Box 1) of the system abruptly changes from one stable state to another (Box 1). This change is also called critical transition (Box 1) or regime shift [1]. Although sudden state transitions such as phase transitions or exothermal reactions are established concepts in physics and chemistry,

increasing awareness rises that also complex biological systems can exhibit abrupt changes in their dynamics. The idea that natural systems might exhibit sudden changes in their dynamical states originated mostly from theoretical models over 50 years ago [2–4] and are based on mathematical catastrophic bifurcation theory [5]. These studies, motivated by descriptions of magnetic systems, laid the concept that natural systems might have alternative stable states and thereby undergo critical transitions (Box 1), typically without obvious warning signals (Figure 1). At that time, these theories lacked robust empirical evidence [6].

Only in the last decade, several studies provided evidence for the existence of critical transitions in natural and societal systems. In ecology [7], there are examples of the desertification of Mediterranean arid ecosystems [8] or the tree abundance in tropical forest and savannah [9]. Critical transitions have also been associated with the eutrophication of lakes [10], collapse of fish populations due to overfishing [11] and algae overgrowth in Caribbean coral reefs [12]. On larger scale such as the earth's climate system, reduction of Greenland ice sheets or melting of arctic sea-ice has been associated with a potential transition in the global climate system, which may or may not be reversible [13]. Similar to ecological systems, humans have complex traits. When considered as complex systems, both present positive and negative feedback loops, inherent nonlinearity and hysteresis (Box 1) [14]. Here, we show how concepts first developed in physics are now increasingly used to describe complex systems in the context of health and disease.

Critical transitions in medicine

Recently the concept of critical transitions and tipping points has been applied to clinical questions and systems medicine. We are convinced that a detailed understanding of critical transitions in disease onset and progression will provide broad applications in health care. The identification of early warning signals (Box 1) for example, can be expected to leverage prevention strategies. Already identified sudden transitions in the medical context have been associated with gut microbiome dysregulation [15^{••}], pulmonary disease [16], depression [17[•]], type 1 and 2 diabetes [18,19], inflammation [20], start [21] and termination [22[•]] of epileptic seizures [23], cancer [24], and cardiovascular events [25]. Examples will be elaborated in more details in the section *Examples of clinical relevance in the context of critical transitions*.

Box 1 Stable state—A stable state of a dynamical (phenotype) system does not change its average phenotypic trait when being exposed to small random perturbations.

Alternative stable states—Distinct stable states of a dynamical system for the same set of environmental conditions. These alternative stable states are separated by a metastable state.

Equilibrium state—The state in which a system stays when no additional external perturbations are applied.

Critical transition—Sudden shift from one stable state to an alternative one where the actual transition can be triggered by small perturbations.

Tipping point—A threshold point at which a system will undergo a critical transition when exposed to perturbations.

Hysteresis—Current system state is depended on both the input and the history of the system.

Nonlinearity—A dynamical system is nonlinear if the set of underlying differential equations exhibit products of variables in at least one equation. Note that systems can obey linear dynamics even if the outcome has a nonlinear form like it is the case of exponential growth.

Early warning signal—An observable variable whose dynamics change considerably before a critical transition occurs.

Emergence—A phenotype that is exhibited at the system level but does not exist when studying the system components individually.

Early warning signals to detect upcoming critical transitions

A critical transition is usually detectable after the transition [26] and difficult to anticipate [27]. Before the critical forward transition, the system's equilibrium state might stay relatively unchanged until the forward tipping point is reached (Figure 2c) [28]. Consequently, static observations might not provide enough information to detect upcoming abrupt transitions [29]. By contrast, changing system dynamics have been suggested as early warning signals (EWS) for critical transitions [30*]. A general challenge for data analysis is the intrinsic noise in biological systems which originates from the stochastic nature of molecular interactions and heterogeneity of individual entities like cells or organisms. A brief explanation for the most commonly used early warning signals can be found in Table 1.

The influence of the random behavior is amplified in the vicinity of tipping points because small perturbations in the vulnerable regime of the system can have large effects (Figure 2). Due to this amplified heterogeneity, an increase in variance [10,31] or coefficient of variation [32] has been associated with upcoming critical transitions. Further, an increase or decrease of lag-1 autocorrelation may indicate the unfolding of an abrupt shift [6,33*]. An increase of flickering activity [34] has been identified as EWS for critical transitions in lake eutrophication [35*,36]. Changing skewness in the distribution of time-series climate data, could be used as a robust indicator for some complex natural systems [37]. Dynamical network biomarkers are a new approach to predict

upcoming transitions and showed promising results for liver cancer [38]. Critical slowing down is found in some ecological systems when approaching a tipping point [39–41,42*]. A transition from vegetation to desertification was preceded by changes in the spatial distribution of vegetative patches [8]. Finally, significant heteroscedasticity [43] was observed one year before a critical transition in a lake [44]. This multitude of early warning signals to detect critical transitions shows that the nonlinearity in different systems are not always accompanied by the same EWS.


Examples of clinical relevance in the context of critical transitions

A recent example of alternative stable states was found in the context of microbiome dysregulation in human intestines. A highly diverse and dynamically evolving microbial ecosystem, mainly including bacteria from the *Firmicutes*, *Actinobacteria* and *Bacteroidetes* phyla, is living in the human gut and its dysregulation can have stark consequences on health [45]. It is thought that certain diseases such as obesity and irritable bowel syndrome might unfold due to transitions in microbial composition [46]. Recently, the question whether these transitions are linear or rather abrupt and nonlinear was raised [47]. Alternative stable states with high resilience were identified in human individuals after repeated prolonged exposure to an antibiotic (Figure 2) [47]. Such alternative stable states of bacterial ecosystems were confirmed in a study covering 1000 individuals [15**].

Acute asthma attacks are characterized by a constriction of the bronchioles which ultimately leads to patchiness in lung ventilation [16] and difficulties to breathe [48]. Such patch clusters can potentially lead to critical transitions via an interaction of feedback mechanisms [16]. One study developed a model of a bronchial tree and simulated incremental airway smooth muscle stimulation [16]. At a stimulation threshold, the system underwent a critical transition and showed severe ventilation defects [48]. In environmental epidemiology, early warning signals in the forms of changing variance and skewness were found for the deterioration of lung activity in humans after exposure to ozone [49*].

Clinical depression is characterized by a wide array of symptoms such as inability to sleep, low mood, loss of interest and suicidal tendencies. Onset and remission of clinical depression can occur suddenly. A recent study suggests that critical slowing down could be an early warning signal for onset and termination of depression [17*]. During the study, subjects were logging their mood states by self-assessment on an emotional scale at random intervals during the day. In follow-up assessments, subjects were re-evaluated using the same scales. Interestingly, existence of critical slowing down based on the collected mood states was confirmed and was indicative of

Figure 1



History of critical transitions. From theory to applications in medicine. Copyrights: Desertification: Google, Map data: DigitalGlobe. All other images were reproduced with written consent or found in the public domain. Hopf bifurcation freely adapted from [5].

upcoming transitions from a normal to a depressed state or vice versa.


Diabetes mellitus has two major sub-types. In type 1 diabetes, patients cannot produce enough insulin and in type 2 diabetes the produced insulin is not used efficiently [50]. Both sub-types can be categorized into multiple disease stages ranging from pre-disease to full onset with clinical diagnosis often occurring only at the latter stage. Dynamical network biomarkers were found to be early warning signals for critical transitions from a pre-disease to a disease state for type 1 [18,51] and type 2 [19] diabetes in mice. On the scale of cellular interactions, critical transitions were identified in insulin-producing pancreatic-cell islets [52]. In this system, a critical amount of β -cell death at which pancreatic islets lose connectivity in the islet network results in systemic failure.

Dynamics of certain pro-inflammatory cytokines play a pivotal role in the unfolding of inflammation [53]. For instance, Interleukin-6 (IL-6) dynamics in pig blood

revealed that variations of this cytokine over time could potentially serve as an early warning signal [20]. Scheff *et al.* showed on a mathematical model that under gradually increasing inflammatory conditions, the state of the system remains stable until a critical threshold at which it evolves from a healthy to an inflammatory state [54]. A mathematical model was created from experimental data where humans underwent an endotoxin application to simulate chronic inflammation. More generally, it has been proposed that early warning signals could predict upcoming epidemic infectious inflammatory disease occurrences in populations [55].

Epileptic seizures start and terminate suddenly and diminish quality of life in patients [56]. Increased variance in spiking patterns of individual neurons has been proposed as an early warning signal to detect the onset of a sudden epilepsy seizure [57]. Further, the mechanism leading to the unfolding of a seizure, when considering groups rather than single neurons, was attributable to a Hopf bifurcation. Recently, Kramer and colleagues

Figure 2



Critical transitions and their characteristics in biomedicine. The general basis of critical transitions is the existence of multistability of the underlying dynamic systems. **(a)** Such a multistability can occur by bifurcations where, for instance, a single stable state is turned into two stable states in dependence on a system parameter during a pitchfork bifurcation. **(b)** Under the assumption of equilibrated systems, the dynamics can be visualized by quasi-energy potentials or stability landscapes where the state of the system describing the phenotype corresponds to a ball in the potential. Changing the condition adiabatically will lead to a modification of the potential and can induce multistability (potentials with grey and black ball) from a single stable point (potentials with only black ball) by imperfect bifurcations. In this regime, the system can perform either induced or spontaneous state transitions by jumping from one well to another. The stability landscapes are projected on the underlying bifurcation diagram. Freely adapted from [7]. **(c)** A typical bifurcation diagram of a bistable system exhibiting hysteresis. The point where a system is switching from one phenotype to another one like in a health-to-disease transition is called the forward tipping point (FTP). The reverse tipping point (RTP) indicates the state where the system can return to the alternative one. For many nonlinear systems the FTP and RTP are not identical and this is known as hysteresis. The area between FTP and RTP corresponds to the regime of coexistence where critical transitions occur which can induce bimodality in populations. These properties may lead to system specific warning signals and emerging properties.

presented evidence of the existence of a critical transition between the ictal and post-ictal states of a seizure, representing alternative stable states [22^{*}]. This critical transition was preceded by critical slowing down, increased autocorrelation and flickering.

Outlook: cross-sectional and longitudinal monitoring of health states

A potential application of using some of the described early warning signals for upcoming critical transitions could be in the generation of a population reference profile and the longitudinal monitoring of healthy people and patients. Similar to the human reference genome, a population reference profile would represent the variation of some of the early warning signals, in time and specific conditions from a representative healthy population. Due to the expected variability between individuals, longitudinal studies are expected to provide clearer signals than studies across populations. Such unique patient reference profiles could, for instance, be obtained during regular visits at the clinic. At each point of sample extraction, the current profile could be compared to

the reference profile. The difference between the profiles would then be indicative of disease progression. At a certain threshold, the clinician could take appropriate measures to prevent a critical transition from a healthy to a disease state in individual patients.

Recently, wearable devices such as Fitbit[®] devices [58], and automated voice recording devices [59,60] have been proposed to be able to provide easy-use monitoring of specific activities in patients. For instance, symptoms such as tiredness, anxiety or speech difficulties, contributing to wearing-off in Parkinson's disease patients are difficult to assess during brief clinic visits [61,62]. Wearing-off means that specific symptoms make a re-appearance before the next scheduled administration of the therapeutic drugs. Wearing-off could become one way to assess critical transitions in Parkinson's disease from a clinical point of view. One could compute variables such as the number of wearing-off occurrences during the day or the time interval between the beginning of wearing-off and the next scheduled drug intake. Consequently, indicators as discussed throughout this review could be

Table 1**Potential indicators for early warning.**

Early warning signal	Definition	Observed system variable
Variance	Scatter of data	Phosphorus concentration in lake [10] Pollutant across multiple regions [31] Interleukin-6 levels [20] Spiking patterns in neurons [57]
Coefficient of variation	Standard deviation normalized by the mean	Population density [32] Spatial heterogeneity of ventilation [16] Infectious population dynamics [55] Ozone levels [49*]
Lag-1 autocorrelation	Correlation of data with itself shifted by one time point	Rate of resource harvesting [33*] Connected population density [32]
Flickering	System states are driven back and forth between alternate stable states by intrinsic noise	Sediment diatom composition [35*] Ice conductivity [34] Phosphorus dynamics [36] Invasive electrocorticogram recordings [22*]
Skewness	Third standardized moment of the distribution of system states	Vegetation biomass [37] Phosphorus density [37]
Dynamical network biomarkers	Evolution over time of difference in molecular networks	Gene expression profiles [18,19,38,51]
Critical slowing down	Recovery rates tend to zero after small external perturbation	Calcium carbonate levels [41] Cyanobacteria population density [42*] Nutrient cycling in lakes [39] Macrophyte cover [39] Vegetation growth [40] Mood dynamics [17*] Vegetation patchiness [8]
Spatial distribution	Non-random distribution of elements in a biological entity	
Conditional heteroscedasticity	Variance is conditional on past time points	<i>E. coli</i> population growth [28] Chlorophyll-a concentration [43]

determined and used to assess the progression of the disease and classify the patient's disease states. Such predictors could allow clinicians to modify treatments accordingly and thereby increase patients' quality of life.

Conclusions

Although a large number of theoretical studies and computational simulations have been carried out to provide evidence for catastrophic shifts, there are still only a few biological and medical studies experimentally or empirically validating the predictions and the relevance of the critical transition concept for medical applications.

Furthermore, error rate estimations for false positive and false negative rates of the models underlying the predictions are rarely available. In order to prevent or revert critical system states this will become more important [63**]. In some disciplines, replicate experiments are often not possible, for example, when analyzing the occurrence of ice-ages, the sudden collapse of fish-populations or catastrophic transitions in the financial markets, as has been witnessed a few years ago. Biological systems and disease pathogenesis however might be studied in depth in animal models, patients with identical or closely related clinical symptoms or adverse responses to specific drug treatments. Due to this feasibility and their importance, biological and biomedical applications might play a

driver role in our attempts to experimentally dissect and understand complex nonlinear natural systems. The development of system specific mathematical models and machine learning tools, integrating a wide range of omics and clinical data, and prior knowledge, for example, from literature, public databases and modern media, will become a central domain in systems medicine.

A note of caution

Critical slowing down might only occur in specific situations. Catastrophic collapse can occur without prior early warning signals in autocorrelation or variance [64**]. Therefore these early warning signals are not always globally applicable and it was suggested that in fact each system might have a characteristic subset of early warning signals [65]. This individuality is based on the intricate interplay of the underlying dynamics which are typically nonlinear and include intrinsic random forces originating, for example, from the stochastic nature of molecular interactions. The existence or occurrence of multiple stable states corresponds to inducible systems many of which exhibit excitable dynamics. Many natural systems exhibit such excitable dynamics like neuron spiking [66] or laser pulsing [67]. Driving such nonlinear dynamics by random perturbations can induce state transitions with a mathematically strictly defined transition rate [68] and lead to non-trivial effects like stochastic and coherence resonance [69,70].

Stochastic resonance describes the scenario where a weak periodic input signal of an excitable nonlinear system is amplified by intrinsic noise. Although first used to describe ice-age periodicity [71] stochastic resonance was subsequently found in a variety of natural and biological systems [72]. In general, stochastic resonance is characterized by a maximum in the signal-to-noise ratio in dependence on the noise intensity and corresponds to a minimum in the coefficient of variation. Similarly, coherence resonance occurs in excitable systems only driven by inherent noise and is also characterized by minimal values of coefficient of variations for optimal random perturbations [70]. These mechanisms can interfere with the general assumptions that critical transitions are accompanied by an increase of variability.

These prominent examples demonstrate that noisy nonlinear systems like those found in biology can obey unintuitive dynamics. The resulting complex behavior including noise induced effects question the general application of the early warning signals mentioned above but emphasize the need to complement experimental investigations with mechanistic theoretical models to fully characterize critical transitions.

Acknowledgments

We would like to thank Linda Wampach for her help with graphical elements in this article. The authors also acknowledge the 'Fonds National de la Recherche Luxembourg (FNR)' for financial support of CT through an AFR grant (3118186). AS received funding from the project 'plan Technologies de la Santé par le Gouvernement du Grand-Duché de Luxembourg' through the Luxembourg Centre for Systems Biomedicine (LCSB), University of Luxembourg. All authors have contributed to the editing of this article.

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Scheffer M: *Critical Transitions in Nature and Society*. Princeton University Press; 2009.
 2. Lewontin RC: **The meaning of stability**. *Brookhaven Symp Biol* 1969, **22**:13-24.
 3. Holling CS: **Resilience and stability of ecological systems**. *Annu Rev Ecol Syst* 1973, **4**:1-23.
 4. May RM: **Thresholds and breakpoints in ecosystems with a multiplicity of stable states**. *Nature* 1977, **269**:471-477.
 5. Strogatz Steven H: *Nonlinear Dynamics And Chaos*. Westview Press; 2001.
 6. Scheffer M, Bascompte J, Brock WA, Brovkin V, Carpenter SR, Dakos V, Held H, van Nes EH, Rietkerk M, Sugihara G: **Early-warning signals for critical transitions**. *Nature* 2009, **461**:53-59.
 7. Scheffer M, Carpenter S, Foley JA, Folke C, Walker B: **Catastrophic shifts in ecosystems**. *Nature* 2001, **413**:591-596.
 8. Kéfi S, Rietkerk M, Alados CL, Pueyo Y, Papanastasis VP, Elaiç A, de Ruiter PC: **Spatial vegetation patterns and imminent desertification in Mediterranean arid ecosystems**. *Nature* 2007, **449**:213-217.
 9. Hirota M, Holmgren M, Van Nes EH, Scheffer M: **Global resilience of tropical forest and savanna to critical transitions**. *Science* 2011, **334**:232-235.
 10. Carpenter SR, Brock WA: **Rising variance: a leading indicator of ecological transition**. *Ecol Lett* 2006, **9**:311-318.
 11. Jackson JB, Kirby MX, Berger WH, Bjorndal KA, Botsford LW, Bourque BJ, Bradbury RH, Cooke R, Erlandson J, Estes JA et al.: **Historical overfishing and the recent collapse of coastal ecosystems**. *Science* 2001, **293**:629-637.
 12. Mumby PJ, Hastings A, Edwards HJ: **Thresholds and the resilience of Caribbean coral reefs**. *Nature* 2007, **450**:98-101.
 13. Lenton TM, Held H, Kriegler E, Hall JW, Lucht W, Rahmstorf S, Schellnhuber HJ: **Tipping elements in the Earth's climate system**. *Proc Natl Acad Sci U S A* 2008, **105**:1786-1793.
 14. Higgins JP: **Nonlinear systems in medicine**. *Yale J Biol Med* 2003, **75**:247-260.
 15. Lahti L, Salojärvi J, Salonen A, Scheffer M, de Vos WM: **Tipping elements in the human intestinal ecosystem**. *Nat Commun* 2014, **5**:1-10.
- In this study to identify nonlinear dynamics in the microbiome of 1000 adults, the authors identified signs of alternative stable states between subgroups of bacterial phyla.
16. Venegas JG, Winkler T, Musch G, Vidal Melo MF, Layfield D, Tgavalekos N, Fischman AJ, Callahan RJ, Bellani G, Harris RS: **Self-organized patchiness in asthma as a prelude to catastrophic shifts**. *Nature* 2005, **434**:777-782.
 17. Van de Leemput I, Wichers M, Cramer AOJ, Borsboom D, Tuerlinckx F, Kuppens P, van Nes EH, Viechtbauer W, Giltay EJ, Aggen SH et al.: **Critical slowing down as early warning for the onset and termination of depression**. *Proc Natl Acad Sci U S A* 2014, **111**:87-92.
- In this medical study, the existence of alternative stable states and associated tipping points was shown using mood information. Interestingly, the authors also provide an attempt to predict the onset and termination of depression based on the mood information.
18. Zeng T, Zhang C, Zhang W, Liu R, Liu J, Chen L: **Deciphering early development of complex diseases by progressive module network**. *Methods* 2014, **67**:334-343.
 19. Li M, Zeng T, Liu R, Chen L: **Detecting tissue-specific early warning signals for complex diseases based on dynamical network biomarkers: study of type 2 diabetes by cross-tissue analysis**. *Brief Bioinform* 2014, **15**:229-243.
 20. Tambuyzer T, De Waele T, Chiers K, Berckmans D, Goddeeris BM, Aerts J-M: **Interleukin-6 dynamics as a basis for an early-warning monitor for sepsis and inflammation in individual pigs**. *Res Vet Sci* 2014, **96**:460-463.
 21. Litt B, Esteller R, Echaüz J, D'Alessandro M, Shor R, Henry T, Pennell P, Epstein C, Bakay R, Dichter M et al.: **Epileptic seizures may begin hours in advance of clinical onset**. *Neuron* 2001, **30**:51-64.
 22. Kramer MA, Truccolo W, Eden UT, Lepage KQ, Hochberg LR, Eskandar EN, Madsen JR, Lee JW, Maheshwari A, Halgren E et al.: **Human seizures self-terminate across spatial scales via a critical transition**. *Proc Natl Acad Sci U S A* 2012, **109**:21116-21121.
- This study has two major outcomes. First it was shown that critical transitions can be studied in a cellular system and second that the termination of epilepsy seizures can be associated to such critical transitions.
23. McSharry PE, Smith La, Tarassenko L: **Prediction of epileptic seizures: are nonlinear methods relevant?** *Nat Med* 2003, **9**:241-242.
 24. Kianercy A, Veltri R, Pienta KJ: **Critical transitions in a game theoretic model of tumour metabolism**. *Interface Focus* 2014, **4**:20140014.
 25. Kowalski WJ, Dur O, Wang Y, Patrick MJ, Tinney JP, Keller BB, Pekkan K: **Critical transitions in early embryonic aortic arch patterning and hemodynamics**. *PLoS One* 2013, **8**:e60271.

26. Boettiger C, Hastings A: **Quantifying limits to detection of early warning for critical transitions.** *J R Soc Interface* 2012, **9**:2527-2539.
 27. Clark JS, Carpenter SR, Barber M, Collins S, Dobson A, Foley JA, Lodge DM, Pascual M, Pielke R, Pizer W *et al.*: **Ecological forecasts: an emerging imperative.** *Science* 2001, **293**:657-660.
 28. Takimoto G: **Early warning signals of demographic regime shifts in invading populations.** *Popul Ecol* 2009, **51**:419-426.
 29. Carpenter SR, Brock WA, Cole JJ, Kitchell ML, Pace JF: **Leading indicators of trophic cascades.** *Ecol Lett* 2008, **11**:128-138.
 30. Dakos V, Carpenter SR, Brock WA, Ellison AM, Guttal V, Ives AR, Kéfi S, Livina V, Seekell DA, van Nes EH *et al.*: **Methods for detecting early warnings of critical transitions in time series illustrated using simulated ecological data.** *PLoS One* 2012, **7**:e41010.
- In this article, the authors provide a practical guide on how to detect EWS in critical transitions from time series data and most importantly made available a toolbox to analyze such data.
31. Brock WA, Carpenter SR: **Variance as a leading indicator of regime shift in ecosystem services.** *Ecol Soc* 2006:11.
 32. Dai L, Korolev KS, Gore J: **Slower recovery in space before collapse of connected populations.** *Nature* 2013, **496**:355-358.
 33. Dakos V, van Nes EH, D'Odorico P, Scheffer M: **Robustness of variance and autocorrelation as indicators of critical slowing down.** *Ecology* 2012, **93**:264-271.
- In this work the authors show that an increase as well as a decrease in variance can be indicative of an upcoming critical transition. Importantly, they show that variance can be under estimated in sparse data sets.
34. Taylor KC, Lamorey GW, Doyle GA, Alley RB, Grootes PM, Mayewski PA, White JWC, Barlow LK: **The flickering switch of late Pleistocene climate change.** *Nature* 1993, **361**:432-436.
 35. Wang R, Dearing JA, Langdon PG, Zhang E, Yang X, Dakos V, Scheffer M: **Flickering gives early warning signals of a critical transition to a eutrophic lake state.** *Nature* 2012, **492**:419-422.
- Wang *et al.* show, using a combination of experimental data and a model, that flickering can occur in an ecological system reflected by bimodality, an increase in variance, and a decrease in autocorrelation and skewness.
36. Dakos V, Nes EH, Scheffer M: **Flickering as an early warning signal.** *Theor Ecol* 2013, **6**:309-317.
 37. Guttal V, Jayaprakash C: **Changing skewness: an early warning signal of regime shifts in ecosystems.** *Ecol Lett* 2008, **11**:450-460.
 38. Wu X, Chen L, Wang X: **Network biomarkers, interaction networks and dynamical network biomarkers in respiratory diseases.** *Clin Transl Med* 2014, **3**:16.
 39. Van Nes EH, Scheffer M: **Slow recovery from perturbations as a generic indicator of a nearby catastrophic shift.** *Am Nat* 2007, **169**:738-747.
 40. Kéfi S, Dakos V, Scheffer M, Van Nes EH, Rietkerk M: **Early warning signals also precede non-catastrophic transitions.** *Oikos* 2013, **122**:641-648.
 41. Dakos V, Scheffer M, van Nes EH, Brovkin V, Petoukhov V, Held H: **Slowing down as an early warning signal for abrupt climate change.** *Proc Natl Acad Sci U S A* 2008, **105**:14308-14312.
 42. Veraart AJ, Faassen EJ, Dakos V, van Nes EH, Lürling M, Scheffer M: **Recovery rates reflect distance to a tipping point in a living system.** *Nature* 2012, **481**:357-359.
- In this fully controllable experimental study, Veraart and colleagues present evidence that critical slowing down occurs before a critical transition in a population of cyanobacteria. The conclusion is that critical slowing down may occur in real complex systems.
43. Seekell DA, Carpenter SR, Cline TJ, Pace ML: **Conditional heteroskedasticity forecasts regime shift in a whole-ecosystem experiment.** *Ecosystems* 2012, **15**:741-747.
 44. Seekell DA, Carpenter SR, Pace ML: **Conditional heteroskedasticity as a leading indicator of ecological regime shifts.** *Am Nat* 2011, **178**:442-451.
 45. De Vos WM, de Vos EAJ: **Role of the intestinal microbiome in health and disease: from correlation to causation.** *Nutr Rev* 2012, **70**(Suppl 1):S45-S56.
 46. Lozupone CA, Stombaugh JI, Gordon JI, Jansson JK, Knight R: **Diversity, stability and resilience of the human gut microbiota.** *Nature* 2012, **489**:220-230.
 47. Dethlefsen L, Relman DA: **Incomplete recovery and individualized responses of the human distal gut microbiota to repeated antibiotic perturbation.** *Proc Natl Acad Sci U S A* 2011, **108**(Suppl):4554-4561.
 48. Winkler T, Venegas JG, Harris RS: **Mathematical modeling of ventilation defects in asthma.** *Drug Discov Today Dis Model* 2014 <http://dx.doi.org/10.1016/j.ddmod.2014.02.008>.
 49. Hsieh N-H, Cheng Y-H, Liao C-M: **Changing variance and skewness as leading indicators for detecting ozone exposure-associated lung function decrement.** *Stoch Environ Res Risk Assess* 2014 <http://dx.doi.org/10.1007/s00477-014-0887-2>.
- For this work, adult subjects were exposed to various levels of ozone. The authors conclude that statistical early warning signals can be used to detect respiratory problems and propose a risk assessment framework to potentially allow for an early identification of the shift from healthy to disease.
50. Polonsky KS: **The past 200 years in diabetes.** *N Engl J Med* 2012, **367**:1332-1340.
 51. Liu X, Liu R, Zhao X-M, Chen L: **Detecting early-warning signals of type 1 diabetes and its leading biomolecular networks by dynamical network biomarkers.** *BMC Med Genomics* 2013, **6**(Suppl 2):S8.
 52. Stamper IJ, Jackson E, Wang X: **Phase transitions in pancreatic islet cellular networks and implications for type-1 diabetes.** *Phys Rev E* 2014, **89**:012719.
 53. Ashley NT, Weil ZM, Nelson RJ: **Inflammation: mechanisms, costs, and natural variation.** *Annu Rev Ecol Evol Syst* 2012, **43**:385-406.
 54. Scheff JD, Calvano SE, Androulakis IP: **Predicting critical transitions in a model of systemic inflammation.** *J Theor Biol* 2013, **338**:9-15.
 55. O'Regan SM, Drake JM: **Theory of early warning signals of disease emergence and leading indicators of elimination.** *Theor Ecol* 2013, **6**:333-357.
 56. Mormann F, Andrzejak RG, Elger CE, Lehnertz K: **Seizure prediction: the long and winding road.** *Brain* 2007, **130**:314-333.
 57. Meisel C, Kuehn C: **Scaling effects and spatio-temporal multilevel dynamics in epileptic seizures.** *PLoS One* 2012, **7**:e30371.
 58. Cook DJ, Thompson JE, Prinsen SK, Dearani JA, Deschamps C: **Functional recovery in the elderly after major surgery: assessment of mobility recovery using wireless technology.** *Ann Thorac Surg* 2013, **96**:1057-1061.
 59. Tsanas A, Little MA, McSharry PE, Ramig LO: **Accurate telemonitoring of Parkinson's disease progression by noninvasive speech tests.** *IEEE Trans Biomed Eng* 2010, **57**:884-893.
 60. Oller DK, Niyogi P, Gray S, Richards Ja, Gilkerson J, Xu D, Yapanel U, Warren SF: **Automated vocal analysis of naturalistic recordings from children with autism, language delay, and typical development.** *Proc Natl Acad Sci U S A* 2010, **107**:13354-13359.
 61. Storch A, Schneider CB, Wolz M, Stürwald Y, Nebe A, Odin P, Mahler A, Fuchs G, Jost WH, Chaudhuri KR *et al.*: **Nonmotor fluctuations in Parkinson disease: severity and correlation with motor complications.** *Neurology* 2013, **80**:800-809.
 62. Stacy M, Bowron A, Guttman M, Hauser R, Hughes K, Larsen JP, LeWitt P, Oertel W, Quinn N, Sethi K *et al.*: **Identification of motor and nonmotor wearing-off in Parkinson's disease: comparison of a patient questionnaire versus a clinician assessment.** *Mov Disord* 2005, **20**:726-733.
 63. Boettiger C, Ross N, Hastings A: **Early warning signals: the charted and uncharted territories.** *Theor Ecol* 2013, **6**:255-264.

This critical review looks at cases where critical transitions are not preceded by specific early warning signals such as critical slowing down and shows published work with contradicting conclusion on the use of EWS. This article highlights the growing observations that there might be no global indicators but rather that certain classes of systems share common dynamics.

64. Boerlijst MC, Oudman T, de Roos AM: **Catastrophic collapse can occur without early warning: examples of silent catastrophes in structured ecological models.** *PLoS One* 2013, **8**:e62033.
 In this original article, Boerlijst shows that critical transitions can occur with no early warning signals. The authors conclude that more theoretical work needs to be performed as the current hypothesis, that generic early warning signals exist in all natural systems, can be refuted.
65. Hastings A, Wysham DB: **Regime shifts in ecological systems can occur with no warning.** *Ecol Lett* 2010, **13**:464-472.
66. Lindner B: **Effects of noise in excitable systems.** *Phys Rep* 2004, **392**:321-424.
67. Giacomelli G, Giudici M, Balle S, Tredicce J: **Experimental evidence of coherence resonance in an optical system.** *Phys Rev Lett* 2000, **84**:3298-3301.
68. Kramers HA: **Brownian motion in a field of force and the diffusion model of chemical reactions.** *Physica* 1940, **7**:284-304.
69. Benzi R, Sutera A, Vulpiani A: **The mechanism of stochastic resonance.** *J Phys A Math Gen* 1981, **14**:L453-L457.
70. Pikovsky A, Kurths J: **Coherence resonance in a noise-driven excitable system.** *Phys Rev Lett* 1997, **78**:775-778.
71. Benzi R, Parisi G, Sutera A, Vulpiani A: **A theory of stochastic resonance in climatic change.** *SIAM J Appl Math* 1983, **43**:565-578.
72. Wiesenfeld K, Moss F: **Stochastic resonance and the benefits of noise: from ice ages to crayfish and SQUIDS.** *Nature* 1995, **373**:33-36.