






Biogeoclimatic Zone

- Boreal Altai Fescue Alpine
- Boreal White and Black Spruce
- Bunchgrass
- Coastal Douglas-fir
- Coastal Mountain-heather Alpine
- Coastal Western Hemlock
- Engelmann Spruce — Subalpine Fir
- Interior Cedar — Hemlock
- Interior Douglas-fir
- Interior Mountain-heather Alpine
- Montane Spruce
- Mountain Hemlock
- Ponderosa Pine
- Spruce — Willow — Birch
- Sub-Boreal Pine — Spruce
- Sub-Boreal Spruce




Fuel Type

- C-1 Spruce-Lichen Woodland
- C-2 Boreal Spruce
- C-3 Mature Jack or Lodgepole Pine
- C-4 Immature Jack or Lodgepole Pine
- C-5 Red and White Pine
- C-7 Ponderosa Pine / Douglas Fir
- M-1/M-2 Boreal Mixedwood (25% Conifer)
- M-1/M-2 Boreal Mixedwood (50% Conifer)
- M-1/M-2 Boreal Mixedwood (75% Conifer)
- M-1 Boreal Mixedwood - Leafless (15% Conifer)
- D-1 Leafless Aspen
- D-1/D-2 Aspen
- O-1a Matted Grass
- Non-fuel
- Vegetated Non-fuel
- Water



Forest Cover Ownership

- Crown
- Crown Lease
- Crown Tenure
- Federal
- Private
- Unknown & Other



Head Fire Intensity (kW/m)

- >0 to 1000
- >1000 to 2000
- >2000 to 4000
- >4000 to 6000
- >6000 to 10000
- >10000 to 18000
- >18000 to 30000
- >30000 to 60000
- >60000 to 100000
- >100000
- Water
- Non-Fuel
- No Data (Private Land)
- No Data (Private Managed Forest Land)