



**Department of Electrical,
Computer, & Biomedical Engineering**
Faculty of Engineering & Architectural Science

Course Title:	
Course Number:	
Semester/Year (e.g.F2016)	

Instructor:	
--------------------	--

<i>Assignment/Lab Number:</i>	
<i>Assignment/Lab Title:</i>	

<i>Submission Date:</i>	
<i>Due Date:</i>	

Student LAST Name	Student FIRST Name	Student Number	Section	Signature*

*By signing above you attest that you have contributed to this written lab report and confirm that all work you have contributed to this lab report is your own work. Any suspicion of copying or plagiarism in this work will result in an investigation of Academic Misconduct and may result in a "0" on the work, an "F" in the course, or possibly more severe penalties, as well as a Disciplinary Notice on your academic record under the Student Code of Academic Conduct, which can be found online at: <http://www.ryerson.ca/senate/current/pol60.pdf>

Problem 1

Part 1

Canonical form of the plane at infinity

$$\pi = (0, 0, 0, 1)^T$$

Part 2

$$\pi'_\infty = H_A^{-T} \pi_\infty = \begin{bmatrix} A^{-T} & 0 \\ -A^+ & 1 \end{bmatrix} \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \end{pmatrix} = \pi_\infty$$

Part 3

A plane as defined by 3 points can be expressed as:

$$\begin{bmatrix} \pi_1^T \\ \pi_2^T \\ \pi_3^T \end{bmatrix} X = 0$$

Undergoing the transform of the 3D points:

$$X' = HX$$

$$\pi' = H^{-T} \pi$$

Problem 2

Part 1

General form:

$$K = \begin{bmatrix} \alpha_x & s & x_0 \\ & \alpha_y & y_0 \\ & & 1 \end{bmatrix}$$

α_x : focal length of x-axis = f_{mx}

α_y : focal length of y axis = f_{my}

s : the skew

x_0, y_0 : principal points.

Part 2

Projection matrix

$$P = [M | P_4]$$

The internal and external parameters can be recovered by

$$P = KR[I] - C = M[I]M^{-1}P_4$$

K : internal parameter

R, t : external parameter

Part 3

The image of the point at infinity can be represented by

$$D = (d^T, 0)^T$$

$$x = PD = [M | P_4] D = M_d$$

Problem 3

Part 1

$$PC = 0$$

or

$$X = \det([p_2, p_3, p_4])$$

$$Z = \det([p_1, p_2, p_4])$$

$$Y = -\det([p_1, p_3, p_4])$$

$$T = -\det([p_1, p_2, p_3])$$

Part 2

$$s_x V_x = p_{xw} = [p_1, p_2, p_3, p_4] \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}^T = p_1$$

$$s_y V_y = p_{yw} = [p_1, p_2, p_3, p_4] \begin{bmatrix} 0 \\ 1 \\ 0 \\ 0 \end{bmatrix}^T = p_2$$

$$s_z V_z = p_{zw} = [p_1, p_2, p_3, p_4] \begin{bmatrix} 0 \\ 0 \\ 1 \\ 0 \end{bmatrix}^T = p_3$$

Part 3

$$\text{Projection plane} = [x, y, 0]$$

$$\begin{bmatrix} x \\ y \\ 0 \end{bmatrix} = \begin{bmatrix} p_1^T \\ p_2^T \\ p_3^T \end{bmatrix} \begin{bmatrix} x \\ y \\ z \\ 1 \end{bmatrix}$$

Problem 4

Part 1

3D Points

$$x = [p_1 \ p_2 \ p_3 \ p_4] \begin{pmatrix} x \\ y \\ 0 \\ 1 \end{pmatrix}$$

2D Points

$$x = [p_1 \ p_2 \ p_4] \begin{pmatrix} x \\ y \\ 1 \end{pmatrix}$$

$$[p_1 \ p_2 \ p_3 \ p_4] \begin{pmatrix} x \\ y \\ 0 \\ 1 \end{pmatrix} = [p_1 \ p_2 \ p_4] \begin{pmatrix} x \\ y \\ 1 \end{pmatrix}$$

$$p = K[R|t]$$

$$H = K[r_1, r_2, t]$$

Part 2

Back projection of a real line in 3D

$$\pi = P^T L$$

x will lie on the line if

$$x^T L = 0$$

real point on the real line in 3D if and only if

$$x^T P^T L = 0$$

therefore

$P^T L$ represents the plane. 2D point x lies on this point if and only if

$$\pi = P^T L$$

Part 3

$$x = K[I|0]X$$

$$x' = K'[I|0]X$$

$$x' = K' K^{-1} (K[I|0]X) = K' K^{-1} x$$

$$x' = Hx$$

$$H = K' K^{-1}$$

Problem 5

Part 1

Real points on the line at infinity:

$$P = KR[I - \tilde{C}]$$

point in the image

$$x = PX_{\infty} = KR[I - \tilde{C}]\begin{pmatrix} 0 \\ 0 \end{pmatrix} = KRd$$

Map observed point from line at infinity $x = Hd$

$$H = KR$$

absolute conic

$$w = (KK^T)^{-1}$$

Part 2

The angle between two rays is

$$\cos(\theta) = \frac{x_1^T w x_2}{\sqrt{x_1^T w x_1} \sqrt{x_2^T w x_2}}$$

if 2 rays are orthogonal

$$x_1^T w x_2 = 0$$

Part 3

$$h_1^T w h_2 = 0$$

$$h_1^T w h_1 = h_2^T w h_2$$

Part 4

The restraints for the square pixel assumption for the image of absolute conic is

$$w_{12} = w_{21} = 0$$

and

$$w_{11} = w_{22}$$