



भा. कृ. अनु. प.- केंद्रीय कपास अनुसंधान संस्थान, नागपुर

ICAR-Central Institute for Cotton Research, Nagpur

An ISO 9001:2015 Certified Organisation



XII Weekly Advisory for Cotton Cultivation from 27th August to 2nd September, 2024

PUNJAB		Actual Rainfall in last week(mm)					Predicted Rainfall in next week (mm)				
		August					Aug / Sep				
		23	24	25	26	27	29	30	31	01	02
	Firozpur						1	8	4	5	1
	Faridkot	0	0	7.6	0.8	0	1	2	4	5	1
	Muktsar						2	1	4	4	1
	Bhatinda	0	0	0	0	26.4	2	2	6	4	2
	Sangrur	0	0	0.5	0	0	2	3	5	2	0
	Ludhiana	1.8	0.2	0	0	15	4	7	5	1	0
Amount of rainfall & colour Code		0.1 to 2.4 mm		2.5 to 15.5 mm		15.6 to 64.4 mm		64.5 to 115.5 mm		115.6 to 204.4	
Rainfall category		Very light rainfall		Light rainfall		Moderate rainfall		Heavy rainfall		Very heavy rainfall	

Crop condition

At Bathinda, the crop is 95 to 115 days old at boll development stage. Intercultural operations, one to two sprays of potassium nitrate (13:0:45) @ 2% in cotton fields where flowering started were taken up during the reporting period. Incidence of whitefly 6-27/three leaves, jassid 0-6/three leaves and around 0-5% flower infestation by pink bollworm was observed. Incidence of leaf curl virus was also recorded in few fields.

At Faridkot, the crop is 101 to 106 days old at boll formation/ boll development stage. Manual intercultural operations and fertiliser application, herbicide and insecticide sprays have been taken up. Leaf hopper and thrips incidence were below ETL and whitefly incidence was moderate to very high and pink bollworm incidence below ETL in most of the fields. Parawilt and CLCuD were also observed a few locations of II-III grade.

Advisory:

At Bathinda, farmers are advised to give four sprays of potassium nitrate (13:0:45) @ 2% at weekly interval in cotton fields to reduce flower drop and to improve boll setting in cotton. Spray the crop with Afidopyropen 50DC @ 400ml/acre or Flonicamid 50WG @ 80g/acre or Clothianidin 50WG @ 20g/acre or Dinotefuran 20%SG @ 60g/acre to control whitefly once infestation crosses ETL. Spray Dinotefuran 20%SG @ 60g/acre or Tolfepryad 15EC @ 400 ml/acre or Fenpyroximate 5EC @ 300 ml/acre against jassid. If thrips population is above ETL, spray Profenofos 50EC @ 600 ml/acre or Spinetoram @ 170ml/acre. Remove and destroy rosette flowers, if any. Spray Profenofos 50EC @ 600 ml/acre or Emamectin benzoate 5SG @ 100g/acre if the incidence of pink bollworm is above 5 per cent (ETL) in flowers or green bolls. Spray Azoxystrobin 18.2%+ Difenoconazole 11.4% SC @ 200 ml per 200 litres of water to manage fungal foliar leaf spots of cotton.

At Faridkot, farmers are advised to spray potassium nitrate (13:0:45) @ 2% to improve boll setting and reduce flower drop in timely sown crop. Avoid brackish/tubewell water for any type of foliar sprays on cotton as excessive salts sometimes burn margins of cotton leaves. Cotton growers who experienced leaf reddening during last season in cotton crop must apply two sprays of Magnesium sulphate @ 1% (through foliar application) at 15 days interval during full bloom and boll development to minimize the issue in Bt cotton. Both chemicals should be sprayed at alternate weeks depending upon weather and rainfall. Drain out excessive water in event of heavy rainfall. To control weeds, spray Glufosinate ammonium 13.5% SL @ 2.5-3.0L/ha as a directed spray between the crop rows. The directed spray can be done by using a protective hood as Glufosinate is non-selective herbicide and can cause injury to the crop if it falls on the crop leaves. Alternatively, do manual or tractor operated intercultural operation to control emerging weeds. In case of high thrips infestation, spray Profenofos 50% EC @ 600ml/acre or Spinetoram @ 170 ml/acre. If whitefly incidence exceeds ETL, spray Flonicamid 50 WG @ 80g/acre or Dinotefuran 20SG @ 60g/acre; in case of higher infestation of whitefly adults, nymphal population and appearance of sooty mould on leaves, spray Afidopyropen 50DC @ 400ml/acre alternately with Pyriproxyfen @ 500ml/acre. Check squares and flowers for the infestation of pink bollworm especially in the early sown crop. Spray Profenofos 50EC @ 600ml/acre or Emamectin benzoate 5SG @ 100g/acre or indoxacarb 14.5SC @ 200ml/acre, if incidence crosses ETL. To monitor pink bollworm, use pheromone trap @ 2 per acre and place it at 30 cm above crop canopy. Replace the lure as per validity. After heavy rains or irrigation, some plants might show wilting due to parawilt which can be checked by spraying Cobalt chloride @ 10 mg/ litre of water on the affected plants at the initial stage of wilting symptom. To manage fungal foliar leaf spot, spray (Azoxystrobin 18.2%+ Difenoconazole 11.4% SC) @ 200 ml per 200 litres of water. To check further spread of cotton leaf curl virus, protect the crop against whitefly vector by using recommended insecticides as given above.

HARYANA		Actual Rainfall in last week(mm)					Predicted Rainfall in next week (mm)				
		August					Aug / Sep				
		23	24	25	26	27	29	30	31	01	02
	Hisar	4.7	0	0	0	12.2	3	2	5	3	2
	Jind	0	1.2	4.2	0	0	3	2	6	2	3
	Sirsa	1.5	0.5	0	0	0	3	2	5	2	3
	Rohtak	0	0	0	0	8	3	4	5	0	0
Amount of rainfall & colour Code		0.1 to 2.4 mm		2.5 to 15.5 mm		15.6 to 64.4 mm		64.5 to 115.5 mm		115.6 to 204.4	
Rainfall category		Very light rainfall		Light rainfall		Moderate rainfall		Heavy rainfall		Very heavy rainfall	

Crop Condition:

At Hisar, the crop is 77 to 126 days at flowering to boll development stage. Flowering is appropriate in areas where rainfall was minimal. Most of the fields are free from weeds but weeds like *motha*, *makra*, *santhi*, *hirankhuri* and *doob* were observed in some of the fields after rainfall. Manual hoeing by *khurpa*/ spade or mechanical hoeing done as per the growth of crop. Population of whitefly is above economic threshold at few locations only where less rainfall was received. Jassid population crossing ETL in few fields and thrips is below ETL in majority of fields. Trap catches of pink bollworm has increased during last week but the infestation of pink bollworm started appearing on flowers and bolls in several fields. Some cases of root rot and wilt were observed. Cotton leaf curl virus disease was also observed in several locations. Sooty mould was also observed in some fields where less rainfall was received. Some cases of cotton leaves malformation were observed due to herbicide contamination.

At Sirsa, the crop is 94 to 119 days at flowering and boll formation stage. Cloudy, rainy and hot humid weather prevailed during the reporting period. Intercultural operations by tractor / bullock, hand hoeing, weeding and application of second split dose of Urea broadcasting and insecticide spray for sucking pests and PBW were taken up. At some locations, tank mixture of insecticides and fungicides spray was observed. Weeds have appeared at all locations. Squaring, flowering and boll formation in full swings at all locations. Whitefly infestation ranged between 11-38/3leaves, thrips below ETL and infestation of pink bollworm reported above ETL (5-20 %) based on green boll damage. Sooty mould and CLCuD incidence were observed at few locations.

Advisory:

At Hisar, farmers are advised to drain excess water after rainfall and apply third split dose of Urea @1bag per acre. Take up manual or mechanical hoeing after irrigation or rainfall. In case of cotton crop more than 100 days old, apply foliar spray of 2.0% Urea+0.5% ZnSO₄ (21%) especially in light soils. Examine at least 150 to 200 flowers and 20 bolls per acre in cotton crop where flowering has started for the infestation of pink bollworm larvae. Install pheromone traps @ 2/acre to monitor pink bollworm. Collect and destroy early season rosette flowers in cotton crop. Manage infestation of pink bollworm by foliar spray of Profenofos 50EC @600ml/acre or Emamectin benzoate 5%SG @100g/acre or Indoxacarb 14.5%SC @200ml/acre or Chlorpyrifos 20%EC @ 500ml/acre. Spray Flonicamid 50 WG @ 80g or Afidopyropen 50 DC @ 400ml/acre for management of sucking pests like whitefly and jassid. Treat root rot affected patches in field by drenching the affected plants with Carbendazim 50WP @2g/litre of water for early symptomatic plants and nearby healthy plants. Confine root rot affected patches by making bunds before flood irrigation so that this disease can be prevented from spreading further. Uproot and bury early season cotton leaf curl virus infected plants. In case of parawilt, spray Cobalt chloride @10mg/ litre of water immediately after the appearance of symptoms on the affected plants. To manage sooty mould, beside spray for whitefly, give a spray of copper oxychloride 50WP @ 2-3 g/litre of water. Monitor fields regularly at weekly intervals and necessarily after rainfall.

At Sirsa, farmers are advised to continue intercultural operations. After irrigation or rain, apply third split dose of nitrogenous fertilizer. For better retention of square, flower and bolls, apply NPK 13:00:45 @ 2kg/100 litres of water and repeat 2-3 times at 10 days interval. If crop is in full swing with square, flowers and bolls and if leaf reddening appears, spray magnesium sulphate @ 1.0 kg in 100 liters of water per acre and repeat after a fortnight. spray 1,250 ml per hectare Paraquat or 2,250 ml per hectare Glufosinate ammonium as a directed spray (using a protective hood) to control weeds in between the crop rows. Use 200- 250 litres of water per acre for spray. Regularly monitor the insect pest infestation. Install pheromone traps @ 2/acre to monitor pink bollworm and 40 low-cost yellow sticky traps for whitefly management. Whitefly adults can be managed by spray of Diafenthiuron 50% WP@ 240 g/acre (field should either be wet due to irrigation or rainfall) or Afidopyropen 50 DC@ 400 ml/acre or Dinotefuran 20SG @ 60g/acre or Flonicamid 50 WG @ 80g or Profenophos 50EC @ 600ml/acre. Spray Spinetoram 11.7%SC @170ml/acre to control thrips. If sooty mould appear, leaves will become sticky or if nymphal population of whitefly is higher, then spray Pyriproxyfen 10 EC @ 400ml or Spiromesifen 22.9 SC @ 240ml/acre 3-5 days after first adult's emergence. Repeat spray of Propiconazole 25EC @1 ml /litre or Mancozeb 30%WDG @ 2.0g/litre or Copper Oxychloride (COC) 50WP@2.5 g/litre water as curative measure. Manage jassid infestation by spraying Dinotefuran 20 SG @ 60g or Flonicamid 50WG @ 80g or Tolfenpyrad 15EC @ 400ml or Fenpyroximate 5% EC @ 300ml/acre. Destroy rosette flower and if PBW infestation crosses ETL based on flower i.e. 10 or more flowers out of 100 observed per acre are infested or 2 out of 20 bolls infested with PBW, or 5-8 male trap catches per night for 3 consecutive nights, then spray spinetoram 11.7 SC @170ml or Profenofos 50EC @ 600ml/acre or

Emamectin benzoate 5SG@ 100 g/acre or Indoxacarb 14.5SC @ 200ml/acre or thiodicarb 75 WP @ 250g in 150 liters of water. In case of appearance of parawilt symptoms, spray cobalt chloride @10mg/litre of water immediately after the appearance of symptoms (within 24 hours) on affected plants only. For boll rot and sooty mould, give prophylactic spray with Carbendazim 50% WP@ 0.4 g or Propineb 70% WP @2.5-3 g/litre or Azoxystrobin 18.2%+ Difenoconazole 11.4% SC@ 1 ml/litre or Propiconazole 25 EC @1 ml/litre or Pyraclostrobin 20% SC @ 1 g/litre or (Fluxapyroxad 167 g/L + Pyraclostrobin 333 g/L SC) @ 0.6 ml/litre or (Metiram 55% + Pyraclostrobin 5% WG) @2 g per litre of water .Avoid spray of tank mixture of insecticides and fungicides. Spray only recommended insecticides or fungicides.

RAJASTHAN		Actual Rainfall in last week(mm)					Predicted Rainfall in next week (mm)				
		August					Aug / Sep				
		23	24	25	26	27	29	30	31	01	02
	Ajmer	3.6	7	79	59.9	16.6	20	20	32	4	9.1
	Jodhpur	0.8	10	4	15.4	0.5	14	8	16	1.3	0
	Nagaur						14	8	26	0.7	2
	Pali	0	0	8	29	26	12	10	10	4	1
	Sri Ganganagar	2	0	0	0	3.3	4.1	4	20	2	0
Amount of rainfall & colour Code		0.1 to 2.4 mm		2.5 to 15.5 mm		15.6 to 64.4 mm		64.5 to 115.5 mm		115.6 to 204.4	
Rainfall category		Very light rainfall		Light rainfall		Moderate rainfall		Heavy rainfall		Very heavy rainfall	

Crop Condition:

In Southern Rajasthan (Banswara, Bhilwara, Chittorgarh, Dungarpur, Pratapgarh, Rajsamand and Udaipur), the crop is 49 to 77 days old at squaring and flowering stage. Intercultural operations for weed management and fertilizer application was done. Fields are free from weeds. Infestation of jassid noticed above ETL and whitefly infestation below ETL. No incidence of diseases reported so far.

At Sriganganagar and Hanumangarh, the crop is 81 to 116 days at square formation, flower initiation and boll development stage. Post sowing irrigation has been applied. Manual hoeing/ hand weeding and intercultural operations are in progress. Jassid population was 0 to 1.8/ 3 leaves, Whitefly 3 to 15.37/3 leaves and thrips population 0 to 9.54/3 leaves were recorded.

Advisory:

In southern Rajasthan (Banswara, Bhilwara, Chittorgarh, Dunarpur, Pratapgarh, Rajsamand and Udaipur), farmers are advised to drain excess rain water from the fields on time. Apply recommended dose of N fertilizers according to crop stage, if required. Keep watch on infestation of sucking pests in early sown cotton. If any sucking pest infestation reported near ETL, spray with Neem based insecticide or NSKE 5%+ Neem oil 5 ml /litre or neem oil-based formulation 5 ml /litre (300 or 1500 ppm) +0.05% surfactant. Install yellow sticky traps 8-10/acre to monitor whitefly and jassid incidence. Spray Flonicamid 50%WG @ 80g/acre or Dinotefuran 20%SG @60g/acre or Imidacloprid 17.8%SL @ 60ml/acre or Tolfenpyrod 15%EC @ 400ml/acre or Fenpyroximate 5%EC @ 300ml/acre against sucking pests when infestation seen above ETL. Install pheromone traps for PBW @ 2 /acre and also watch rosette flowers. Change lures of pheromone traps as per validity. If pink bollworm infestation observed (60-90 DAS), spray Profenofos 50%EC @ 30ml/10L (1500 ml/ha) or Emamectin benzoate 5%SG @ 5g/10L (250 g/ha) or Indoxacarb 14.5%SC @10ml/10L (500ml/ha) or Chlorpyrifos 20 % EC @ 25ml/10L (1250 ml/ha).

At Sriganganagar and Hanumangarh, farmers are advised to give foliar application of potassium nitrate @2% wherever the crop is above 80 days old. Monitor the crop for insect pests and disease regularly. Spray NSKE 5% + Neem formulation @ 5ml /litre or neem oil-based formulation 5 ml /litre (300 or1500 ppm) + 0.05% surfactant to control sucking pests and PBW infestation at initial level of incidence. Spray Flonicamid 50%WG @ 80g/acre Or Dinotefuran 20%SG @ 60g/acre Or Imidacloprid 17.8%SL @ 60ml/acre Or Tolfenpyrod 15%EC @ 400ml/acre Or Fenpyroximate 5%EC @300ml/acre against sucking pests when infestation seen above ETL. In case of higher thrips infestation, spray Spinetoram 11.7SC @170 ml/acre or Profenofos 50 EC @ 600ml/acre. Whenever PBW population cross ETL, spray any of the insecticides like., Emamectin benzoate 5%SG @ 100g/acre or Profenofos 50%EC @ 600ml/acre. The locations found infested with pink bollworm last year are to be monitored closely. Install pheromone traps @ 2 per acre to monitor pink bollworm activity. Foliar spray of Propineb 70% WP @2.5-3 g/litre or Propiconazole 25 EC @1 ml/litre or Carbendazim 50 WP @0.4 g per litre of water is recommended to manage boll rot disease and leaf spots in cotton.

ODISHA		Actual Rainfall in last week(mm)					Predicted Rainfall in next week (mm)				
		August					Aug / Sep				
		23	24	25	26	27	29	30	31	01	02
	Koraput	0	5	2.6	5.5	3	5	6	15	35	45
	Kalahandi	1	5	0	2	0	6	6	10	25	35
	Balangir	52	0.3	18.1	0.7	1.5	5	6	6	20	25
Amount of rainfall & colour Code		0.1 to 2.4 mm		2.5 to 15.5 mm		15.6 to 64.4 mm		64.5 to 115.5 mm		115.6 to 204.4	
Rainfall category		Very light rainfall		Light rainfall		Moderate rainfall		Heavy rainfall		Very heavy rainfall	

Crop Condition:

At Odisha, crop is 49 to 59 days old at square formation and flower initiation stage. Partially cloudy and intermittent rainfall was prevalent during reporting period. Weeding, earthing up, second dose of fertilizers, intercultural operations and spray of neem-based pesticides and application of NAA have been taken up. Weeds of all types i.e., broad leaf, grasses and sedges have emerged in the cotton fields. Incidence of aphids, jassid, Spodoptera, semi loopers and leaf miners reported from some areas. Wilting reported from some places due to continuous rains and water logging in some places.

Advisory:

Farmers are advised to take care to drain excess water from the field. Apply third dose of fertilizer in the early sown crop (after 50-55 DAS) In case of moist soil condition where manual weeding is not possible, go for application of total killer translocated herbicide, Glufosinate Ammonium 13.5% SL @ 2.5-3.0 L/ha as directed spray using hoods for control of all types of weeds. Regularly monitor for any incidence of pests and diseases. If any sucking pest infestation reported near ETL, spray with Neem based insecticide or NSKE 5%+Neem oil 5 ml /litre or neem oil-based formulation 5 ml /litre (300 or 1500 ppm) + 0.05% surfactant. If parawilt or sudden wilt occur due to heavy rain and water logging after dry spell, go for application of cobalt chloride @10mg/litre (10ppm) on affected plants. Spray 100g magnesium sulphate mixed with 100g urea in 10 litres of water in waterlogged areas to reduce leaf reddening in cotton. Apply alfa NAA @3.5 ml/15 litres of water at 60 DAS to prevent square and flower drop. Drench the soil of early symptomatic and nearby plants wilt and root rot diseases with Carbendazim 50 WP@ 2 g/litre of water to manage any soil borne disease including root rot and wilt disease. Foliar spray of Carbendazim 12%+Mancozeb 63% WP@30 g or Propineb 70% WP @2.5-3 g/litre or Propiconazole 25 EC @1 ml/litre or Carbendazim 50 WP @0.4 g per litre of water is recommended to manage boll rot disease and leaf spots in cotton.

GUJARAT		Actual Rainfall in last week(mm)					Predicted Rainfall in next week (mm)				
		August					Aug / Sep				
		23	24	25	26	27	29	30	31	01	02
	Amreli	0	3.6	5	12.8	44.4	215	205	65	30	10
	Bhavnagar	8	16	39.2	31	19.8	200	200	65	15	5
	Jamnagar	0	0	8.1	0	105	225	215	65	20	10
	Rajkot	0	0	40	228	318	235	215	80	35	25
	Junagadh	0	0.8	5.8	6.6	62.8	205	200	65	25	15
	Sabarkantha						70	65	15	19	40
	Surendranagar	0	0	12	137	96.8	205	200	65	15	5
	Ahmedabad	15.4	65	46	84.3	138	125	65	25	15	5
	Baroda	3	21	61.5	91.3	282	125	68	15	18	25
	Patan						70	65	10	5	5
	Mehesana						70	65	15	10	5
Amount of rainfall & colour Code		0.1 to 2.4 mm		2.5 to 15.5 mm		15.6 to 64.4 mm		64.5 to 115.5 mm		115.6 to 204.4	
Rainfall category		Very light rainfall		Light rainfall		Moderate rainfall		Heavy rainfall		Very heavy rainfall	

Crop Condition:

At Surat, sown crop is 56 to 62 days old at vegetative stage. Heavy rains received during the reporting week. Hand weeding and intercultural operations were taken up wherever possible. Weeds have infested the fields. No incidence of diseases. Thrips infestation seen above ETL.

At Junagadh, the sown crop is 35 to 55 days old at vegetative and square formation stage. Heavy rains received during the week. Intercultural, operations, fertilizers and pesticide spray were carried out wherever possible. Fields are infested with weeds. Minor infestations of thrips, jassid and whitefly were noticed in few locations.

Advisory:

At Surat, farmers are advised to drain out the excess rain water from the cotton fields. Keep fields free from debris. To manage sucking pest infestation, spray Neem based insecticide or NSKE 5%+ Neem oil 5ml/litre or neem oil-based formulation 5ml /litre (300 or 1500 ppm) +0.05% surfactant. Install pheromone traps @ 2/acre to monitor pink bollworm. Check *Fusarium* wilt patches along with one-meter radius of healthy plants in desi cotton fields. If parawilt symptoms found in fields, drench the affected plants with Urea (2%) near root zone of plant by manually making 3 to 4 holes for proper aeration and recovery of the plants. Further, to prevent the entry of pathogens, apply Kresoxim methyl 44.3 SC @10 ml or Propineb 70WP @ 25g or Propiconazole 25 EC @10ml or (Metiram 55% + Pyraclostrobin 5%WG) @20g or (Azoxystrobin18.2% w/w + Difenoconazole 11.4% w/w SC) @ 10ml or (Fluxapyroxad167 g/l + Pyraclostrobin 333g/l SC) @ 6 ml mixed in 10 litres of water to manage fungal leaf spots and blight diseases. Foliar spray of Carbendazim 12%+Mancozeb 63% WP@30 g or Propineb 70% WP @2.5-3 g/litre or Propiconazole 25 EC @1 ml/litre or Carbendazim 50 WP @0.4 g per litre of water is recommended to manage boll rot disease complex.

At Junagadh, farmers are advised to drain excess water from the cotton fields. Apply Ammonium sulphate @ 10kg/acre or Urea 15kg/acre. Take up intercultural operations and weeding, earthing up along with application of fertilizers (25kg MOP+25kg Urea/acre in irrigated condition) to the soil. In case of moist soil condition where manual weeding is not possible, go for application of herbicides like Quizalofop ethyl 5%EC @ 2ml /litre of water if the field is infested with grassy weeds or Pyriithobac sodium 10%EC @1.5ml/litre of water for broad leaved weeds or Pyriithobac sodium 6% + Quizalofop Ethyl 4% MEC @2-2.5 ml /litre of water to control both grassy and broad-leaved weeds. If thrips infestation crosses threshold level, spray Thiamethoxam 25%WG @2g/10litres (100g/ha) or Spinetoram 11.7%SC @8.4ml/10 litres (420 ml/ha). Manage fungal foliar diseases, if any, by giving a spray of (Fluxapyroxad 167g/litre + Pyraclostrobin 333g/litre SC) @ 0.6 ml/litre or (Metiram 55%+ Pyraclostrobin 5%WG) @ 2g/litre of water. Foliar spray of Carbendazim 12%+Mancozeb 63% WP@30 g or Propineb 70% WP @2.5-3 g/litre or Propiconazole 25 EC @1 ml/litre or Carbendazim 50 WP @0.4 g per litre of water is recommended to manage boll rot disease complex.

MADHYA PRADESH		Actual Rainfall in last week(mm)					Predicted Rainfall in next week (mm)				
		August					Aug / Sep				
		23	24	25	26	27	29	30	31	01	02
	Khargaon										
	Dhar	11.6	11	6.4	34.3	21.7	11	13	16	22	60
	Khandwa										
Amount of rainfall & colour Code		0.1 to 2.4 mm		2.5 to 15.5 mm		15.6 to 64.4 mm		64.5 to 115.5 mm		115.6 to 204.4	
Rainfall category		Very light rainfall		Light rainfall		Moderate rainfall		Heavy rainfall		Very heavy rainfall	

Crop Condition:

At Khandwa, the sown crop is 56 to 105 days old at vegetative / square / flower / boll formation stage. Spot weeding, fertigation and plant protection measures were done based on the feasibility of field conditions. Irrigation was not given as enough rains have been received. Weeds have dominated the fields. Infestation of jassid and whitefly have been observed in traces in some fields. Occurrence of Bacterial blight, *Corynespora* and *Cercospora* leaf spots have been noticed in few locations. At some of the areas in Dhar, Barwani and Chhindwara districts, sudden drying symptoms have been observed

Advisory:

At Khandwa, farmers are advised to give third dose of chemical fertilizer. Start weeding with bullock drawn *Kolpain* those area where crop is more than 35 days old. In case of moist soil condition where manual weeding is not possible, go for application of herbicides like Quizalofop ethyl 5 %EC @ 2 ml /litre of water if the field is infested with grassy weeds or Pyriithobac sodium 10%EC @ 1.5 ml/lit water for broad leaved weeds or Pyriithobac sodium 6% + Quizalofop Ethyl 4% MEC @ 2-2.5 ml /litre of water to control both grassy and broad-leaved weeds. Install pheromone traps @ 2/acre to monitor pink bollworm infestation and yellow sticky traps @ 8/acre to monitor whitefly. In those crops that has crossed 90 days with sucking pest infestation above ETL, spray Diafenthiuron 50% WP@ 240 g/acre or Dinotefuran 20 SG @ 60g/acre or Flonicamid 50 WG @ 80 g or Profenophos 50 EC @600 ml/acre. Foliar spray of Carbendazim 12%+Mancozeb 63% WP@30 g or Propineb 70% WP @2.5-3 g/litre or Propiconazole 25 EC @1 ml/litre or Carbendazim 50 WP @0.4 g per litre of water is recommended to manage boll rot disease complex and leaf spots in cotton.

MAHARASHTRA		Actual Rainfall in last week(mm)					Predicted Rainfall in next week (mm)				
		August					Aug / Sep				
		23	24	25	26	27	29	30	31	01	02
	Dhule	0.5	6	42	71.5	0	15	23	18	8	10
	Nandurbar	24	2.5	173	89		40	27	18	10	8
	Jalgaon	15.4	8.6	49	11.7	0	8	9	10	15	18
	Ahmednagar	7.6	74	11.4	0	0.5	8	10	12	5	6
	Aurangabad	0.6	4.3	28.7	0	0	4	4	8	9	14
	Jalna	0.5	0	2	1.5	0.5	5	3	5	3	16
	Beed	0	0	0	0.4		3	6	9	5	8
	Nanded	1	48	0.6	0	1.6	5	4	8	15	25
	Parbhani	0	55	2.3	0	0	5	4	5	12	20
	Hingoli	0	0	0	0		5	4	5	12	20
	Buldhana	75	10	12.2	0	0	12	14	29	41.6	43.5
	Akola	48.6	23	25.6	0.5	0	12	13	26	40.2	41.5
	Washim	0	24	13	0	0	12	13	26	42.5	43.6
	Amravati	36	8	26.2	5.8	0	18	19	25	42.5	43.6
	Yavatmal	9	0	59	0	0	13	14	27	42.3	76.5
	Wardha	17	0	18.2	0	1.8	14	18	26	42.5	43.4
	Nagpur	2	1.2	21.4	3.2	4	19	19	27	42.1	43.5
	Chandrapur	26	4	88.8	0	1	19	19	29	75.2	119
Amount of rainfall & colour Code		0.1 to 2.4 mm		2.5 to 15.5 mm		15.6 to 64.4 mm		64.5 to 115.5 mm		115.6 to 204.4	
Rainfall category		Very light rainfall		Light rainfall		Moderate rainfall		Heavy rainfall		Very heavy rainfall	

Crop Condition:

At Akola, sown crop is 42 to 47 days old at square initiation/ formation stage. Different intercultural operations like hoeing, weeding, weedicide and insecticide spraying and top dressing of fertilizers were carried out during reporting week. Some fields are infested with monocots and dicot weeds. Incidence of sucking pests like aphids and jassid noticed in some fields. Infestation of pink bollworm in flower was in the range of 7-8% in early sown cotton fields. No incidence of diseases so far.

At Nanded, the crop is 63 to 83 days at vegetative growth and squaring stage. Weed grasses, broad-leaved weeds and sedges have infested the fields. Intercultural operations, weeding, fertilizer application and plant protection spray were taken up. Incidence of jassid, aphids, thrips and pink bollworm were noticed but below ETL. *Fusarium* wilt was noticed in few patches.

At Rahuri, the crop is at 63 to 86 days at vegetative growth to flowering stage. Intercultural operations, weeding and hoeing were taken up. Weeds have infested the fields. Sucking pest incidence noticed below ETL except thrips which was above ETL. Incidence of TSV at 5-10% was recorded

Advisory:

At Akola, farmers are advised to drain out the excess water from fields in the area where heavy rainfall occurred. For symptoms of parawilt, drench the affected plants with Copper Oxychloride 50 WP@25g or Carbendazim 50 WP@12 g+ Urea 100g in 10 litres of water. Give a spray 2% Urea at flowering stage of cotton with first spray at 45 DAS and 2% spray of DAP at boll development stage of cotton. Spray alpha NAA @ 3 ml /10 litres of water to avoid natural shedding of squares and flowers of cotton. Apply second split of 90 Kg Urea per ha for irrigated hybrid *Bt* cotton. Open the furrow in cotton fields at 50-55 DAS by tying rope to tyne of the hoe to conserve moisture. In case of moist soil condition where manual weeding is not possible, go for application of herbicides like Quizalofop ethyl 5%EC @ 400ml /acre if the field is infested with grassy weeds or Pyriothobac sodium 10% EC @ 300ml/acre for broad leaved weeds or Pyriothobac sodium 6% + Quizalofop Ethyl 4% MEC @ 400-500 ml/acre to control both grassy and broad-leaved weeds. Carry out intercultural operations like hoeing and weeding in cotton fields.

At Nanded, farmers are advised to take up intercultural operations to manage weeds in field condition. Give top-dressing @ 130 kg urea/ha at 60 DAS under irrigated condition and @ 75 kg/ha to rainfed crop. Spray post emergence weedicide, Pyriothobac sodium 6% + Quizalofop ethyl 4% @ 2 – 2.5 litre when weeds are at 2-4 leaf stage to keep crop weed free under labour shortage condition. To prevent sucking pests like aphids, spray NSKE 5% + Neem formulation 5 ml /litre or neem oil-based formulation 5 ml /litre (300 or 1500 ppm) + 0.05% surfactant at 50 DAS. When crop crosses 60 days, spray Flonicamid 50% WG @ 80g/acre or Tolfenpyrod 15% EC @ 400ml/acre to manage jassid when crosses ETL. Install pheromone traps @5/ha to monitor pink bollworm. Collect and destroy rosette flowers. Spray Profenophos 50%EC @ 30ml/10 litre once it crosses ETL i.e. 10% rosette flowers. Drench

with *Trichoderma viridae* 1% WP @ 50g/10 litres water or Carbendazim 50% WP @ 12 g/10 litres water to manage *Fusarium* wilt. Spray Mepiquat chloride @ 500 ml / ha (10 ml / 10 L) as growth retardant under high density planting as well as where excess growth of crop is observed in normal spaced cotton fields. Open furrows at 60 DAS under rainfed condition to conserve rainwater. Do detopping (nipping) at 80-90 DAS.

At Rahuri, farmers are advised to carryout intercultural operations. Install yellow sticky traps @ 8/acre. Spray NSKE 5% + Neem oil 5 ml /litre or neem oil-based formulation 5 ml /litre (300 or 1500 ppm) + 0.05% surfactant to check sucking pests and avoid laying eggs of pink bollworm. Spray Flonicamid 50 WG @ 4g or Buprofezin 25SC @ 20ml or Difenturon 50WP 12 g per 10 liter of water once sucking pests crosses ETL. Install pheromone traps @ 2 per acre to monitor pink bollworm moth activity. Change lure as per validity. Foliar spray of Carbendazim 12%+Mancozeb 63% WP@30 g or Propineb 70% WP @2.5-3 g/litre or Propiconazole 25 EC @1 ml/litre or Carbendazim 50 WP @0.4 g per litre of water is recommended to manage boll rot disease complex and leaf spots in cotton.

TELANGANA		Actual Rainfall in last week(mm)					Predicted Rainfall in next week (mm)				
		August					Aug / Sep				
		23	24	25	26	27	29	30	31	01	02
	Adilabad	0	22	8	0	0	5	12	13	61	27
	Warangal	0	10	8	0	0	11	13	29	28	9
	Khammam	2	0	0	5	0	7	15	14	11	5
	Karimnagar	5.7	24	10.4	4.5	0	7	13	30	35	20
	Mahabubnagar	0	0	4	12	0	9	9	10	25	15
Amount of rainfall & colour Code		0.1 to 2.4 mm		2.5 to 15.5 mm		15.6 to 64.4 mm		64.5 to 115.5 mm		115.6 to 204.4	
Rainfall category		Very light rainfall		Light rainfall		Moderate rainfall		Heavy rainfall		Very heavy rainfall	

Crop Condition:

At Warangal, the sown crop is 57 to 67 days at vegetative to square formation stage. Foliar application of NPK nutrients, pesticide sprays against pests and diseases were done. Infestation of thrips was above ETL, jassids, aphids and mealy bug infestation was noticed below ETL.

At Adilabad, the sown crop is 61 to 66 days old at vegetative stage. Foliar application of nutrients, pest management, manual weeding and intercultural operations have been completed. Infestation of aphids was noticed above ETL but thrips and jassid were below ETL. PBW Infestation has just started in the form of rosette flowers

Advisory:

At Warangal, farmers are advised to give foliar application of 19-19-19/13-0-45 @10g/litre of water. If any sucking pest infestation reported near ETL, spray with Neem based insecticide or NSKE 5%+ Neem oil 5 ml /litre or neem oil-based formulation 5 ml /litre (300 or 1500 ppm) + 0.05% surfactant. Install yellow sticky traps @ 8/acre to monitor whitefly and jassid infestation. In case of moist soil condition where manual weeding is not possible, go for application of herbicides like Quizalofop ethyl 5 % EC @ 2 ml /litre of water if the field is infested with grassy weeds or Pyriothobac sodium 10 % EC @ 1.5 ml/litre of water for broad leaved weeds or Pyriothobac sodium 6% + Quizalofop Ethyl 4% MEC @ 2-2.5 ml/litre of water to control both grassy and broad-leaved weeds. Spray Flonicamid 50%WG @ 80g/acre or Dinotefuran 20%SG @ 60g/acre or Imidacloprid 17.8%SL @ 60ml/acre or Tolfenpyrod 15%EC @ 400ml/acre or Fenpyroximate 5%EC @ 300ml/acre against sucking pests when infestation seen above ETL. Foliar spray of Carbendazim 12%+Mancozeb 63% WP@30 g or Propineb 70% WP @2.5-3 g/litre or Propiconazole 25 EC @1 ml/litre or Carbendazim 50 WP @0.4 g per litre of water is recommended to manage boll rot disease complex and leaf spots in cotton.

At Adilabad, farmers are advised to give foliar application of nutrients 19-19-19/13-0-45 @10g/lit of water. Install yellow and blue sticky traps @ 8/ acre to monitor thrips. Do not spray any insecticides to control early stage sucking pests which might kill entire natural enemy population and delay the crop maturity. Instead, spray NSKE 5% + Neem formulation 5ml /litre or neem oil-based formulation 5ml /litre (300 or 1500 ppm) + 0.05% surfactant at 50 DAS. Spray Flonicamid 50%WG @ 80g/acre or Dinotefuran 20%SG @ 60g/acre or Imidacloprid 17.8%SL @ 60ml/acre or Tolfenpyrod 15%EC @ 400ml/acre or Fenpyroximate 5%EC @ 300ml/acre against sucking pests if infestation seen above ETL. Install pheromone traps @ 2 per acre to monitor pink bollworm moth activity, change lure as per validity. Collect and destroy rosette flowers. Spray Profenofos 50%EC @ 30ml/10L (1500 ml/ha) Or Emamectin benzoate 5%SG @ 5g/10L (250 g/ha) to manage pink bollworm. Foliar spray of Carbendazim 12%+Mancozeb 63% WP@30 g or Propineb 70% WP @2.5-3 g/litre or Propiconazole 25 EC @1 ml/litre or Carbendazim 50 WP @0.4 g per litre of water is recommended to manage boll rot disease complex and leaf spots in cotton.

ANDHRA PRADESH		Actual Rainfall in last week(mm)					Predicted Rainfall in next week (mm)				
		August					Aug / Sep				
		23	24	25	26	27	29	30	31	01	02
	Guntur	0.1	0.1	0	0.1	0	2	1	8	20	12
	Prakasam	2.3	0.6	0	0	0	1	1	8	12	8
Amount of rainfall & colour Code		0.1 to 2.4 mm		2.5 to 15.5 mm		15.6 to 64.4 mm		64.5 to 115.5 mm		115.6 to 204.4	
Rainfall category		Very light rainfall		Light rainfall		Moderate rainfall		Heavy rainfall		Very heavy rainfall	

Crop Condition:

At Guntur, sown crop is 35 days old at vegetative stage. Sowing is still in progress. Infestation of thrips and leaf hoppers noticed above ETL. Pheromone traps @4/acre was erected where the crop is at 40 days old.

At Nandyal, the crop is 60 to 94 days old at flowering and boll formation stage. Grassy weeds have dominated the fields. Weeding, top dressing of fertilizer and insecticide sprays were given. Infestation of thrips and jassids noticed above ETL.

Advisory:

At Guntur, farmers who are yet to take up sowing are advised to treat seeds with imidacloprid 600 FS@9 ml/kg seed and *Pseudomonas fluorescens* @10g/kg or *Trichoderma viride*@8 g/Kg seed or Carboxin 37.5% + Thiram37.5% DS @3.5 g per kg of seeds (root rot and bacteria diseases) or Fluxapyroxad (333 g/L FS) @1.5 ml per kg seed or Tetraconazole 11.6% W/W (12.5% w/v) SL @1.5 ml per kg of seeds to manage seed and soil borne diseases. Give pre-emergence application of Pendimethalin 30 EC @ 1.5 litre/acre within 48 hours of sowing cotton and basal application of P₂O₅ @ 60 kg/acre and 50 kg N and 20 Kg K₂O per acre within 30 DAS. In the field where crop is already sown which is in vegetative stage, manage sucking pests by spraying NSKE 5%+ Neem oil 5 ml /litre or neem oil-based formulation 5 ml /litre (300 or 1500 ppm) + 0.05% surfactant or if needed, spray flonicamid 50 WG @ 4 g per 10 litres of water.

At Nandyal, farmers are advised to give a spray of Urea 20g/litre of water or 19-19-19 @5g/litre of water to tide over moisture stress condition. In case of moist soil condition where manual weeding is not possible, go for application of herbicides like Quizalofop ethyl 5% EC @ 2 ml /litre of water if the field is infested with grassy weeds or Pyriproxyfen sodium 10% EC @1.5 ml/litre of water for broad leaved weeds or Pyriproxyfen sodium 6% + Quizalofop Ethyl 4% MEC @2-2.5 ml /litre of water to control both grassy and broad-leaved weeds. Spray neem-based pesticide 1 L /acre, install yellow and blue sticky traps@ 8/acre along to against sucking pests and pheromone traps @2/acre to monitor pink bollworm incidence. Change lure as per validity. As a prophylactic measure, give a spray of Carbendazim 12%+Mancozeb 63% WP@30 g or Propineb 70% WP @2.5-3 g/litre or Propiconazole 25 EC @1 ml/litre or Carbendazim 50 WP @0.4 g per litre of water may be taken is recommended to manage boll rot disease complex, leafspots due to humidity and congenial weather factors. To prevent sucking pests like aphids, spray NSKE 5% + Neem formulation 5 ml /litre or neem oil-based formulation 5 ml /litre (300 or 1500 ppm) + 0.05% surfactant. In case of high sucking pest infestation, spray Flonicamid 50%WG @ 80g/acre or Dinotefuran 20%SG @ 60g/acre or Imidacloprid 17.8%SL @ 60ml/acre or Buprofezin 25SC @ 400ml/acre. Spray Profenofos @ 400ml and Chlorpyrifos @ 500ml/acre to manage pink bollworm in cotton.

KARNATAKA		Actual Rainfall in last week(mm)					Predicted Rainfall in next week (mm)				
		August					Aug / Sep				
		23	24	25	26	27	29	30	31	01	02
	Dharwad	3.2	8.4	17.4	5.4		2	2	4	4	4
	Haveri						2	3	4	4	4
	Mysore	0	0	0	2	0	2	3	5	5	4
Amount of rainfall & colour Code		0.1 to 2.4 mm		2.5 to 15.5 mm		15.6 to 64.4 mm		64.5 to 115.5 mm		115.6 to 204.4	
Rainfall category		Very light rainfall		Light rainfall		Moderate rainfall		Heavy rainfall		Very heavy rainfall	

Crop Condition:

At Dharwad, the sown crop is 74 to 79 days old at flowering to square formation stage. Sporadic rainfall received in surrounding locations. Weeds like *Cyanodondactylon* and *Cyperusrotundus* have dominated the fields. Hand weeding, intercultural operations and spraying of nutrients and insecticides were taken up. Sucking pests like thrips and aphids, shoot weevil infestations were noticed in few pockets. No incidence of diseases.

At Raichur, the early sown crop is 70 to 76 days old at flowering stage, late sown 48 to 53 days old at squaring stage and very late sown 33 to 38 days old at vegetative stage. Due to continuous rains, the cotton fields are infested with more weeds. Thrips and jassid infestations were noticed at a few locations.

At Chamarajanagar, the crop is 93 to 108 days old at flowering to boll formation stage. Top-dressing of urea, weeding, intercultivation and earthing-up operations were done. *Cyprus* spp. Weeds have dominated the fields. Infestation of leafhopper (4-5/3 leaves), Aphids (20-30/3 leaves) and Mirid bug (6-7/25 squares) was noticed.

Advisory:

At Dharwad, farmers are advised to take up hand weeding and intercultivation to manage weeds in the fields wherever cotton crop is sown. Spray Flonicamid 50 WG @ 80g/acre or Buprofezin 25SC @ 400ml/acre or Diafenthiuron 50WP @ 240g/acre once sucking pests' crosses ETL. Install pheromone traps @ 2 per acre to monitor pink bollworm moth activity, change lure as per validity. Spray Profenofos 50%EC @ 30ml/10L (1500 ml/ha) Or Emamectin benzoate 5%SG @ 5g/10L (250 g/ha) of water to manage pink bollworm if infestation crosses ETL. Foliar spray of Carbendazim 12%+Mancozeb 63% WP@30 g or Propineb 70% WP @2.5-3 g/litre or Propiconazole 25 EC @1 ml/litre or Carbendazim 50 WP @0.4 g per litre of water is recommended to manage boll rot disease complex and leaf spots if noticed in the cotton fields.

At Raichur, farmers are advised to take up weeding on time. Give recommended dose of fertilizers (DAP, MOP and MgSO₄) for the very late sown and early sown crop as per the stipulated crop duration period. Spray post emergent application of Pyriothobac Sodium herbicide @ 1.5 ml/litre of water for 25-30 days old crop. Manage leafhoppers by spraying Flonicamid @0.4g or Thiamethoxam @ 0.2g in 1 litre of water. In case of sucking pests above ETL, spray Flonicamid 50%WG @ 80g/acre or Dinotefuran 20%SG @ 60g/acre or Imidacloprid 17.8%SL @ 60ml/acre or Buprofezin 25SC @ 400ml/acre. Spray Profenofos @ 400ml and Chlorpyrifos @ 500ml/acre to manage pink bollworm in cotton. For early sown crop, install pheromone traps @4/acre to monitor pink boll worm population. To early sown crop, give first foliar spray of 19:19:19 (10 grams in 1 litre of water) + MgSO₄ (10 grams in 1 litre of water) for better growth, boll retention and to manage leaf reddening problem. Spray Dinotefuran @0.4 g/litre of water against leafhoppers and Profenophos @ 3 ml in 1 litre of water as a precautionary spray against pink boll worm in 60 to 70 days old crop. Foliar spray of Carbendazim 12%+Mancozeb 63% WP@30 g or Propineb 70% WP @2.5-3 g/litre or Propiconazole 25 EC @1 ml/litre or Carbendazim 50 WP @0.4 g per litre of water is recommended to manage boll rot disease complex and leaf spots in cotton.

At Chamarajanagar, farmers are advised to take up hand weeding and intercultivation operations to control weeds. In case of sucking pests above ETL, spray Flonicamid 50%WG @ 80g/acre or Dinotefuran 20%SG @ 60g/acre or Imidacloprid 17.8%SL @ 60ml/acre or Buprofezin 25SC @ 400ml/acre. Install pheromone traps @ 8/ acre to monitor pink bollworm attack. Spray Profenofos @ 400ml and Chlorpyrifos @ 500ml/acre to manage pink bollworm in cotton once it crosses ETL. Foliar spray of Carbendazim 12%+Mancozeb 63% WP@30 g or Propineb 70% WP @2.5-3 g/litre or Propiconazole 25 EC @1 ml/litre or Carbendazim 50 WP @0.4 g per litre of water is recommended to manage boll rot disease complex and leaf spots in cotton.

TAMIL NADU		Actual Rainfall in last week(mm)					Predicted Rainfall in next week (mm)				
		August					Aug / Sep				
		23	24	25	26	27	29	30	31	01	02
	Perambalur	0	0	0			0	7	1	2	0
	Salem	0	0	0	0	0	3	8	4	10	5
	Trichy						0	3	0	3	2
	Virudhunagar						0	2	0	0	0
Amount of rainfall & colour Code		0.1 to 2.4 mm		2.5 to 15.5 mm		15.6 to 64.4 mm		64.5 to 115.5 mm		115.6 to 204.4	
Rainfall category		Very light rainfall		Light rainfall		Moderate rainfall		Heavy rainfall		Very heavy rainfall	

Crop Condition:

At Coimbatore and surrounding areas, the crop is around 7 to 35 days old at seedling to initial vegetative growth stage. Sowing is still in progress. Gap filling, thinning, hand weeding were done. Weeds like *Cyperusrotundus* and *Parthenium* have spread in the fields. Collar rot was noticed in few fields.

At Srivilliputhur in and around fields, sowing is yet to commenced. Final land preparation is in progress.

Advisory:

At Coimbatore and surrounding areas, farmers are advised to clean bunds, irrigation and drainage channels. Take up gap filling 10 DAS and thinning of seedlings on 15 days old crop. Do hand weeding on more than 25 days old crop depending on the weed intensity. Monitor the crop scrupulously for pest and disease incidence. Do spot drenching with Carbendazim 50 WP@12 g/10 litres of water at the base of affected plants and surrounding healthy plants to manage collar rot incidence. Drench the soil with *Bacillus subtilis* @ 10ml/litre of water to manage soil borne diseases.

At Srivilliputhur in and around fields, farmers are advised to use the showers for field preparation and sowing and drain the excess water. Remove and destroy previous cotton crop stubbles to avoid stem weevil and root rot infestation. Trim the bunds and bring the field in good tilth condition by using rotavator. Form ridges and furrows at 75 cm and 45 cm spacing for irrigated and rainfed cotton, respectively. Follow crop rotation where heavy infestation of stem weevil and bollworm complex are prominent. During final plough, apply neem cake @250 kg/ha to prevent stem weevil infestation. Clean the farm bunds and other premises to destroy the alternate host of insect pests such as *Abutilon* and other grassy weeds

--/--