



भा. कृ. अनु. प.- केंद्रीय कपास अनुसंधान संस्थान, नागपुर
ICAR-Central Institute for Cotton Research, Nagpur
 An ISO 9001:2015 Certified Organisation



XIV Weekly Advisory for Cotton Cultivation from 10th to 16th September, 2024

PUNJAB		Actual Rainfall in last week(mm)					Predicted Rainfall in next week (mm)				
		September					September				
		06	07	08	09	10	12	13	14	15	16
	Firozpur						0	0	0	0	0
	Faridkot	0	0	0	0		0	0	0	0	0
	Muktsar						0	0	0	0	0
	Bhatinda	0	2.4	0	0		0	0	0	0	0
	Sangrur	0	0	0	0	0	0	0.5	0.1	0	0
	Ludhiana	0	25	0	3.2	0	0	0	0	0	0
Amount of rainfall & colour Code		0.1 to 2.4 mm		2.5 to 15.5 mm		15.6 to 64.4 mm		64.5 to 115.5 mm		115.6 to 204.4	
Rainfall category		Very light rainfall		Light rainfall		Moderate rainfall		Heavy rainfall		Very heavy rainfall	

Crop condition

At Bathinda, the crop is 109 to 129 days old at boll development stage. Foliar spray of Potassium nitrate (13:0:45) @ 2% in cotton fields where flowering started and insecticide and fungal sprays were taken up during the reporting period. Incidence of 0-5% in green bolls and 5% flower infestation by pink bollworm was observed. Incidence of leaf curl virus and wilting symptoms was also recorded in few fields.

At Faridkot, the crop is 115 to 120 days old at peak flowering/ boll formation/ boll development stage. Intercultural operations, insecticide and foliar sprays were taken up. Jassid and thrips incidence were below ETL and whitefly incidence was moderate to very high and pink bollworm incidence was near ETL in most of the fields. Incidence of CLCuD was also observed at most of the locations in II-III grade.

Advisory:

At Bathinda, farmers are advised to give four sprays of Potassium nitrate (13:0:45) @ 2% at weekly interval in cotton fields to reduce flower drop and to improve boll setting in cotton. Spray the crop with Afidopyropen 50 DC @ 400 ml/acre or Flonicamid 50 WG @ 80 g/acre or Dinotefuran 20% SG @ 60 g/acre to control whitefly once infestation crosses ETL. If jassid population is above ETL, spray Dinotefuran 20% SG @ 60 g/acre or Tolfepryad 15 EC @ 400 ml/acre or Fenpyroximate 5 EC @ 300 ml/acre. If thrips population is above ETL, spray Profenofos 50EC @ 600 ml/acre or Spinetoram @ 170 ml/acre. Remove and destroy rosette flowers, if any. Spray Profenofos 50 EC @ 600 ml/acre or Emamectin benzoate 5 SG @ 100 g/acre if the incidence of pink bollworm is above 5 per cent (ETL) in flowers or green bolls. Spray Azoxystrobin 18.2%+ Difenconazole 11.4% SC @ 200 ml per 200 litres of water to manage fungal foliar leaf spots of cotton. Apply two sprays of Magnesium Sulphate @ 1% at 15 days interval if leaf reddening appeared.

At Faridkot, farmers are advised to spray Potassium nitrate (13:0:45) @ 2% to improve boll setting and reduce flower drop at 7 to 10 days interval in timely sown crop. Avoid brackish/tube well water for any type of foliar sprays on cotton as excessive salts sometimes burn margins of cotton leaves. Cotton growers who experienced leaf reddening during last season in cotton crop must apply two sprays of Magnesium sulphate @1% (through foliar application) at 15 days interval during full bloom and boll development to minimize the issue in Bt cotton. Both chemicals should be sprayed at alternate weeks depending upon weather and rainfall. In case of high thrips infestation, spray Profenofos 50% EC @ 600 ml/acre or Spinetoram @ 170 ml/acre. If whitefly incidence exceeds ETL, spray Flonicamid 50WG @80g/acre or Dinotefuran 20SG @ 60g/acre; in case of higher infestation of whitefly adults, nymphal population and appearance of sooty mould on leaves, spray Afidopyropen 50 DC @ 400ml/acre alternately with Pyriproxyfen @ 500 ml/acre. Check squares and flowers for the infestation of pink bollworm especially in the early sown crop. Spray Profenofos 50EC @600ml/acre or Emamectin benzoate 5SG @ 100g/acre or Indoxacarb 14.5SC @ 200 ml/acre, if incidence crosses ETL. To monitor pink bollworm, use pheromone trap @ 2 per acre and place it at 30 cm above crop canopy. Replace the lure as per validity. After heavy rains or irrigation, some plants might show wilting due to parawilt which can be checked by spraying Cobalt chloride @ 10 mg/ litre of water on the affected plants at the initial stage of wilting symptom. To manage fungal foliar leaf spot, spray (Azoxystrobin 18.2%+ Difenconazole 11.4% SC) @ 200 ml/ 200 lit of water. To check further spread of cotton leaf curl virus, protect the crop against whitefly vector by using recommended insecticides as given above.

HARYANA		Actual Rainfall in last week(mm)					Predicted Rainfall in next week (mm)				
		September					September				
		06	07	08	09	10	12	13	14	15	16
	Hisar	19.2	0	0.5	0		0	0	0	0	0
	Jind	0	0	0	0	0	0.3	5.4	1.5	0	0
	Sirsa	0	0	0	0		0	0.3	0	0	0
	Rohtak	0	4	0.6	2		1.2	12	1.1	0	0
Amount of rainfall & colour Code											
Rainfall category		Very light rainfall		Light rainfall		Moderate rainfall		Heavy rainfall		Very heavy rainfall	

At Hisar, the crop is 91 to 140 days at flowering to boll development stage. The crop condition is deteriorating due to continuous rainfall and problems like, lodging, weed infestation, boll rot, jassid infestation, wilting of plants have started to appear. Crop is in good condition and bearing good fruiting bodies in areas where rainfall was minimal. Most of the fields are free from weeds but weeds like *motha*, *makra*, *santhi*, *hirankhuri* and *doob* were observed in some of the fields after rainfall managed through weedicide spray. Population of whitefly is above economic threshold at few locations only where less rainfall was received. However, jassid incidence crossed ETL in large number of fields. Trap catches of pink bollworm has increased during last week and infestation has started appearing on flowers and bolls in several fields crossing ETL. Some cases of root rot and wilt were observed. Cotton leaf curl virus disease was observed in several locations. Boll rot was also observed in some fields where high rainfall was received.

At Sirsa, the crop is 110 days at flowering and boll formation stage. Cloudy, rainy and hot humid weather prevailed during the reporting period. Intercultural operations by tractor / bullock, hand hoeing, weeding and application of final split dose of Urea broadcasting and insecticide spray for sucking pests, pink bollworm and sooty mould were taken up. At some locations, tank mixture of insecticides and fungicides spray was observed. Some farmers sprayed micronutrient i.e. NPK, Boron, Zn etc. mixed with insecticide in tank. Weeds have appeared at all locations. Squaring, flowering and boll formation in full swings at all locations. Whitefly infestation ranged between 18-51/3leaves, jassid 0-12/3 leaves, thrips below ETL and infestation of pink bollworm reported above ETL (5-17 %) based on green boll damage. Sooty mould (very less) and CLCuD incidences were observed at few locations.

At Hisar, farmers are advised to drain excess water after rainfall and keep fields free from weeds as per the feasibility. Give two foliar sprays of 13:00:45 @ 2kg per acre in cotton crop bearing good fruiting after 90-100 days of sowing at an interval of 10 days. In case of cotton crop is more than 100 days old, give alternate foliar spray of 2.5% Urea+0.5% ZnSo₄ (21%) especially in light soils. Examine at least 150 to 200 flowers and 20 bolls per acre in cotton crop where flowering has started for the infestation of pink bollworm larvae. Install pheromone traps @ 2/acre to monitor pink bollworm. Collect and destroy early season rosette flowers in cotton crop. Spray Profenofos 50 EC @ 600 ml/acre or Eamectin benzoate 5 SG @ 100 g/acre if the incidence of pink bollworm is above 5 per cent (ETL) in flowers or green bolls. Spray Flonicamid 50 WG @ 80 g or Afidopyropen 50 DC @ 400 ml/acre for management of sucking pests like whitefly and jassid. Do not use synthetic pyrethroids in cotton fields where whitefly is crossing threshold level till 15th September. Treat root rot affected patches in field by drenching the affected plants with Carbendazim 50 WP @2 g/litre of water for early symptomatic plants and nearby healthy plants. Confine root rot affected patches by making bunds before flood irrigation so that this disease can be prevented from spreading further. Uproot and bury early season cotton leaf curl virus infected plants. In case of parawilt, spray Cobalt chloride @10 mg/ litre of water immediately after the appearance of symptoms (within 24 hrs.) on the affected plants. Monitor fields regularly at weekly intervals and necessarily after rainfall.

At Sirsa, farmers are advised to continue intercultural operations. After irrigation or rain, apply third split dose of nitrogenous fertilizer. For better retention of square, flower and bolls, apply NPK 13:00:45 @ 2 kg/100 litres of water and repeat 2-3 times at 10 days interval. If crop is in full swing with square, flowers and bolls and if leaf reddening appears, spray Magnesium sulphate @ 1.0 kg in 100 liters of water per acre and repeat after a fortnight during full bloom and boll development stages. Use 200- 250 litres of water per acre for spray. Regularly monitor the insect pest infestation. Install pheromone traps @ 2/acre to monitor pink bollworm and 40 low-cost yellow sticky traps for whitefly management. Control whitefly adults by spraying Diafenthiuron 50% WP @ 240 g/acre (field should either be wet due to irrigation or rainfall) or Afidopyropen 50 DC @ 400 ml/acre or Dinotefuran 20 SG @ 60 g/acre or Flonicamid 50 WG @ 80 g or Profenophos 50 EC @ 600 ml/acre. If sooty mold appear, leaves will become sticky or if nymphal population of whitefly is higher, then spray Pyriproxyfen 10 EC @ 400 ml or Spiromesifen 22.9 SC @ 240 ml/acre 3-5 days after first adult's emergence. Repeat spray of Propiconazole 25 EC @ 1 ml /litre or Mancozeb 30% WDG @ 2.0 g/litre or Copper Oxychloride (COC) 50 WP @ 2.5 g/litre water as curative measure. Manage jassid infestation by spraying Dinotefuran 20 SG @ 60 g or Flonicamid 50 WG @ 80 g or Tolfenpyrad 15 EC @ 400 ml or Fenpyroximate 5% EC @ 300 ml/acre. Destroy rosette flower

and if pink bollworm infestation crosses ETL based on flower i.e. 10 or more flowers out of 100 observed per acre are infested or 2 out of 20 bolls infested with pink bollworm, or 5-8 male trap catches per night for 3 consecutive nights, then spray Profenofos 50 EC @ 600 ml/acre or Emamectin benzoate 5 SG @ 100 g/acre or Indoxacarb 14.5 SC @ 200 ml/acre or thiodicarb 75 WP @ 250 g in 150 liters of water. In case of appearance of parawilt symptoms, spray cobalt chloride @ 10 mg/litre of water immediately after the appearance of symptoms (within 24 hours) on affected plants only. For boll rot and sooty mould, give prophylactic spray with Carbendazim 50% WP @ 0.4 g or Propineb 70% WP @ 2.5-3 g/litre or (Azoxystrobin 18.2% + Difenoconazole 11.4% SC) @ 1 ml/litre or Propiconazole 25 EC @ 1 ml/litre or Pyraclostrobin 20% SC @ 1 g/litre or (Fluxapyroxad 167 g/L + Pyraclostrobin 333 g/L SC) @ 0.6 ml/litre or (Metiram 55% + Pyraclostrobin 5% WG) @ 2 g per litre of water. Avoid spray of tank mixture of insecticides and fungicides. Spray only recommended insecticides or fungicides.

RAJASTHAN		Actual Rainfall in last week(mm)					Predicted Rainfall in next week (mm)				
		September					September				
		06	07	08	09	10	12	13	14	15	16
	Ajmer	20.1	28	8.6	6.2		2.5	0	0.2	0	0
	Jodhpur	0.1	0.9	9	0		0	0	0	0	0
	Nagaur						0.4	0	0	0	0
	Pali	0	0	0			0.6	0.1	0.1	0	0
	Sri Ganganagar	0	0	0	0		0	0	0	0	0
Amount of rainfall & colour Code		0.1 to 2.4 mm		2.5 to 15.5 mm		15.6 to 64.4 mm		64.5 to 115.5 mm		115.6 to 204.4	
Rainfall category		Very light rainfall		Light rainfall		Moderate rainfall		Heavy rainfall		Very heavy rainfall	

Crop Condition:

In Southern Rajasthan (Banswara, Bhilwara, Chittorgarh, Dungarpur, Pratapgarh, Rajsamand and Udaipur), the crop is 63 to 91 days old at squaring and flowering stage. Intercultural operations for weed management and fertilizer application was done. Fields are free from weeds. Infestation of jassid and whitefly infestation noticed below ETL. No incidence of diseases reported so far.

At Sriganganagar and Hanumangarh, the crop is 95 to 130 days at square formation, flower initiation and boll development stage. Post sowing irrigation has been applied. Manual hoeing/ hand weeding and intercultural operations are in progress. Jassid population was 0.0-2.0/ 3 leaves, Whitefly 2.0-8.2/3 leaves and thrips population 0.0 to 3.22/3 leaves were recorded. Occurrence of CLCuD up to grade IV to V has been noticed

Advisory:

In southern Rajasthan (Banswara, Bhilwara, Chittorgarh, Dunarpur, Pratapgarh, Rajsamand and Udaipur), farmers are advised to drain excess rain water from the fields on time due to continuous heavy rains. Apply recommended dose of N fertilizers according to crop stage, if required. Keep watch on infestation of sucking pests in early sown cotton. If any sucking pest infestation reported near ETL, spray with Neem based insecticide or NSKE 5%+ Neem oil 5 ml /litre or neem oil-based formulation 5 ml /litre (300 or 1500 ppm) + 0.05% surfactant. Install yellow sticky traps 8-10/acre to monitor whitefly and jassid incidence. Spray Flonicamid 50% WG @ 80 g/acre or Dinotefuran 20% SG @ 60 g/acre or Imidacloprid 17.8% SL @ 60 ml/acre or Tolfenpyrod 15% EC @ 400 ml/acre or Fenpyroximate 5% EC @ 300 ml/acre against sucking pests when infestation seen above ETL. Install pheromone traps for PINK BOLLWORM @ 2 /acre. Change lures of pheromone traps as per validity. Remove and destroy rosette flowers, if any. Spray Profenofos 50 EC @ 600 ml/acre or Emamectin benzoate 5 SG @ 100 g/acre if the incidence of pink bollworm is above 5 per cent (ETL) in flowers or green bolls. Do not repeat same insecticides as well same group of insecticides. Avoid tank mixture of two or more insecticides.

At Sriganganagar and Hanumangarh, farmers are advised to give foliar application of potassium nitrate @ 2% wherever the crop is above 80 days old. Monitor the crop for insect pests and disease regularly. Spray NSKE 5% + Neem formulation @ 5 ml /litre or neem oil-based formulation 5 ml /litre (300 or 1500 ppm) + 0.05% surfactant to control sucking pests and pink bollworm infestation at initial level of incidence. Spray Flonicamid 50% WG @ 80 g/acre Or Dinotefuran 20% SG @ 60 g/acre Or Tolfenpyrod 15% EC @ 400 ml/acre Or Fenpyroximate 5% EC @ 300 ml/acre against sucking pests when infestation seen above ETL. In case of higher thrips infestation, spray Spinetoram 11.7 SC @ 170 ml/acre or Profenofos 50 EC @ 600 ml/acre. Whenever pink bollworm population cross ETL, spray any of the insecticides like Emamectin benzoate 5% SG @ 100 g/acre or Profenofos 50% EC @ 600 ml/acre. The locations found infested with pink bollworm last year are to be monitored closely. Install pheromone traps @ 2 per acre to monitor pink bollworm activity. Foliar spray of Propineb 70% WP @ 2.5-3 g/litre or Propiconazole 25 EC @ 1 ml/litre or Carbendazim 50 WP @ 1.0 g per litre of water is recommended to manage boll rot disease and fungal leaf spots in cotton.

ODISHA	Actual Rainfall in last week(mm)					Predicted Rainfall in next week (mm)				
	September					September				

		06	07	08	09	10	12	13	14	15	16
	Koraput	0	0	0	0		2	13	14	2.1	6.3
	Kalahandi	17	17	6	24		1.4	39	16	25.4	13.7
	Balangir	0.5	4.7	2.6	39.9		6.3	40	20	19.6	18.2
Amount of rainfall & colour Code		0.1 to 2.4 mm		2.5 to 15.5 mm		15.6 to 64.4 mm		64.5 to 115.5 mm		115.6 to 204.4	
Rainfall category		Very light rainfall		Light rainfall		Moderate rainfall		Heavy rainfall		Very heavy rainfall	

Crop Condition:

At Odisha, crop is 63 to 69 days old at flowering and boll formation stage. Partially cloudy with intermittent rainfall was prevalent during reporting period. Weeding, third dose of fertilizers, intercultural operations, pesticide and fungal sprays have been taken up. Weed infestation seen in all the cotton fields. Incidence of aphids, jassid, Spodoptera, semiloopers and leaf miners reported from some areas. Wilting reported from Balangir and Nuapada districts due to continuous rains and water logging conditions. Root rot, leaf spot and Bacterial leaf blight reported from Kalahandi and Rayagada districts.

Advisory:

Farmers are advised to drain excess water from the field. Apply third dose of fertilizer by 60 DAS to reduce more vegetative growth. In case of moist soil condition where manual weeding is not possible, go for application of herbicides like Quizalofop ethyl 5 % EC @ 2 ml /litre of water if the field is infested with grassy weeds or Pyriothibac sodium 10%EC @ 1.5 ml/lit water for broad leaved weeds or Pyriothibac sodium 6% + Quizalofop Ethyl 4% MEC @ 2-2.5 ml /litre of water to control both grassy and broad-leaved weeds. In case of water logging conditions, ensure proper drainage in the fields to drain out excess rainwater by providing adequate slope and draining excessive stagnating water around the root system at the earliest possible. Due to waterlogged situation, cotton crop suffers from inadequacy of nutrients causing reddening and chlorosis in leaves due to poor uptake by roots under anerobic conditions in soil. Give foliar application of 1.0 % Urea or DAP or 19:19:19 at 15 days interval. Regularly monitor for any incidence of pests and diseases. If any sucking pest infestation reported near ETL, spray with Neem based insecticide or NSKE 5%+Neem oil 5 ml /litre or neem oil-based formulation 5 ml /litre (300 or 1500 ppm) + 0.05% surfactant. If parawilt or sudden wilt occur due to heavy rain and water logging after dry spell, drenching to be done for affected plants/rows with a mixture of Copper oxychloride 25 g + 200 g Urea in 10 liters of water immediately after occurrence of Parawilt within 48 h can revive the plants. Spray 100g Magnesium sulphate mixed with 100g Urea in 10 litres of water in waterlogged areas to reduce leaf reddening in cotton. Apply alfa NAA @4 ml/10 lit of water at 60 DAS to prevent square and flower drop. Drench the soil of early symptomatic and nearby plants wilt and root rot diseases with Carbendazim 50 WP@ 2 g/litre of water to manage any soil borne disease including root rot and wilt disease. Give foliar spray of (Carbendazim 12%+Mancozeb 63% WP) @30g or Propineb 70%WP @2.5-3g/lit or Propiconazole 25EC @1ml/lit or Carbendazim 50WP @0.4g/lit of water to manage boll rot disease and leaf spots in cotton.

GUJARAT		Actual Rainfall in last week(mm)					Predicted Rainfall in next week (mm)				
		September					September				
		06	07	08	09	10	12	13	14	15	16
	Amreli	31.8	0	0	2		0.2	0.6	0.6	1.3	0.5
	Bhavnagar	8.2	23	0	0		0.2	0.4	3.6	5.5	1
	Jamnagar	0	0	0	0		0	0	0	0	0
	Rajkot	28.4	0	0	0		0	0	0	0	0
	Junagadh	1.6	2.5	1.4	10		0	0	0	0	0
	Sabarkantha						0	0	0	0	0
	Surendranagar	0	0.4	0	0		0	0	0	0	0
	Ahmedabad	25.1	0.6	0	0		0	0	0	0	0
	Baroda	9.6	7.2	0.3	0		0	0.6	1.2	2.7	2.8
	Patan						0.1	0.4	0.5	0.4	0.2
	Mehesana						0	0	0	0	0
Amount of rainfall & colour Code		0.1 to 2.4 mm		2.5 to 15.5 mm		15.6 to 64.4 mm		64.5 to 115.5 mm		115.6 to 204.4	
Rainfall category		Very light rainfall		Light rainfall		Moderate rainfall		Heavy rainfall		Very heavy rainfall	

Crop Condition:

At Surat, sown crop is 70 to 76 days old at vegetative and square formation stage. The weather was cloudy and rainy. Hand weeding and intercultural operations were taken up wherever possible. Weeds have infested the fields. No incidence of diseases. Thrips infestation seen above ETL.

At Junagadh, the sown crop is at vegetative and square formation stage. Intercultural, operations, fertilizers and pesticide spray

could not be done. Fields are infested with weeds. Minor infestations of thrips, jassid and whitefly were noticed in few locations. Parawilt is the major issue at majority of farmers' fields.

Advisory:

At Surat, farmers are advised to drain out the excess rain water from the cotton fields. Keep fields free from debris. On receiving sufficient rain, apply second split dose of Nitrogenous fertilizer. To manage sucking pest infestation, spray Neem based insecticide or NSKE 5%+ Neem oil 5 ml/litre or neem oil-based formulation 5 ml /litre (300 or 1500 ppm) +0.05% surfactant. Install pheromone traps @ 2/acre to monitor pink bollworm. Check *Fusarium* wilt patches along with one-meter radius of healthy plants in desi cotton fields. If parawilt symptoms found in fields, drenching to be done for affected plants/rows with a mixture of Copper oxychloride 25 g + 200 g Urea in 10 liters of water immediately after occurrence of Parawilt within 48 h can revive the plants. Further, to prevent the entry of pathogens, apply Kresoxim methyl 44.3 SC @10 ml or Propineb 70 WP @ 25 g or Propiconazole 25 EC @10ml or (Metiram 55% + Pyraclostrobin 5% WG) @20 g or (Azoxystrobin 18.2% w/w + Difenconazole 11.4% w/w SC) @ 10 ml or (Fluxapyroxad 167 g/l + Pyraclostrobin 333 g/l SC) @ 6 ml mixed in 10 litres of water to manage fungal leaf spots and blight diseases. Foliar spray of (Carbendazim 12%+Mancozeb 63% WP) @30 g or Propineb 70% WP @2.5-3 g/litre or Propiconazole 25 EC @1 ml/litre or Carbendazim 50 WP @0.4 g per litre of water is recommended to manage boll rot disease complex.

At Junagadh, farmers are advised to drain excess water from the cotton fields. Apply Ammonium sulphate @ 10 kg/acre or Urea 15 kg/acre. Take up intercultural operations, weeding along with application of fertilizers (25 kg MOP+25 kg Urea/acre in irrigated condition) to the soil. In case of moist soil condition where manual weeding is not possible, go for application of herbicides like Quizalofop ethyl 5% EC @ 2 ml /litre of water if the field is infested with grassy weeds or Pyriothobac sodium 10% EC @1.5 ml/litre of water for broad leaved weeds or Pyriothobac sodium 6% + Quizalofop Ethyl 4% MEC @2-2.5 ml /litre of water to control both grassy and broad-leaved weeds. If thrips infestation crosses threshold level, spray Thiamethoxam 25% WG @ 2 g/10 litres (100 g/ha) or Spinetoram 11.7% SC @ 8.4 ml/10 litres (420 ml/ha). Apply spray of (Fluxapyroxad 167 g/litre + Pyraclostrobin 333 g/litre SC) @ 0.6 ml/litre or (Metiram 55%+ Pyraclostrobin 5% WG) @ 2 g/litre of water against fungal foliar diseases. Foliar spray of (Carbendazim 12%+Mancozeb 63% WP) @30 g or Propineb 70% WP @2.5-3 g/litre or Propiconazole 25 EC @1 ml/litre or Carbendazim 50 WP @0.4 g per litre of water is recommended to manage boll rot disease complex. In case of water logging conditions, ensure proper drainage in the fields to drain out excess rainwater by providing adequate slope and draining excessive stagnating water around the root system at the earliest possible. Due to waterlogged situation cotton crop suffers from inadequacy of nutrients causes reddening and chlorosis in leaves due to poor uptake by roots under anerobic conditions in soil. Foliar application of 1.0 % Urea or DAP or 19:19:19 at 15 days interval is recommended

MADHYA PRADESH		Actual Rainfall in last week(mm)					Predicted Rainfall in next week (mm)				
		September					September				
		06	07	08	09	10	12	13	14	15	16
	Khargaon										
	Dhar	0	16	0.3	0.2		2.5	0.2	0.7	0.3	0.1
	Khandwa										
Amount of rainfall & colour Code		0.1 to 2.4 mm		2.5 to 15.5 mm		15.6 to 64.4 mm		64.5 to 115.5 mm		115.6 to 204.4	
Rainfall category		Very light rainfall		Light rainfall		Moderate rainfall		Heavy rainfall		Very heavy rainfall	

Crop Condition:

At Khandwa, the sown crop is 70 to 118 days old at square / flower / boll formation stage. Spot weeding, fertigation and plant protection measures were done based on the feasibility of field conditions. Irrigation was not given as enough rains have been received. Weeds have dominated the fields. Infestation of jassid and whitefly have been observed in some fields. Occurrence of Bacterial blight, *Corynespora* and *Cercospora* leaf spots have been noticed in few locations. In few areas of Dhar, Barwani and Chhindwara districts, sudden drying symptoms have been observed.

Advisory:

At Khandwa, farmers are advised to give third and fourth dose of chemical fertilizer as per requirement. Start weeding with bullock drawn *Kolpa*. In case of moist soil condition where manual weeding is not possible, go for application of herbicides like Quizalofop ethyl 5 % EC @ 2 ml /litre of water if the field is infested with grassy weeds or Pyriothobac sodium 10%EC @ 1.5 ml/lit water for broad leaved weeds or Pyriothobac sodium 6% + Quizalofop Ethyl 4% MEC @ 2-2.5 ml /litre of water to control both grassy and broad-leaved weeds. Install pheromone traps @ 2/acre to monitor pink bollworm infestation and yellow sticky traps @ 8/acre to monitor whitefly. In those crops that has crossed 90 days with sucking pest infestation above ETL, spray Diafenthiuron 50% WP @ 240 g/acre or Dinotefuran 20 SG @ 60 g/acre or Flonicamid 50 WG @ 80 g/acre. Foliar spray of (Carbendazim 12%+Mancozeb 63% WP) @30 g or Propineb 70% WP @2.5-3 g/litre or Propiconazole 25 EC @1 ml/litre or Carbendazim 50 WP @0.4 g per litre of

water is recommended to manage boll rot disease complex and leaf spots in cotton. There is excessive growth of cotton and less fruiting in some fields. Spray these fields with Mepiquat Chloride 5% @ 200g/acre Or 1 ml per litre of water. If parawilt symptoms found in fields, drenching to be done for affected plants/rows with a mixture of Copper oxychloride 25 g + 200 g Urea in 10 liters of water immediately after occurrence of Parawilt within 48 h can revive the plants.

MAHARASHTRA		Actual Rainfall in last week(mm)					Predicted Rainfall in next week (mm)				
		September					September				
		06	07	08	09	10	12	13	14	15	16
	Dhule	0	0	0	0	0.5	2.1	1	1.5	0.2	0.2
	Nandurbar	25	32	17.5	0	9	1.1	1.2	2.1	0.7	0.9
	Jalgaon	0	11	0	0		5.4	1.2	1.4	0.9	0.2
	Ahmednagar	0	0	2.5	0	0	0.3	0.6	0.2	0.2	0.2
	Aurangabad	1.1	0.6	15.2	0.3	0	2.8	1.1	1.5	1.8	1.2
	Jalna						1.4	0.6	0.4	1.3	0.1
	Beed	0	5	0			0.4	1.3	0.6	0.8	0.9
	Nanded	1	4	6.5	0	0	1.7	0.6	0.4	1.7	0
	Parbhani	8	2.8	0.3	0.6		0.5	0.2	0	1.5	0
	Hingoli		0	0	0	0.5	1	0.4	0.1	2	0
	Buldhana	21.3	22	0	0		6.1	0.6	0.4	1.3	0
	Akola	0	2.9	4.2	0		6.8	0.1	0	0.4	0
	Washim	24	10	0	0		6.5	0.4	0.1	1	0
	Amravati	1	6	2	0		9.4	2.4	0.3	0.6	1.4
	Yavatmal	0	0	6.3	0		5.8	0.2	0.1	0.5	0.8
	Wardha	0	0	0	2		7.4	2.2	0.1	0.1	3.8
	Nagpur	0	0	6.6	0		14	12	6.4	0.2	6.2
	Chandrapur	0	0	15	4		5.6	6.5	1.2	0.2	4.9
Amount of rainfall & colour Code		0.1 to 2.4 mm		2.5 to 15.5 mm		15.6 to 64.4 mm		64.5 to 115.5 mm		115.6 to 204.4	
Rainfall category		Very light rainfall		Light rainfall		Moderate rainfall		Heavy rainfall		Very heavy rainfall	

Crop Condition:

At Akola, sown crop is at various stages depending upon the time sown. The crop is at vegetative to boll development stage. The weather during the reporting period was mostly cloudy with less sunshine hours, and heavy rainfall was received during this week. Different intercultural operations like hoeing and insecticide spray were carried out during reporting week. Some fields are infested with monocots and dicot weeds. Incidence of sucking pests like thrips and jassid noticed in some fields. Infestation of spotted and pink bollworm in flower were observed. No incidence of diseases so far.

At Nanded, the crop is 77 to 97 days at flowering/ boll formation stage. Weed grasses, broad-leaved weeds and sedges have infested the fields. Intercultural operations and plant protection spray were taken up. Incidence of jassid, thrips and pink bollworm were noticed but below ETL. Parawilt was noticed in few patches.

At Rahuri, the crop is at 77 to 100 days at squaring to flowering stage. Intercultural operations, weeding and hoeing were taken up. Weeds have infested the fields. Sucking pest incidence noticed below ETL. Incidence of TSV at 8-15% was recorded

Advisory:

At Akola, farmers are advised to drain out the excess water from fields in the area where heavy rainfall occurred. Dead furrow should be opened in June and July sown crop if not opened. Take up weeding to avoid competition of weeds with crop. Give a spray of 2% Urea at flowering stage and 2% spray of 13-0-45 at boll development stage of cotton. Spray NAA @ 4 ml /10 lit of water to avoid natural shedding of squares and flowers of cotton and mepiquat chloride @ 10 ml/10 lit of water to restrict the excess vegetative growth of cotton. To manage sucking pests of cotton above ETL, spray Diafenthiuron 50% WP @ 240 g/acre or Dinotefuran 20 SG @ 60 g/acre or Flonicamid 50 WG @ 80 g/acre. Install 2 pheromone traps/acre for pink bollworm monitoring. Spray Profenofos 50% EC @ 30 ml/10 L (1500 ml/ha) Or Emamectin benzoate 5% SG @ 5 g/10 L (250 g/ha) to manage pink bollworm.

At Nanded, farmers are advised to drain excess water by opening trenches. Take up intercultural operations to manage weeds in field condition. Spray post emergence weedicide, Pyriithiobac sodium 6% + Quizalofop ethyl 4% MEC @ 2 – 2.5 litre when weeds are at 2-4 leaf stage to keep crop weed free under labour shortage condition. Spray NAA @ 4ml /10 lit of water to avoid natural

shedding of squares and flowers of cotton. To prevent sucking pests like aphids, spray NSKE 5% + Neem formulation 5 ml /litre or neem oil-based formulation 5 ml /litre (300 or 1500 ppm) + 0.05% surfactant. Spray Flonicamid 50% WG @ 80 g/acre or Tolfe nypyr od 15% EC @ 400 ml/acre to manage jassid whencrosses ETL. Install pheromone traps @ 5/ha to monitor pink bollworm. Collect and destroy rosette flowers. Spray Profenophos 50% EC @ 30 ml/10 lit once it crosses ETL i.e. 10% rosette flowers. Spray Mepiquat chloride @ 500 ml / ha (10 ml / 10 lit) as growth retardant under high density planting as well as where excess growth of crop is observed in normal spaced cotton fields. Open furrows at 60 DAS under rainfed condition to conserve rainwater. Do detopping (nipping) at 80-90 DAS under irrigated condition and at 75 DAS in rainfed HDPS cotton fields. Drench the affected plants with Copper oxychloride 50 WP @ 25 g + Urea 200g/10 lit water to manage parawilt in cotton fields.

At Rahuri, farmers are advised to carryout intercultural operations. Install yellow sticky traps @ 8/acre. Spray NSKE 5% + Neem oil 5 ml /litre or neem oil-based formulation 5 ml /litre (300 or 1500 ppm) + 0.05% surfactant to check sucking pests and avoid laying eggs of pink bollworm. Spray Flonicamid 50 WG @ 4 g or Buprofezin 25 SC @ 20 ml or Difenthrun 50 WP 12 g per 10 liter of water once sucking pests crosses ETL. Install pheromone traps @ 2 per acre to monitor pink bollworm moth activity. Change lure as per validity. Foliar spray of Carbendazim 12%+Mancozeb 63% WP@30 g or Propineb 70% WP @2.5-3 g/litre or Propiconazole 25 EC @1 ml/litre or Carbendazim 50 WP @0.4 g per litre of water is recommended to manage boll rot disease complex and leaf spots in cotton. In case of water logging conditions, ensure proper drainage in the fields to drain out excess rainwater by providing adequate slope and draining excessive stagnating water around the root system at the earliest possible. Due to waterlogged situation, cotton crop suffers from inadequacy of nutrients causing reddening and chlorosis in leaves due to poor uptake by roots under anerobic conditions in soil. Foliar application of 1.0 % Urea or 13-0-45 or 19:19:19 at 15 days interval is recommended. Spray NAA @ 4ml /10 lit of water to avoid natural shedding of squares and flowers of cotton. If parawilt or sudden wilt occur due to heavy rain and water logging after dry spell, drenching to be done for affected plants/rows with a mixture of Copper oxychloride 50WP @ 25 g + 200 g Urea in 10 liters of water immediately after occurrence of Parawilt within 48 hrs to revive the plants.

TELANGANA		Actual Rainfall in last week(mm)					Predicted Rainfall in next week (mm)				
		September					September				
		06	07	08	09	10	12	13	14	15	16
	Adilabad	0	3	15	0		4	0.5	0.4	0.5	0.2
	Warangal	0	22	2	18		5.4	0.1	0.6	0	0
	Khammam	0	19	60	12		4.8	0.1	1	0	0
	Karimnagar	0	0.5	0.2	5.5		0.1	0	0.2	0.1	0
	Mahabubnagar	38	0.4	0	2		0	0.1	0.3	0	0
Amount of rainfall & colour Code		0.1 to 2.4 mm		2.5 to 15.5 mm		15.6 to 64.4 mm		64.5 to 115.5 mm		115.6 to 204.4	
Rainfall category		Very light rainfall		Light rainfall		Moderate rainfall		Heavy rainfall		Very heavy rainfall	

Crop Condition:

At Warangal, the sown crop is 71 to 81 days at square formation / flowering and boll development stage. Foliar application of NPK nutrients, pesticide sprays against pests and diseases were done. Infestation of thrips was noticed above ETL, jassid and aphids below ETL and TSV (5-10%). Parawilt was noticed due to water logged conditions in the field.

At Adilabad, the sown crop is 75 to 80 days old at vegetative stage. Foliar application of nutrients, pest management, manual weeding and intercultural operations have been completed. Infestation of aphids was noticed below ETL but thrips and jassid were above ETL. Pink bollworm Infestation has just started in the form of rosette flowers.

Advisory:

At Warangal, farmers are advised to give foliar application of 19-19-19 or 13-0-45 @10 g/lit of water at crop under dry or inundation/excess soil moisture conditions. In case of moist soil condition where manual weeding is not possible, go for application of herbicides like Quizalofop ethyl 5 % EC @ 2 ml /lit of water if the field is infested with grassy weeds or Pyri thio bac sodium 10 % EC @ 1.5 ml/litre of water for broad leaved weeds or Pyri thio bac sodium 6% + Quizalofop Ethyl 4% MEC @ 2-2.5 ml/litre of water to control both grassy and broad-leaved weeds. Spray Magnesium Sulphate @10g/ lit of water two times at 10 days interval for Magnesium deficiency and spray Boran @ 2-3g/lit of water to arrest the flower and boll drop problem. If any sucking pest infestation reported near ETL, spray with Neem based insecticide or NSKE 5%+ Neem oil 5 ml /lit or neem oil-based formulation 5 ml /lit (300 or 1500 ppm) + 0.05% surfactant. Install yellow sticky traps @ 8/acre to monitor whitefly and jassid Infestation. Spray Flonicamid 50% WG @ 80 g/acre or Dinotefuran 20% SG @ 60 g/acre or Tolfe nypyr od 15% EC @ 400 ml/acre or Fenpyroximate 5% EC @ 300 ml/acre against sucking pests when infestation seen above ETL. Install pheromone traps @ 2/acre to monitor pink bollworm. Spray Profenofos 50% EC @ 30 ml/10 L (1500 ml/ha) Or Emamectin benzoate 5% SG @ 5 g/10 L (250 g/ha) to manage pink bollworm. Collect and destroy rosette flowers to arrest the spread of pink bollworm. Give foliar spray of (Carbendazim 12%+Mancozeb 63% WP) @30g or Propineb 70% WP @2.5-3 g/lit or Propiconazole 25 EC @1 ml/lit or

Carbendazim 50 WP @0.4g/lit of water to manage boll rot disease complex and leaf spots in cotton.

At Adilabad, farmers are advised to give foliar application of nutrients 19-19-19 or 13-0-45 @10 g/lit of water. In case of water logging conditions, ensure proper drainage in the fields to drain out excess rainwater by providing adequate slope and draining excessive stagnating water around the root system at the earliest possible. Due to waterlogged situation, cotton crop suffers from inadequacy of nutrients causing reddening and chlorosis in leaves due to poor uptake by roots under anaerobic conditions in soil. Foliar application of 1.0 % Urea or DAP or 19:19:19 at 15 days interval is recommended. Install yellow and blue sticky traps @ 8/ acre to monitor thrips. Do not spray any insecticides to control early stage sucking pests which might kill entire natural enemy population and delay the crop maturity. Instead, spray NSKE 5% + Neem formulation 5 ml /litre or neem oil-based formulation 5 ml /litre (300 or 1500 ppm) + 0.05% surfactant at 50 DAS. Spray Flonicamid 50% WG @ 80 g/acre or Dinotefuran 20% SG @ 60 g/acre or Tolfe pyrod 15% EC @ 400 ml/acre or Fenpyroximate 5%EC @ 300 ml/acre if infestation of sucking pests seen above ETL. Install pheromone traps @ 2 per acre to monitor pink bollworm moth activity, change lure as per validity. Collect and destroy rosette flowers. Spray Profenofos 50% EC @ 30 ml/10 L (1500 ml/ha) Or Emamectin benzoate 5% SG @ 5 g/10 L (250 g/ha) to manage pink bollworm. Give foliar spray of Carbendazim 12%+Mancozeb 63% WP@30 g or Propineb 70% WP @2.5-3 g/litre or Propiconazole 25 EC @1 ml/litre or Carbendazim 50 WP @0.4g/lit of water to manage boll rot disease complex and leaf spots in cotton. If parawilt or sudden wilt occur due to heavy rain and water logging after dry spell, drench the affected plants/rows with a mixture of Copper oxychloride 50WP @ 25 g + 200 g Urea in 10 lit of water immediately after occurrence of parawilt within 48hrs to revive the plants.

ANDHRA PRADESH		Actual Rainfall in last week(mm)					Predicted Rainfall in next week (mm)				
		September					September				
		06	07	08	09	10	12	13	14	15	16
	Guntur	53	3.8				0	0	1.7	0	0
	Prakasam	3.3	0	0	0		0	0	0.1	0.1	0.8
Amount of rainfall & colour Code		0.1 to 2.4 mm		2.5 to 15.5 mm		15.6 to 64.4 mm		64.5 to 115.5 mm		115.6 to 204.4	
Rainfall category		Very light rainfall		Light rainfall		Moderate rainfall		Heavy rainfall		Very heavy rainfall	

Crop Condition:

At Guntur, sown crop is 49 to 67 days old at vegetative to square formation stage. Due to heavy rains field were completely inundated with water. Water was drained out from the fields immediately after receding the rains.

At Nandyal, the crop is at boll formation to first picking stage. Excess rain water was drained out from the fields and managed flooding related disorders. Incidence of boll rot was noticed in few fields.

Advisory:

At Guntur, farmers are advised to drain out excess water from the standing crop. Apply N and K as a booster dose @ 25-30 kg urea with 10 kg MoP/acre. Give foliar application of 1- 2% Potassium Nitrate or 1- 2% urea + 1% MoP or 1-2% DAP. In case of any micronutrient deficiencies, spray 1% MgSO₄ (10 g/l) and Borax or Boric acid @1-1.5 g/lit to avoid flower and young boll drop. In case of moist soil condition where manual weeding is not possible, go for application of herbicides like Quizalofop ethyl 5 % EC @ 2 ml /litre of water if the field is infested with grassy weeds or Pyri thio bac sodium 10%EC @ 1.5 ml/lit water for broad leaved weeds or Pyri thio bac sodium 6% + Quizalofop Ethyl 4% MEC @ 2-2.5 ml /litre of water to control both grassy and broad-leaved weeds. Spray Mepiquat chloride @1 ml/litre of water at 75-100 days crop or follow nipping. to prevent excess vegetative growth. If reddening is noticed, spray 1- 2% Potassium Nitrate + spray 1% MgSO₄ (10 g/l) twice at 15 days interval. Spray NAA @ 4 ml/ 10 lit of water to prevent flower drop. Manage initial population of sucking pests by spraying NSKE 5%+ Neem oil 5 ml /litre or neem oil-based formulation 5 ml /litre (300 or 1500 ppm) + 0.05% surfactant. Spray Flonicamid 50% WG @ 80 g/acre or Dinotefuran 20% SG @ 60 g/acre or Tolfe pyrod 15% EC @ 400 ml/acre or Fenpyroximate 5%EC @ 300 ml/acre if infestation of sucking pests seen above ETL. Spray 600g/acre of Copper oxychloride as prophylactic measure to prevent boll rot diseases in cotton and (Carbendazim 12% + Mancozeb 63% WP) @ 2.5-3.0 g/lit to prevent foliar diseases. Give spot drenching of COC twice at 10-15 days interval @ 3g/lit or Carbendazim @1g/lit if wilting of plants is noticed.

At Nandyal, farmers are advised to drain out the water from the fields and spray Potassium nitrate @ 20g/litre of water and 19-19-19 @ 5 g/litre of water to manage flooding related disorders. Install yellow and blue sticky traps @ 8/acre against sucking pests and pheromone traps @2/acre to monitor pink bollworm moth activity. Change lure as per validity. During initial infestation of sucking pests spray NSKE 5% + Neem formulation 5 ml /litre or neem oil-based formulation 5 ml /litre (300 or 1500 ppm) + 0.05%

surfactant. In case sucking pests' infestation crosses ETL, spray Flonicamid 50% WG @ 80 g/acre or Dinotefuran 20% SG @ 60 g/acre or Imidacloprid 17.8% SL @ 60 ml/acre or Buprofezin 25 SC @ 400 ml/acre. Spray Profenofos 50% EC @ 30 ml/10 L (1500 ml/ha) Or Emamectin benzoate 5% SG @ 5 g/10 L (250 g/ha) to manage pink bollworm. As a prophylactic measure, give a spray of (Carbendazim 12%+Mancozeb 63% WP) @30 g or Propineb 70% WP @2.5-3 g/litre or Propiconazole 25 EC @1 ml/litre or Carbendazim 50 WP @0.4 g per litre of water to manage boll rot disease complex, leaf spots due to humidity and congenial weather factors. If parawilt or sudden wilt occur due to heavy rain and water logging after dry spell, drench the affected plants/rows with a mixture of Copper oxychloride 50WP @ 25 g + 200 g Urea in 10 lit of water immediately after occurrence of Parawilt within 48hrs to revive the plants.

KARNATAKA		Actual Rainfall in last week(mm)					Predicted Rainfall in next week (mm)				
		September					September				
		06	07	08	09	10	12	13	14	15	16
	Dharwad	0.8	1	0.4	0.8		1	0.5	0.2	0.1	0.4
	Haveri						0.9	0.5	0.5	0.6	0.7
	Mysore										
Amount of rainfall & colour Code		0.1 to 2.4 mm		2.5 to 15.5 mm		15.6 to 64.4 mm		64.5 to 115.5 mm		115.6 to 204.4	
Rainfall category		Very light rainfall		Light rainfall		Moderate rainfall	Heavy rainfall		Very heavy rainfall		

Crop Condition:

At Dharwad, the sown crop is 88 to 93 days old at flowering to square formation stage. Weeds have dominated the fields. Hand weeding, intercultural operations and spraying of nutrients and insecticides were taken up. Sucking pests like thrips, aphids and jassid infestations were noticed in few pockets. Incidence of Alternaria blight (*Alternaria macrospora*) was reported in moderate status.

At Raichur, the early sown crop is 84 to 90 days old at flowering stage, late sown 62 to 67 days old at squaring stage and very late sown 47 to 52 days old at vegetative stage. Due to continuous rains, the cotton fields are infested with more weeds. Manual weeding and intercultural operations were taken up. Incidence of boll rot was noticed in some fields.

At Chamarajanagar, the crop is 107 to 122 days old at boll formation to boll development stage. Top-dressing of Potash fertilizer, weeding, inter cultivation and earthing-up operations were done. *Cyprus* spp. weeds have dominated the fields. Infestations of jassid (7-8/3leaves), Aphids (35-45/3 leaves), Mirid bug (6-7/25 squares) and pink bollworm 5-6% were noticed.

Advisory:

At Dharwad, farmers are advised to take up hand weeding and intercultivation to manage weeds in the fields wherever cotton crop is sown. Spray Flonicamid 50 WG @ 80 g/acre or Buprofezin 25 SC @ 400 ml/acre or Diafenthiuron 50 WP @ 240 g/acre once sucking pests' crosses ETL. Install pheromone traps @ 2 per acre to monitor pink bollworm moth activity, change lure as per validity. Spray Profenofos 50% EC @ 30 ml/10 lit (1500 ml/ha) or Emamectin benzoate 5% SG @ 5g/10 lit (250 g/ha) of water to manage pink bollworm if infestation crosses ETL. Give foliar spray of (Carbendazim 12%+ Mancozeb 63% WP) @30 g or Propineb 70%WP @2.5-3 g/lit or Propiconazole 25 EC @1 ml/lit or Carbendazim 50 WP @0.4g/ lit of water to manage boll rot disease complex and leaf spots. if noticed in the cotton fields. Spray (Pyroclostrabin 5%+ Metiram 55% WG) @3.5 g/lit of water to manage *Alternaria* blight.

At Raichur, farmers are advised to drain out the excess water from the fields. Take up weeding as and when required. Give recommended dose of fertilizers (DAP, MOP and MgSO₄) for the very late sown and early sown crop as per the stipulated crop duration period. In case of moist soil condition where manual weeding is not possible, go for application of herbicides like Quizalofop ethyl 5 % EC @ 2 ml /litre of water if the field is infested with grassy weeds or Pyriathiobac sodium 10%EC @ 1.5 ml/lit water for broad leaved weeds or Pyriathiobac sodium 6% + Quizalofop Ethyl 4% MEC @ 2-2.5 ml /litre of water to control both grassy and broad-leaved weeds. Manage jassid by spraying Flonicamid @ 4g or Thiamethoxam @ 2g per 10 lit of water. In case of sucking pests above ETL, spray Flonicamid 50 % WG @ 80 g/acre or Dinotefuran 20% SG @ 60 g/acre or Imidacloprid 17.8% SL @ 60 ml/acre or Buprofezin 25 SC @ 400 ml/acre. Spray Profenofos @ 600 ml/acre or Chlorpyrifos @ 500 ml/acre to manage pink bollworm in cotton. For early sown crop, install pheromone traps @ 2/acre to monitor pink bollworm population. To early sown crop, give first foliar spray of 19:19:19 (10 grams in 1 litre of water) + MgSO₄ (10 grams in 1 litre of water) for better growth, boll retention and to manage leaf reddening. Foliar spray of (Carbendazim 12%+Mancozeb 63% WP) @30 g or Propineb 70% WP @2.5-3 g/litre or Propiconazole 25 EC @1 ml/litre or Carbendazim 50 WP @0.4 g per litre of water is recommended to manage boll rot disease complex and leaf spots in cotton.

At Chamarajanagar, farmers are advised to take up hand weeding and intercultivation operations to control weeds. In case of sucking pests above ETL, spray Flonicamid 50%WG @ 4g/10L (200g/ha) Or Dinotefuran 20%SG @ 3g/10L (150g/ha). Install pheromone traps @ 2/ acre to monitor pink bollworm attack. Spray Profenofos 50% EC @ 30 ml/10 L (1500 ml/ha) Or Emamectin benzoate 5% SG @ 5 g/10 L (250 g/ha) to manage pink bollworm. Foliar spray of (Carbendazim 12%+Mancozeb 63% WP) @30 g or Propineb 70% WP @2.5-3 g/litre or Propiconazole 25 EC @1 ml/litre or Carbendazim 50 WP @0.4 g per litre of water is recommended to manage boll rot disease complex and leaf spots in cotton.

TAMIL NADU		Actual Rainfall in last week (mm)					Predicted Rainfall in next week (mm)				
		August / September					September				
		06	07	08	09	10	12	13	14	15	16
	Perambalur	0	0	0	0		0.9	1.7	1.2	0.4	0.4
	Salem	0	0	0	0		0.2	1.5	1.8	1.5	0.5
	Trichy						0.5	0.2	3	0.6	0.4
	Virudhunagar						0.6	0.2	2.4	1.9	1
Amount of rainfall & colour Code		0.1 to 2.4 mm		2.5 to 15.5 mm		15.6 to 64.4 mm		64.5 to 115.5 mm		115.6 to 204.4	
Rainfall category		Very light rainfall		Light rainfall		Moderate rainfall		Heavy rainfall		Very heavy rainfall	

Crop Condition:

At Coimbatore and surrounding areas, the crop is around 21 to 49 days old at initial to vegetative growth stage. Sowing is still in progress. Earthing up, weeding, top dressing and foliar spray to manage sucking pests were done during the reporting period. Weeds like *Cyperus rotundus*, *Tridax procumbens* and *Parthenium* have spread in the fields. Incidence of thrips, aphids and jassid were recorded below ETL.

At Srivilliputhur in and around fields, sowing has commenced. The sown cotton is 7 to 17 days old at germination and initial growth stage. Land preparation and sowing in some parts are in progress. Weeds have started to appear in the fields. No incidence of pests and diseases.

Advisory

At Coimbatore and surrounding areas, farmers are advised to do earthing up. Irrigation and drainage channel should be kept clean to avoid stagnation of water due to excess rain. Take up gap filling 10 DAS and thinning of seedlings on 15 days old crop. Do hand weeding on more than 25 days old crop depending on the weed intensity. Give top dressing with Urea and Potash @50:25 kg/ha for hybrids and for variety @35:20kg/ha. In case of moist soil condition where manual weeding is not possible, go for application of herbicides like Quizalofop ethyl 5 % EC @ 2 ml /litre of water if the field is infested with grassy weeds or Pyriothobac sodium 10%EC @ 1.5 ml/lit water for broad leaved weeds or Pyriothobac sodium 6% + Quizalofop Ethyl 4% MEC @ 2-2.5 ml /litre of water to control both grassy and broad-leaved weeds. Monitor the crop scrupulously for pest and disease incidence. At 50DAS, spray the crop with Neem based insecticide or NSKE 5%+ Neem oil 5 ml /litre or neem oil-based formulation 5 ml /litre (300 or 1500 ppm) +0.05% surfactant. Install yellow sticky traps 8-10/acre to monitor whitefly and jassid incidence. Do spot drenching with Carbendazim 50 WP@12 g/10 lit of water or Trifloxystrobin + Tebuconazole @ 0.75g/lit at the base of affected plants and surrounding healthy plants to manage collar rot incidence. Drench the soil with *Bacillus subtilis* @10ml/lit of water to manage soil borne diseases. Cotton fields should be maintained without *Parthenium* weeds which serves as the source of TSV inoculum

At Srivilliputhur in and around fields, farmers are advised to follow optimum seed rate viz., 15, 20 and 2.5 kg/ha for varieties, rainfed and hybrid cotton respectively. Treat the seeds with 600g of Azospirillum and 600g of Phosphobacteria or 600g of Azophos for one hectare. To maintain optimum plant population, go for recommended spacing of 75x30 cm, 45x15 cm and 120 or 90x60 cm for varieties, rainfed and hybrid cotton, respectively. Follow soil test based fertilizer recommendation or blanket recommendation of 80:40:40, 40:20:20 and 120:60:60 kg/NPK/ha for varieties, rainfed and hybrid cotton, respectively. At the time of sowing, apply basal full dose of P and half dose of N and K. Follow crop rotation where heavy infestation of stem weevil and bollworm complex are prominent. During final plough, apply neem cake @ 250 kg/ha to prevent stem weevil infestation. Clean the farm bunds and other premises to destroy the alternate host of insect pests such as *Abutilon* and other grassy weeds. Treat the seeds with *Trichoderma asperellum* @4g/kg of seeds and soil application at 2.5 kg /ha to prevent the plants from root rot disease