

Updated February 11, 2026

Homework problems for AMAT 300 (Intro to Proofs), Spring 2026. Over the course of the semester I'll add problems to this list, with each problem's due date specified. Each problem is worth 2 points.

Solutions will be gradually added (and may be hastily written without proofreading).

Problem 1 (due Weds 2/4): Let $A = \{20a - 26b \mid a, b \in \mathbb{Z}\}$ and $B = \{2c \mid c \in \mathbb{Z}\}$. Prove that $A = B$. (Do it directly, don't use any number theory tricks you might happen to know.)

Solution: ($\subseteq$): Let $x \in A$, so $x = 20a - 26b$ for some $a, b \in \mathbb{Z}$. Set $c = 10a - 13b$, so $c \in \mathbb{Z}$. Now $x = 2c$, so $x \in B$.

($\supseteq$): Let $x \in B$, so $x = 2c$ for some $c \in \mathbb{Z}$. Set $a = 4c$ and $b = 3c$, so $a, b \in \mathbb{Z}$. Then $x = 2c = 80c - 78c = 20(4c) - 26(3c) = 20a - 26b$, so $x \in A$. $\square$

Problem 2 (due Weds 2/4): Prove that the cardinality of $\{1, \{1\}, \{1, \{1\}\}, \{1, 1\}, \{\{1\}, 1\}\}$ is three. (You need to argue that it has exactly three elements, no more, no less.)

Solution: Since $\{\{1\}, 1\} = \{1, \{1\}\}$ and $\{1, 1\} = \{1\}$, we know that our set equals $\{1, \{1\}, \{1, \{1\}\}\}$, hence has at most 3 elements. Also, $1 \neq \{1\}$ since the latter is a set (with one element) and the former is not, which implies $\{1, \{1\}\}$ has two elements, and we conclude that these three elements are distinct. Thus the original set has exactly 3 elements. $\square$

Problem 3 (due Weds 2/4): Let A and B be non-empty finite sets such that $|\mathcal{P}(A \times B)| = |\mathcal{P}(A) \times \mathcal{P}(B)|$. Prove that A and B each have exactly two elements. (Careful not to *assume* they each have exactly two elements, you have to *prove* that they do.)

Solution: Set $a = |\mathcal{P}(A)|$ and $b = |\mathcal{P}(B)|$. Since $|\mathcal{P}(X)| = 2^{|X|}$ and $|X \times Y| = |X||Y|$ for any finite sets X and Y , we compute that $2^{ab} = 2^a 2^b$, hence $2^{ab} = 2^{a+b}$. Taking $\log_2$ we get $ab = a + b$. Since $a, b \in \mathbb{N}$, dividing by a yields $b = 1 + \frac{b}{a}$ and dividing by b yields $a = \frac{a}{b} + 1$, so $\frac{a}{b}$ and $\frac{b}{a}$ are both integers, which implies $a = b$. Now we solve $a^2 = 2a$ to get $a = b = 2$. $\square$

Problem 4 (due Weds 2/11): Let A and B be sets such that $\mathcal{P}(A \cup B) = \mathcal{P}(A) \cup \mathcal{P}(B)$ and suppose there exists $b_0 \in B \setminus A$. Prove that $A \subseteq B$.

Problem 5 (due Weds 2/11): Let $\Lambda = \{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 \leq 1\}$ be the closed disk of radius 1 centered at the origin. For each $\alpha \in \Lambda$, say $\alpha = (x_0, y_0)$, let $X_\alpha = \{(x, y) \in \mathbb{R}^2 \mid (x - x_0)^2 + (y - y_0)^2 = 1\}$ be the closed disk of radius 1 centered at α . Prove (rigorously) that $\bigcup_{\alpha \in \Lambda} X_\alpha = D$, where $D := \{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 \leq 4\}$ is the closed disk of radius 2 centered at the origin.

Problem 6 (due Weds 2/11): With the same setup as the previous problem, prove (rigorously) that $\bigcap_{\alpha \in \Lambda} X_\alpha = \{(0, 0)\}$.

Problem 7 (due Weds 2/18): Construct truth tables for $\sim (P \Leftrightarrow Q)$ and $(P \vee Q) \wedge (\sim (P \wedge Q))$ (showing the intermediate steps), and compare them. (Which logical operation do these represent?)

Problem 8 (due Weds 2/18): Prove that $n \in \mathbb{Z}$ is a multiple of 24 only if it is a multiple of 4 and 6. (Careful, this is not an “iff”.)

Problem 9 (due Weds 2/18): Let X be a set. Prove that $|\mathcal{P}(X)| \leq 31$ if and only if $|X| \leq 4$.

(Reminder: There's an exam in class on Weds 2/25.)

Problem 10 (due Weds 2/25):

Problem 11 (due Weds 2/25):

Problem 12 (due Weds 2/25):
