

Statistical analysis of experimental data

Parameter Inference

Aleksander Filip Żarnecki



Lecture 06

November 6, 2025

Parameter Inference

- 1 Maximum Likelihood Method
- 2 Confidence intervals
- 3 Real life example
- 4 Homework

General case

Monte Carlo method can be used for very efficient calculation of the arbitrary shape volume with **well controlled precision**. **Works very well also for high-dimensional integrals**

In the general case we can use it to determine an expectation value of a function $h(\mathbf{x})$ of random variable vector $\mathbf{x}$ described by $f(\mathbf{x})$ pdf:

$$\mu_h \equiv \mathbb{E}_f[h(\mathbf{x})] = \int d\mathbf{x} h(\mathbf{x}) f(\mathbf{x})$$

Monte Carlo determination of μ_h assumes we can generate random variables according to $f(\mathbf{x})$. We can then calculate:

$$\mu_{MC} = \lim_{N \rightarrow \infty} \frac{1}{N} \sum_i h(\mathbf{x}_i)$$

where $\mathbf{x}_i$, $i = 1, \dots, N$ are random (input) variables generated from $f(\mathbf{x})$

Weighted mean

For uncorrelated measurements, the best estimate of the expected value is: $\bar{x} = \sigma^2 \sum_i \frac{x_i}{\sigma_i^2}$

In general case, we minimize the variance of the weighted mean: $\mathbb{I} = (1, \dots, 1)^T$

$$\sigma_{\bar{x}}^2 = \mathbf{a}^T \mathbb{C}_x \mathbf{a} - 2 \lambda (\mathbf{a}^T \mathbb{I} - 1)$$

Comparing partial derivatives to zero we get

$$\mathbb{C}_x \mathbf{a} = \lambda \cdot \mathbb{I}$$

where λ can be constrained from the boundary condition $\mathbf{a}^T \mathbb{I} = 1$.

This is a linear set of equations, which can be solved:

$$\mathbf{a} = \frac{\mathbb{C}_x^{-1} \mathbb{I}}{\mathbb{I}^T \mathbb{C}_x^{-1} \mathbb{I}}$$

Maximum Likelihood Method

The product:

$$L = \prod_{j=1}^N f(\mathbf{x}^{(j)}; \lambda)$$

is called a **likelihood function**.

The most commonly used approach to parameter estimation is the **maximum likelihood approach**: as the **best estimate of the parameter set λ** we choose the parameter values for which the **likelihood function** has a (global) maximum.

Frequently used is also log-likelihood function

$$\ell = \ln L = \sum_{j=1}^N \ln f(\mathbf{x}^{(j)}; \lambda)$$

we can look for maximum value of ℓ or minimum of $-2\ell = -2\ln L$

Parameter Inference

- 1 Maximum Likelihood Method
- 2 Confidence intervals
- 3 Real life example
- 4 Homework

Mean estimate

from lecture 05

Let us consider N independent measurements of variable X with non-uniform uncertainties. Assuming measurement fluctuations are described by Gaussian pdf, the likelihood function is:

$$L = \prod_{i=1}^N G(x_i; \mu, \sigma_i) = \prod_{i=1}^N \frac{1}{\sigma_i \sqrt{2\pi}} \exp \left(-\frac{1}{2} \frac{(x_i - \mu)^2}{\sigma_i^2} \right)$$

Mean estimate

from lecture 05

Let us consider N independent measurements of variable X with non-uniform uncertainties. Assuming measurement fluctuations are described by Gaussian pdf, the likelihood function is:

$$L = \prod_{i=1}^N G(x_i; \mu, \sigma_i) = \prod_{i=1}^N \frac{1}{\sigma_i \sqrt{2\pi}} \exp \left(-\frac{1}{2} \frac{(x_i - \mu)^2}{\sigma_i^2} \right)$$

Log-likelihood:

$$\ell = -\frac{1}{2} \sum \frac{(x_i - \mu)^2}{\sigma_i^2} + \text{const}$$

$$\frac{\partial \ell}{\partial \mu} = \sum \frac{x_i - \mu}{\sigma_i^2} = 0$$

$$\Rightarrow \mu = \sigma^2 \sum \frac{x_i}{\sigma_i^2} \quad \text{with } \frac{1}{\sigma^2} = \sum \frac{1}{\sigma_i^2}$$

Mean estimate

modified

Let us consider N independent measurements of variable X with **uniform** uncertainty. Assuming measurement fluctuations are described by Gaussian pdf, the likelihood function is:

$$L = \prod_{i=1}^N G(x_i; \mu, \sigma) = \prod_{i=1}^N \frac{1}{\sigma\sqrt{2\pi}} \exp\left(-\frac{1}{2} \frac{(x_i - \mu)^2}{\sigma^2}\right)$$

Log-likelihood:

$$\ell = -\frac{1}{2\sigma^2} \sum (x_i - \mu)^2 + \text{const}$$

$$\frac{\partial \ell}{\partial \mu} = \frac{1}{\sigma^2} \sum (x_i - \mu) = 0$$

$$\Rightarrow \mu = \frac{1}{N} \sum x_i$$

Variance estimate

The same method can be also used to **estimate the variance** of the Gaussian distribution. Consider partial derivative with respect to σ^2 (we can not neglect normalization now):

$$\ell = -\frac{1}{2\sigma^2} \sum (x_i - \mu)^2 - \frac{N}{2} \ln(2\pi\sigma^2)$$

Variance estimate

The same method can be also used to **estimate the variance** of the Gaussian distribution. Consider partial derivative with respect to σ^2 (we can not neglect normalization now):

$$\begin{aligned}\ell &= -\frac{1}{2\sigma^2} \sum (x_i - \mu)^2 - \frac{N}{2} \ln(2\pi\sigma^2) \\ \frac{\partial \ell}{\partial \sigma^2} &= \frac{1}{2\sigma^4} \sum (x_i - \mu)^2 - \frac{N}{2\sigma^2}\end{aligned}$$

Variance estimate

The same method can be also used to **estimate the variance** of the Gaussian distribution. Consider partial derivative with respect to σ^2 (we can not neglect normalization now):

$$\begin{aligned}\ell &= -\frac{1}{2\sigma^2} \sum (x_i - \mu)^2 - \frac{N}{2} \ln(2\pi\sigma^2) \\ \frac{\partial \ell}{\partial \sigma^2} &= \frac{1}{2\sigma^4} \sum (x_i - \mu)^2 - \frac{N}{2\sigma^2} = 0 \\ \Rightarrow \sigma^2 &= \frac{1}{N} \sum (x_i - \mu)^2\end{aligned}$$

Variance estimate

The same method can be also used to **estimate the variance** of the Gaussian distribution. Consider partial derivative with respect to σ^2 (we can not neglect normalization now):

$$\begin{aligned}\ell &= -\frac{1}{2\sigma^2} \sum (x_i - \mu)^2 - \frac{N}{2} \ln(2\pi\sigma^2) \\ \frac{\partial \ell}{\partial \sigma^2} &= \frac{1}{2\sigma^4} \sum (x_i - \mu)^2 - \frac{N}{2\sigma^2} = 0 \\ \Rightarrow \sigma^2 &= \frac{1}{N} \sum (x_i - \mu)^2\end{aligned}$$

If we extract both μ and σ^2 from the same set of measurements:

$$\sigma^2 = \frac{1}{N} \sum (x_i - \bar{x})^2$$

$\Rightarrow$ ML variance estimator is biased!!! Bessel's correction missing! See lecture 04.

Multiple parameter estimate

Likelihood function (and log-likelihood) can depend on multiple parameters:

$$\boldsymbol{\lambda} = (\lambda_1 \dots \lambda_p) \quad L = \prod_{j=1}^N f(\mathbf{x}^{(j)}; \boldsymbol{\lambda}) \quad \ell = \sum_{j=1}^N \ln f(\mathbf{x}^{(j)}; \boldsymbol{\lambda})$$

Multiple parameter estimate

Likelihood function (and log-likelihood) can depend on multiple parameters:

$$\boldsymbol{\lambda} = (\lambda_1 \dots \lambda_p) \quad L = \prod_{j=1}^N f(\mathbf{x}^{(j)}; \boldsymbol{\lambda}) \quad \ell = \sum_{j=1}^N \ln f(\mathbf{x}^{(j)}; \boldsymbol{\lambda})$$

Best estimate of $\boldsymbol{\lambda}$, for given set of experimental results $\mathbf{x}^{(j)}$, corresponds to maximum of the likelihood function, which can be found by solving a system of equations:

$$\left. \frac{\partial \ell}{\partial \lambda_i} \right|_{i=1 \dots p} = 0$$

Multiple parameter estimate

Likelihood function (and log-likelihood) can depend on multiple parameters:

$$\boldsymbol{\lambda} = (\lambda_1 \dots \lambda_p) \quad L = \prod_{j=1}^N f(\mathbf{x}^{(j)}; \boldsymbol{\lambda}) \quad \ell = \sum_{j=1}^N \ln f(\mathbf{x}^{(j)}; \boldsymbol{\lambda})$$

Best estimate of $\boldsymbol{\lambda}$, for given set of experimental results $\mathbf{x}^{(j)}$, corresponds to maximum of the likelihood function, which can be found by solving a system of equations:

$$\left. \frac{\partial \ell}{\partial \lambda_i} \right|_{i=1 \dots p} = 0$$

The Likelihood Principle

G. Bohm and G. Zech

Given a p.d.f. $f(\mathbf{x}; \boldsymbol{\lambda})$ containing an unknown parameters of interest $\boldsymbol{\lambda}$ and observations $\mathbf{x}^{(j)}$, all information relevant for the estimation of the parameters $\boldsymbol{\lambda}$ is contained in the likelihood function $L(\boldsymbol{\lambda}; \mathbf{x}) = \prod f(\mathbf{x}^{(j)}; \boldsymbol{\lambda})$.

Multivariate Normal Distribution

Consider experiment resulting in a measurement $\mathbf{x} = (x_1, \dots, x_n)$.

If we assume each variable follows Gaussian p.d.f, the most general form of the joint probability distribution is:

$$f(\mathbf{x}; \boldsymbol{\lambda}) = A \exp \left[-\frac{1}{2} (\mathbf{x} - \boldsymbol{\lambda})^\top \mathbb{B} (\mathbf{x} - \boldsymbol{\lambda}) \right]$$

where $\boldsymbol{\lambda}$ is parameter vector and $\mathbb{B}$ is an $n \times n$ matrix.

Since p.d.f. is symmetric about the point $\mathbf{x} = \boldsymbol{\lambda}$:

$$\mathbb{E}(\mathbf{x} - \boldsymbol{\lambda}) = \int d\mathbf{x} (\mathbf{x} - \boldsymbol{\lambda}) f(\mathbf{x}; \boldsymbol{\lambda}) = 0$$

$$\Rightarrow \mathbb{E}(\mathbf{x}) = \boldsymbol{\lambda}$$

Multivariate Normal Distribution

We should note that the derivative of the probability distribution:

$$\frac{\partial f}{\partial \lambda_i} = [(\mathbf{x} - \boldsymbol{\lambda})^\top \mathbb{B}]_i \cdot f(\mathbf{x}; \boldsymbol{\lambda})$$

Multivariate Normal Distribution

We should note that the derivative of the probability distribution:

$$\frac{\partial f}{\partial \lambda_i} = [(\mathbf{x} - \boldsymbol{\lambda})^\top \mathbb{B}]_i \cdot f(\mathbf{x}; \boldsymbol{\lambda})$$

We can now differentiate the formula for $\mathbb{E}(\mathbf{x} - \boldsymbol{\lambda})$ with respect to $\boldsymbol{\lambda}$:

$$\frac{\partial}{\partial \boldsymbol{\lambda}} \int d\mathbf{x} (\mathbf{x} - \boldsymbol{\lambda}) f(\mathbf{x}; \boldsymbol{\lambda}) = \int d\mathbf{x} [(\mathbf{x} - \boldsymbol{\lambda}) (\mathbf{x} - \boldsymbol{\lambda})^\top \mathbb{B} - \mathbb{I}] \cdot f(\mathbf{x}; \boldsymbol{\lambda}) = 0$$

Multivariate Normal Distribution

We should note that the derivative of the probability distribution:

$$\frac{\partial f}{\partial \lambda_i} = [(\mathbf{x} - \boldsymbol{\lambda})^\top \mathbb{B}]_i \cdot f(\mathbf{x}; \boldsymbol{\lambda})$$

We can now differentiate the formula for $\mathbb{E}(\mathbf{x} - \boldsymbol{\lambda})$ with respect to $\boldsymbol{\lambda}$:

$$\frac{\partial}{\partial \boldsymbol{\lambda}} \int d\mathbf{x} (\mathbf{x} - \boldsymbol{\lambda}) f(\mathbf{x}; \boldsymbol{\lambda}) = \int d\mathbf{x} [(\mathbf{x} - \boldsymbol{\lambda}) (\mathbf{x} - \boldsymbol{\lambda})^\top \mathbb{B} - \mathbb{I}] \cdot f(\mathbf{x}; \boldsymbol{\lambda}) = 0$$

and realizing that $\mathbb{B}$ and $\mathbb{I}$ are constant we get

$$\left[\int d\mathbf{x} (\mathbf{x} - \boldsymbol{\lambda}) (\mathbf{x} - \boldsymbol{\lambda})^\top \cdot f(\mathbf{x}; \boldsymbol{\lambda}) \right] \mathbb{B} = \mathbb{I}$$

Multivariate Normal Distribution

We should note that the derivative of the probability distribution:

$$\frac{\partial f}{\partial \lambda_i} = [(\mathbf{x} - \boldsymbol{\lambda})^\top \mathbb{B}]_i \cdot f(\mathbf{x}; \boldsymbol{\lambda})$$

We can now differentiate the formula for $\mathbb{E}(\mathbf{x} - \boldsymbol{\lambda})$ with respect to $\boldsymbol{\lambda}$:

$$\frac{\partial}{\partial \boldsymbol{\lambda}} \int d\mathbf{x} (\mathbf{x} - \boldsymbol{\lambda}) f(\mathbf{x}; \boldsymbol{\lambda}) = \int d\mathbf{x} [(\mathbf{x} - \boldsymbol{\lambda}) (\mathbf{x} - \boldsymbol{\lambda})^\top \mathbb{B} - \mathbb{I}] \cdot f(\mathbf{x}; \boldsymbol{\lambda}) = 0$$

and realizing that $\mathbb{B}$ and $\mathbb{I}$ are constant we get

$$\begin{aligned} \left[\int d\mathbf{x} (\mathbf{x} - \boldsymbol{\lambda}) (\mathbf{x} - \boldsymbol{\lambda})^\top \cdot f(\mathbf{x}; \boldsymbol{\lambda}) \right] \mathbb{B} &= \mathbb{I} \\ \Rightarrow \mathbb{C}_{\mathbf{x}} \mathbb{B} &= \mathbb{I} \quad \Rightarrow \quad \mathbb{B} = \mathbb{C}_{\mathbf{x}}^{-1} \end{aligned}$$

Multivariate Normal Distribution

We can now write the joint probability distribution as:

$$f(\mathbf{x}; \boldsymbol{\lambda}) = A \exp \left[-\frac{1}{2} (\mathbf{x} - \boldsymbol{\lambda})^\top \mathbb{C}_{\mathbf{x}}^{-1} (\mathbf{x} - \boldsymbol{\lambda}) \right]$$

where $\mathbb{C}_{\mathbf{x}}$ is the covariance matrix of variables $\mathbf{x}$.

Multivariate Normal Distribution

We can now write the joint probability distribution as:

$$f(\mathbf{x}; \boldsymbol{\lambda}) = A \exp \left[-\frac{1}{2} (\mathbf{x} - \boldsymbol{\lambda})^\top \mathbb{C}_{\mathbf{x}}^{-1} (\mathbf{x} - \boldsymbol{\lambda}) \right]$$

where $\mathbb{C}_{\mathbf{x}}$ is the covariance matrix of variables $\mathbf{x}$. Log-likelihood:

$$\ell(\boldsymbol{\lambda}; \mathbf{x}) = -\frac{1}{2} (\mathbf{x} - \boldsymbol{\lambda})^\top \mathbb{C}_{\mathbf{x}}^{-1} (\mathbf{x} - \boldsymbol{\lambda}) + \text{const}$$

$$\frac{\partial \ell}{\partial \lambda_i} = [(\mathbf{x} - \boldsymbol{\lambda})^\top \mathbb{C}_{\mathbf{x}}^{-1}]_i \quad \text{similar to p.d.f. derivative}$$

Multivariate Normal Distribution

We can now write the joint probability distribution as:

$$f(\mathbf{x}; \boldsymbol{\lambda}) = A \exp \left[-\frac{1}{2} (\mathbf{x} - \boldsymbol{\lambda})^\top \mathbb{C}_{\mathbf{x}}^{-1} (\mathbf{x} - \boldsymbol{\lambda}) \right]$$

where $\mathbb{C}_{\mathbf{x}}$ is the covariance matrix of variables $\mathbf{x}$. **Log-likelihood:**

$$\ell(\boldsymbol{\lambda}; \mathbf{x}) = -\frac{1}{2} (\mathbf{x} - \boldsymbol{\lambda})^\top \mathbb{C}_{\mathbf{x}}^{-1} (\mathbf{x} - \boldsymbol{\lambda}) + \text{const}$$

$$\frac{\partial \ell}{\partial \lambda_i} = [(\mathbf{x} - \boldsymbol{\lambda})^\top \mathbb{C}_{\mathbf{x}}^{-1}]_i \quad \text{similar to p.d.f. derivative}$$

$$\frac{\partial^2 \ell}{\partial \lambda_i \partial \lambda_j} = - [\mathbb{C}_{\mathbf{x}}^{-1}]_{ij} \quad \Rightarrow \quad \mathbb{C}_{\mathbf{x}} = \left(-\frac{\partial^2 \ell}{\partial \lambda_i \partial \lambda_j} \right)^{-1}$$

Parameter covariance matrix

For the considered case of multivariate normal distribution, best parameter estimates $\hat{\lambda}$ are given by the measured variable values $\mathbf{x}$ (single measurement!)

Unlike parameters λ , parameter estimates $\hat{\lambda}$ are random variables (functions of $\mathbf{x}$ in general) and so we can conclude about the covariance matrix for $\hat{\lambda}$:

$$\mathbb{C}_{\mathbf{x}} = \mathbb{C}_{\hat{\lambda}} = \left(-\frac{\partial^2 \ell}{\partial \lambda_i \partial \lambda_j} \right)^{-1}$$

Parameter covariance matrix

For the considered case of multivariate normal distribution, best parameter estimates $\hat{\lambda}$ are given by the measured variable values $\mathbf{x}$ (single measurement!)

Unlike parameters λ , parameter estimates $\hat{\lambda}$ are random variables (functions of $\mathbf{x}$ in general) and so we can conclude about the covariance matrix for $\hat{\lambda}$:

$$\mathbb{C}_{\mathbf{x}} = \mathbb{C}_{\hat{\lambda}} = \left(-\frac{\partial^2 \ell}{\partial \lambda_i \partial \lambda_j} \right)^{-1}$$

Knowing the likelihood function, we can not only estimate parameter values, but also extract uncertainties and correlations of these estimates!

Parameter covariance matrix

For the considered case of multivariate normal distribution, best parameter estimates $\hat{\lambda}$ are given by the measured variable values $\mathbf{x}$ (single measurement!)

Unlike parameters λ , parameter estimates $\hat{\lambda}$ are random variables (functions of $\mathbf{x}$ in general) and so we can conclude about the covariance matrix for $\hat{\lambda}$:

$$\mathbb{C}_{\mathbf{x}} = \mathbb{C}_{\hat{\lambda}} = \left(-\frac{\partial^2 \ell}{\partial \lambda_i \partial \lambda_j} \right)^{-1}$$

Knowing the likelihood function, we can not only estimate parameter values, but also extract uncertainties and correlations of these estimates!

For the uncorrelated parameters (diagonal covariance matrix):

$$\sigma_{\hat{\lambda}_i} = \left(-\frac{\partial^2 \ell}{\partial \lambda_i^2} \right)^{-1/2}$$

Parameter covariance matrix

Considered example was based on the Gaussian distribution.

Standard deviation is one of the parameters of the p.d.f., can be easily extracted from log-likelihood:

$$\sigma_i = \sqrt{C_{ii}}$$

However, this procedure works only in the Gaussian approximation.

How to define parameter uncertainty in the general case?

Parameter covariance matrix

Considered example was based on the Gaussian distribution.

Standard deviation is one of the parameters of the p.d.f., can be easily extracted from log-likelihood:

$$\sigma_i = \sqrt{C_{ii}}$$

However, this procedure works only in the Gaussian approximation.

How to define parameter uncertainty in the general case?

For 1-D case we have:

$$\ell(\hat{\lambda} + \sigma; x) = \ln f(x; \hat{\lambda} + \sigma) = \ell(\hat{\lambda}; x) - \frac{1}{2}$$

Parameter covariance matrix

Considered example was based on the Gaussian distribution.

Standard deviation is one of the parameters of the p.d.f., can be easily extracted from log-likelihood:

$$\sigma_i = \sqrt{C_{ii}}$$

However, this procedure works only in the Gaussian approximation.

How to define parameter uncertainty in the general case?

For 1-D case we have:

$$\ell(\hat{\lambda} + 2\sigma; x) = \ln f(x; \lambda + 2\sigma) = \ell(\hat{\lambda}; x) - \frac{4}{2}$$

Parameter covariance matrix

Considered example was based on the Gaussian distribution.

Standard deviation is one of the parameters of the p.d.f., can be easily extracted from log-likelihood:

$$\sigma_i = \sqrt{C_{ii}}$$

However, this procedure works only in the Gaussian approximation.

How to define parameter uncertainty in the general case?

For 1-D case we have:

$$\ell(\hat{\lambda} + 3\sigma; x) = \ln f(x; \hat{\lambda} + 3\sigma) = \ell(\hat{\lambda}; x) - \frac{9}{2}$$

Parameter covariance matrix

Considered example was based on the Gaussian distribution.

Standard deviation is one of the parameters of the p.d.f., can be easily extracted from log-likelihood:

$$\sigma_i = \sqrt{C_{ii}}$$

However, this procedure works only in the Gaussian approximation.

How to define parameter uncertainty in the general case?

Recipe for a parameter uncertainty

G. Bohm and G. Zech

Standard error intervals of the extracted parameter are defined by the decrease of the log-likelihood function by 0.5 for one, by 2 for two and by 4.5 for three standard deviations.

This definition works for arbitrary p.d.f. shape, also for multiple parameters

Multiple parameter estimate


Example log-likelihood function contours

$$\ell(\mu, \sigma; \hat{\mu}, \hat{\sigma})$$

for a sample of 10 events from normal distribution, with extracted parameter values $\hat{\mu} = 1$ and $\hat{\sigma} = 2$.

Figure from:
G. Bohm and G. Zech, Introduction to
Statistics and Data Analysis for Physicists,
Verlag Deutsches Elektronen-Synchrotron,
3rd edition

Do we understand the shape of ℓ ?



Multiple parameter estimate

For the set $\mathbf{x}$ of N measurements we can write:

$$\begin{aligned}\sum (x_i - \mu)^2 &= \sum (x_i^2 - 2\mu x_i + \mu^2) \\ &= N (\langle x^2 \rangle - 2\mu \langle x \rangle + \mu^2)\end{aligned}$$

Multiple parameter estimate

For the set $\mathbf{x}$ of N measurements we can write:

$$\begin{aligned}\sum (x_i - \mu)^2 &= \sum (x_i^2 - 2\mu x_i + \mu^2) \\ &= N (\langle x^2 \rangle - 2\mu \langle x \rangle + \mu^2) \\ &= N (\hat{\sigma}^2 + \hat{\mu}^2 - 2\mu \hat{\mu} + \mu^2) \quad \hat{\sigma}^2 = \langle x^2 \rangle - \langle x \rangle^2\end{aligned}$$

Multiple parameter estimate

For the set $\mathbf{x}$ of N measurements we can write:

$$\begin{aligned}\sum (x_i - \mu)^2 &= \sum (x_i^2 - 2\mu x_i + \mu^2) \\ &= N (\langle x^2 \rangle - 2\mu \langle x \rangle + \mu^2) \\ &= N (\hat{\sigma}^2 + \hat{\mu}^2 - 2\mu \hat{\mu} + \mu^2) \quad \hat{\sigma}^2 = \langle x^2 \rangle - \langle x \rangle^2\end{aligned}$$

Log-likelihood function for the example is then:

$$\ell(\mu, \sigma; \hat{\mu}, \hat{\sigma}) = -\frac{1}{2\sigma^2} \sum (x_i - \mu)^2 - \frac{N}{2} \ln(2\pi\sigma^2)$$

Multiple parameter estimate

For the set $\mathbf{x}$ of N measurements we can write:

$$\begin{aligned}\sum (x_i - \mu)^2 &= \sum (x_i^2 - 2\mu x_i + \mu^2) \\ &= N (\langle x^2 \rangle - 2\mu \langle x \rangle + \mu^2) \\ &= N (\hat{\sigma}^2 + \hat{\mu}^2 - 2\mu \hat{\mu} + \mu^2) \quad \hat{\sigma}^2 = \langle x^2 \rangle - \langle x \rangle^2\end{aligned}$$

Log-likelihood function for the example is then:

$$\ell(\mu, \sigma; \hat{\mu}, \hat{\sigma}) = -\frac{N}{2\sigma^2} (\hat{\sigma}^2 + (\mu - \hat{\mu})^2) - \frac{N}{2} \ln(2\pi\sigma^2)$$

This corresponds to the Gaussian shape for μ , but very asymmetric for σ ...

Also, the two parameters are not independent!


Multiple parameter estimate

06_mlm_func.ipynb

 Open in Colab

Example log-likelihood function contours for a **sample of 10 events** from normal distribution, with extracted parameter values $\hat{\mu} = 1$ and $\hat{\sigma} = 2$.

Log-likelihood contours for example measurement



Result from G. Bohm and G. Zech is nicely reproduced...

Multiple parameter estimate

As mentioned before, $\hat{\mu}$ and $\hat{\sigma}$ (extracted from measurements $\mathbf{x}$) are random variables! We can calculate their joint probability distribution:

$$f(\hat{\mu}, \hat{\sigma}^2; \mu, \sigma) = A \cdot \exp\left(-\frac{N(\hat{\mu} - \mu)^2}{2\sigma^2}\right) \cdot \left(\frac{N\hat{\sigma}^2}{\sigma^2}\right)^{k-1} \exp\left(-\lambda \frac{N\hat{\sigma}^2}{\sigma^2}\right)$$

where $\frac{N\hat{\sigma}^2}{\sigma^2}$ is distributed according to the Gamma distribution with $k = (N - 1)/2$ and $\lambda = 1/2$ (particular case referred to as χ^2 distribution; we will discuss it at the next lectures).

Multiple parameter estimate

As mentioned before, $\hat{\mu}$ and $\hat{\sigma}$ (extracted from measurements $\mathbf{x}$) are random variables! We can calculate their joint probability distribution:

$$f(\hat{\mu}, \hat{\sigma}^2; \mu, \sigma) = A \cdot \exp\left(-\frac{N(\hat{\mu} - \mu)^2}{2\sigma^2}\right) \cdot \left(\frac{N\hat{\sigma}^2}{\sigma^2}\right)^{k-1} \exp\left(-\lambda \frac{N\hat{\sigma}^2}{\sigma^2}\right)$$

where $\frac{N\hat{\sigma}^2}{\sigma^2}$ is distributed according to the Gamma distribution with $k = (N - 1)/2$ and $\lambda = 1/2$ (particular case referred to as χ^2 distribution; we will discuss it at the next lectures).

The two variables, $\hat{\mu}$ and $\hat{\sigma}$, are independent!

PDF for $\hat{\sigma}$ is asymmetric, but much less than the likelihood function !!!

This simple example demonstrates also that the likelihood function for the parameters can be very different than the probability distribution for the measurements...

Multiple parameter estimate


06_mlm_func2.ipynb

 Open in Colab

One needs to stress that likelihood function for p.d.f. parameters are not equivalent to probability distribution of parameter estimators!

Contours of the joint probability distribution function $f(\hat{\mu}, \hat{\sigma}; \mu, \sigma)$ for $\mu = 1$ and $\sigma = 2$

Log-likelihood contours for example measurement



Bessel's factor clearly visible!

Contours correspond to log f decrease by 0.5, 2 and 4.5 from maximum.

Multiple parameter estimate


06_mlm_mc_2d.ipynb

 Open in Colab

One needs to stress that likelihood function for p.d.f. parameters are not equivalent to probability distribution of parameter estimators!

Results of Monte Carlo simulation (contours from 1 000 000 experiments)

Distribution of estimated mean vs sigma



In each experiment, mean and sigma are calculated from 10 generated numbers

Parameter Inference

- 1 Maximum Likelihood Method
- 2 Confidence intervals
- 3 Real life example
- 4 Homework

Presenting measurement results

When doing the measurement, we usually quote the final result as numerical value (with units) and estimated uncertainty:

$$x \pm \sigma_x$$

We can often calculate the uncertainty from the data itself (eg. when result is obtained by averaging a large number of independent measurements) or from the variation of the log-likelihood function.

Attributing proper uncertainty to the result is crucial!


But what does it tell us after all?!

Normal distribution

Meaning of σ is well defined for **Gaussian distribution**.

Probability for the experimental result to be consistent with the true value within $\pm N\sigma$:

	$1 - \alpha$	
$\pm 1 \sigma$	$\Rightarrow 68.27$	%
$\pm 2 \sigma$	$\Rightarrow 95.45$	%
$\pm 3 \sigma$	$\Rightarrow 99.73$	%
$\pm 4 \sigma$	$\Rightarrow 99.9937$	%
$\pm 5 \sigma$	$\Rightarrow 99.999943$	%




There is a non-zero chance for deviation greater than 5σ , but it is extremely small

Normal distribution

Meaning of σ is well defined for **Gaussian distribution**.

Probability for the experimental result to **differ from the true** value by more than $N\sigma$:

	α	
$\pm 1 \sigma$	$\Rightarrow 31.73$	%
$\pm 2 \sigma$	$\Rightarrow 4.55$	%
$\pm 3 \sigma$	$\Rightarrow 0.27$	%
$\pm 4 \sigma$	$\Rightarrow 0.0063$	%
$\pm 5 \sigma$	$\Rightarrow 0.000057$	%



Fluctuations up and down are observed with equal probability...

Normal distribution


06_interval.ipynb

 Open in Colab

Results of the Monte Carlo test

($\mu = 0$, $\sigma = 1$, 100 000 000 generations)

Probability distribution from MC



Down fluctuations:

$< -1 \sigma$ $p = 0.15864225$

$< -2 \sigma$ $p = 0.0227686$

$< -3 \sigma$ $p = 0.00134872$

$< -4 \sigma$ $p = 3.204E-05$

$< -5 \sigma$ $p = 3.2E-07$

Fluctuations up:

$> 1 \sigma$ $p = 0.15861607$

$> 2 \sigma$ $p = 0.02275479$

$> 3 \sigma$ $p = 0.00135162$

$> 4 \sigma$ $p = 3.263E-05$


$> 5 \sigma$ $p = 2.8E-07$

Good agreement with expectations

Poisson distribution

Results of the Monte Carlo test
($\mu = 10$, 100 000 000 generations)

Probability distribution from MC



06_interval_2.ipynb

 Open in Colab

Down fluctuations:

$< -1 \sigma$ $p = 0.1301325$

$< -2 \sigma$ $p = 0.01033324$

$< -3 \sigma$ $p = 4.55\text{E-}05$

$< -4 \sigma$ $p = 0$

$< -5 \sigma$ $p = 0$

Fluctuations up:

$> 1 \sigma$ $p = 0.13553469$

$> 2 \sigma$ $p = 0.02702754$

$> 3 \sigma$ $p = 0.00344483$

$> 4 \sigma$ $p = 0.00029364$

$> 5 \sigma$ $p = 1.774\text{e-}05$

Much longer tail of positive fluctuations!


Gamma distribution

06_interval_3.ipynb

 Open in Colab

Results of the Monte Carlo test
($\mu = \sigma^2 = 10$, 10 000 000 generations)

Probability distribution from MC



Down fluctuations:

$< -1 \sigma$	$p = 0.1533757$
$< -2 \sigma$	$p = 0.0046445$
$< -3 \sigma$	$p = 0$
$< -4 \sigma$	$p = 0$
$< -5 \sigma$	$p = 0$

Fluctuations up:


$> 1 \sigma$	$p = 0.1554444$
$> 2 \sigma$	$p = 0.0368358$
$> 3 \sigma$	$p = 0.0067406$
$> 4 \sigma$	$p = 0.0010105$
$> 5 \sigma$	$p = 0.0001305$

Even longer, 5σ fluctuations not excluded

Normal distribution in N-D

It is also important to notice that the fractions presented previously (eg. 68% within $\pm 1\sigma$) refer to one-dimensional normal distribution only!

If we consider 2-D distribution



Less than 40% is contained inside 1σ contour...

Fractions within $N\sigma$ contours:

Deviation	Dimension			
	1	2	3	4
1σ	0.683	0.393	0.199	0.090
2σ	0.954	0.865	0.739	0.594
3σ	0.997	0.989	0.971	0.939
4σ	1.	1.	0.999	0.997

1σ fraction above 50% only for $N=1$!

G. Bohm and G. Zech

Interpreting results

As demonstrated above, quoting numerical result with uncertainties gives only partial information on the measurement...

It is sufficient, if we can assume normal distribution of the variable.

Also, we need to assume that the width of the distribution does not depend on the measured parameter. Only then the likelihood function will be Gaussian as well...

Interpreting results

As demonstrated above, quoting numerical result with uncertainties gives only partial information on the measurement...

It is sufficient, if we can assume normal distribution of the variable.

Also, we need to assume that the width of the distribution does not depend on the measured parameter. Only then the likelihood function will be Gaussian as well...

In the general case, parameter uncertainty does not give us full information on the shape (in particular the tails) of the distribution...

How should we present results of the experiment, if we are more concerned about the probability of (large) result fluctuations?...

Interpreting results

There is also another problem, which has to be noticed!

So far we have only considered distribution of experimental results for given probability distribution, $f(\mathbf{x}; \boldsymbol{\lambda})$, when the parameter values $\boldsymbol{\lambda}$ are known.

Interpreting results

There is also another problem, which has to be noticed!

So far we have only considered distribution of experimental results for given probability distribution, $f(\mathbf{x}; \boldsymbol{\lambda})$, when the parameter values $\boldsymbol{\lambda}$ are known.

The actual situation is usually different: for given set of measurements $\mathbf{x}$ we extract estimates of the parameter values $\hat{\boldsymbol{\lambda}}$.

Uncertainties estimated from log-likelihood variation indicate the expected level of agreement (in Gaussian approximation) between our estimate $\hat{\boldsymbol{\lambda}}$ and the true parameter values $\boldsymbol{\lambda}$.

Can we present measurement results in a way which gives us more precise information about the possible fluctuations in the estimate $\hat{\boldsymbol{\lambda}}$?

Interpreting results

There is also another problem, which has to be noticed!

So far we have only considered distribution of experimental results for given probability distribution, $f(\mathbf{x}; \boldsymbol{\lambda})$, when the parameter values $\boldsymbol{\lambda}$ are known.

The actual situation is usually different: for given set of measurements $\mathbf{x}$ we extract estimates of the parameter values $\hat{\boldsymbol{\lambda}}$.

Uncertainties estimated from log-likelihood variation indicate the expected level of agreement (in Gaussian approximation) between our estimate $\hat{\boldsymbol{\lambda}}$ and the true parameter values $\boldsymbol{\lambda}$.

Can we present measurement results in a way which gives us more precise information about the possible fluctuations in the estimate $\hat{\boldsymbol{\lambda}}$?

Yes, but we need to define the problem differently... We should not consider probability of $\hat{\boldsymbol{\lambda}}$...

Frequentist confidence intervals

Classical (frequentist) definition of the confidence interval refers directly to the probability distribution of the experimental results, $f(\mathbf{x}; \lambda)$.

We do not try to make any prediction (nor guess) about the “probability” (degree of belief) of given parameter value λ . This is the Bayesian concept we will discuss later...

In the frequentist approach we consider individually each λ value.

Given value of λ is allowed (on given confidence level, C.L.), if the actual outcome of our experiment, x_m , is within the corresponding **probability interval** for **variable x** for this value of λ .

Frequentist confidence intervals

Classical (frequentist) definition of the confidence interval refers directly to the probability distribution of the experimental results, $f(\mathbf{x}; \lambda)$.

We do not try to make any prediction (nor guess) about the “probability” (degree of belief) of given parameter value λ . This is the Bayesian concept we will discuss later...

In the frequentist approach we consider individually each λ value.

Given value of λ is allowed (on given confidence level, C.L.), if the actual outcome of our experiment, x_m , is within the corresponding **probability interval** for **variable x** for this value of λ .

This definition clearly depends on the way we define probability intervals for x .

So this is rather a general concept, more assumptions are needed.

We always refer to probability distribution $f(\mathbf{x}; \lambda)$ for random variable x !

Frequentist confidence intervals

It is interesting to note that the modern concept of confidence intervals was proposed by Polish statistician **Jerzy Neyman** only in 1937

Jerzy Neyman obtained his PhD (1924) and habilitation (1928) at UW

X—Outline of a Theory of Statistical Estimation Based on the
Classical Theory of Probability

By J. NEYMAN

Reader in Statistics, University College, London

(Communicated by H. JEFFREYS, F.R.S.—Received 20 November, 1936—Read 17 June, 1937)

CONTENTS

	Page
I—INTRODUCTORY	333
(a) General Remarks, Notation, and Definitions	333
(b) Review of the Solutions of the Problem of Estimation Advanced Hereto	343
(c) Estimation by Unique Estimate and by Interval	346
II—CONFIDENCE INTERVALS	347
(a) Statement of the Problem	347
(b) Solution of the Problem of Confidence Intervals	350
(c) Example I	356
(d) Example II	362
(e) Family of Similar Regions Based on a Sufficient System of Statistics	364
(f) Example IIa	367

J.Neyman, Phil. Trans. Royal Soc. London,
Series A, 236 333-80 (1937).

Frequentist confidence intervals

As mentioned above, to define confidence interval for our parameter, we need to define how the **probability interval** for our measurement is defined.

There are three “natural” choices:

$$CL = 1 - \alpha$$
$$\alpha \ll 1$$

- We constrain the measurement from above, define upper limit x_{ul} :

$$\int_{-\infty}^{x_{ul}} dx f(x; \lambda) = CL \quad \text{or} \quad \int_{x_{ul}}^{+\infty} dx f(x; \lambda) = \alpha$$

- We constrain the measurement from below, define lower limit x_{ll} :

$$\int_{-\infty}^{x_{ll}} dx f(x; \lambda) = \alpha \quad \text{or} \quad \int_{x_{ll}}^{+\infty} dx f(x; \lambda) = CL$$

Frequentist confidence intervals

As mentioned above, to define confidence interval for our parameter, we need to define how the **probability interval** for our measurement is defined.

There are three “natural” choices:

$$CL = 1 - \alpha$$

$$\alpha \ll 1$$

- We constrain the measurement from above, define upper limit x_{ul} :

$$\int_{-\infty}^{x_{ul}} dx f(x; \lambda) = CL \quad \text{or} \quad \int_{x_{ul}}^{+\infty} dx f(x; \lambda) = \alpha$$

- We constrain the measurement from below, define lower limit x_{ll} :

$$\int_{-\infty}^{x_{ll}} dx f(x; \lambda) = \alpha \quad \text{or} \quad \int_{x_{ll}}^{+\infty} dx f(x; \lambda) = CL$$

- We define central probability interval $[x_1, x_2]$:

as presented for Gaussian pdf

$$\int_{-\infty}^{x_1} dx f(x; \lambda) = \alpha/2 \quad \text{and} \quad \int_{x_2}^{+\infty} dx f(x; \lambda) = \alpha/2$$

Frequentist confidence intervals

Confidence intervals for parameter λ reflect the probability intervals for $\mathbf{x}$.

Let us consider the simplest example, Gaussian pdf: $f(x; \lambda, \sigma) = \frac{1}{\sigma\sqrt{2\pi}} \exp\left(-\frac{1}{2} \frac{(x-\lambda)^2}{\sigma^2}\right)$

Assuming σ is known (and fixed), **what is the central 90% C.L. interval for λ ?** $\alpha = 0.1$

Frequentist confidence intervals

Confidence intervals for parameter λ reflect the probability intervals for $\mathbf{x}$.

Let us consider the simplest example, Gaussian pdf: $f(x; \lambda, \sigma) = \frac{1}{\sigma\sqrt{2\pi}} \exp\left(-\frac{1}{2} \frac{(x-\lambda)^2}{\sigma^2}\right)$

Assuming σ is known (and fixed), **what is the central 90% C.L. interval for λ ?** $\alpha = 0.1$

From the Gaussian pdf properties we can directly obtain:

$$x_1 = \lambda - 1.64\sigma \quad x_2 = \lambda + 1.64\sigma$$

Frequentist confidence intervals

Confidence intervals for parameter λ reflect the probability intervals for $\mathbf{x}$.

Let us consider the simplest example, Gaussian pdf: $f(x; \lambda, \sigma) = \frac{1}{\sigma\sqrt{2\pi}} \exp\left(-\frac{1}{2} \frac{(x-\lambda)^2}{\sigma^2}\right)$

Assuming σ is known (and fixed), **what is the central 90% C.L. interval for λ ?** $\alpha = 0.1$

From the Gaussian pdf properties we can directly obtain:

$$x_1 = \lambda - 1.64\sigma \quad x_2 = \lambda + 1.64\sigma$$

Definition of the confidence interval for λ is based on the condition: **we measure $x = x_m$**

$$x_1 < x_m < x_2$$


Which is fulfilled for all λ in range:

$$\lambda_1 = x_m - 1.64\sigma < \lambda < x_m + 1.64\sigma = \lambda_2$$

So the central 90% C.L. interval for λ is $[x_m - 1.64\sigma, x_m + 1.64\sigma]$...

Frequentist confidence intervals


Graphical presentation of the procedure for Gaussian pdf with $\sigma \equiv 1$



G.J.Feldman, R.D.Cousins,
A Unified Approach to the Classical Statistical Analysis of Small Signals,
Phys.Rev.D57:3873-3889,1998; arXiv:physics/9711021

Frequentist confidence intervals


Graphical presentation of the general procedure



G.J.Feldman, R.D.Cousins,
A Unified Approach to the Classical Statistical Analysis of Small Signals,
Phys.Rev.D57:3873-3889,1998; arXiv:physics/9711021

Frequentist confidence intervals


General procedure



- calculate limits of probability intervals for x , $x_1(\theta)$ and $x_2(\theta)$, for different values of θ

Frequentist confidence intervals


General procedure



- calculate limits of probability intervals for x , $x_1(\theta)$ and $x_2(\theta)$, for different values of θ
- calculated intervals define the “accepted region” in (θ, x)

Frequentist confidence intervals


General procedure



- calculate limits of probability intervals for x , $x_1(\theta)$ and $x_2(\theta)$, for different values of θ
- calculated intervals define the “accepted region” in (θ, x)
- confidence interval for θ is defined by drawing line $x = x_m$ in the accepted region

Frequentist confidence intervals

General procedure



- calculate limits of probability intervals for x , $x_1(\theta)$ and $x_2(\theta)$, for different values of θ
 - calculated intervals define the “accepted region” in (θ, x)
 - confidence interval for θ is defined by drawing line $x = x_m$ in the accepted region
- $\Rightarrow$ limit on θ for given x_m , $\theta_1(x_m)$, corresponds to limit on x for given θ : $x_m = x_1(\theta_1)$.

Frequentist limits

Use of central interval limits is one of the possible definitions, other approaches are possible...

The procedure is more unique, if we want to constrain the parameter from above or below.

We first define an upper or lower limit for the measured value x , for given λ (at $CL = 1 - \alpha$):

$$\int_{x_{ul}(\lambda)}^{+\infty} dx f(x; \lambda) = \alpha \quad \text{or} \quad \int_{-\infty}^{x_{ll}(\lambda)} dx f(x; \lambda) = \alpha$$

Frequentist limits

Use of central interval limits is one of the possible definitions, other approaches are possible...
The procedure is more unique, if we want to constrain the parameter from above or below.
We first define an upper or lower limit for the measured value x , for given λ (at CL = $1 - \alpha$):

$$\int_{x_{ul}(\lambda)}^{+\infty} dx f(x; \lambda) = \alpha \quad \text{or} \quad \int_{-\infty}^{x_{ll}(\lambda)} dx f(x; \lambda) = \alpha$$

Assuming that $\langle x \rangle$ increases with λ , and the measurement resulted in value x_m :

- Upper limit on parameter λ can be defined by the condition: $x_{ll}(\lambda_{ul}) = x_m$

For $\lambda > \lambda_{ul}$, the probability that the experiment result is not larger than x_m is less than α .

$\Rightarrow$ We state that values $\lambda > \lambda_{ul}$ are excluded at $(1 - \alpha)$ confidence level (CL)

Frequentist limits

Use of central interval limits is one of the possible definitions, other approaches are possible...
The procedure is more unique, if we want to constrain the parameter from above or below.
We first define an upper or lower limit for the measured value x , for given λ (at CL = $1 - \alpha$):

$$\int_{x_{ul}(\lambda)}^{+\infty} dx f(x; \lambda) = \alpha \quad \text{or} \quad \int_{-\infty}^{x_{ll}(\lambda)} dx f(x; \lambda) = \alpha$$

Assuming that $\langle x \rangle$ increases with λ , and the measurement resulted in value x_m :

- Upper limit on parameter λ can be defined by the condition: $x_{ll}(\lambda_{ul}) = x_m$

For $\lambda > \lambda_{ul}$, the probability that the experiment result is not larger than x_m is less than α .

⇒ We state that values $\lambda > \lambda_{ul}$ are excluded at $(1 - \alpha)$ confidence level (CL)

- Lower limit on parameter λ can be defined by the condition: $x_{ul}(\lambda_{ll}) = x_m$

Procedure example


06_gauss_interval.ipynb

 Open in Colab

Let us consider the 90% CL interval (or 95% CL limits) for Gaussian pdf: width fixed $\sigma \equiv 1$

$$f(x; \mu, \sigma) = \frac{1}{\sigma\sqrt{2\pi}} \exp\left(-\frac{1}{2} \frac{(x - \mu)^2}{\sigma^2}\right)$$

Confidence intervals



- calculate limits of probability intervals for x : $x_1(\mu)$ and $x_2(\mu)$, for different values of μ

Procedure example


06_gauss_interval.ipynb

 Open in Colab

Let us consider the 90% CL interval (or 95% CL limits) for Gaussian pdf: width fixed $\sigma \equiv 1$

$$f(x; \mu, \sigma) = \frac{1}{\sigma\sqrt{2\pi}} \exp\left(-\frac{1}{2} \frac{(x - \mu)^2}{\sigma^2}\right)$$

Confidence intervals



- calculate limits of probability intervals for x : $x_1(\mu)$ and $x_2(\mu)$, for different values of μ
- calculated intervals define the “accepted region” in the (μ, x) plane

Procedure example


06_gauss_interval.ipynb

 Open in Colab

Let us consider the 90% CL interval (or 95% CL limits) for Gaussian pdf: width fixed $\sigma \equiv 1$

$$f(x; \mu, \sigma) = \frac{1}{\sigma\sqrt{2\pi}} \exp\left(-\frac{1}{2} \frac{(x - \mu)^2}{\sigma^2}\right)$$

Confidence intervals



- calculate limits of probability intervals for x : $x_1(\mu)$ and $x_2(\mu)$, for different values of μ
- calculated intervals define the “accepted region” in the (μ, x) plane

Procedure example


06_gauss_interval.ipynb

 Open in Colab

Let us consider the 90% CL interval (or 95% CL limits) for Gaussian pdf: width fixed $\sigma \equiv 1$

$$f(x; \mu, \sigma) = \frac{1}{\sigma\sqrt{2\pi}} \exp\left(-\frac{1}{2} \frac{(x - \mu)^2}{\sigma^2}\right)$$

Confidence intervals



- calculate limits of probability intervals for x : $x_1(\mu)$ and $x_2(\mu)$, for different values of μ
- calculated intervals define the “accepted region” in the (μ, x) plane

Procedure example


06_gauss_interval.ipynb

 Open in Colab

Let us consider the 90% CL interval (or 95% CL limits) for Gaussian pdf: width fixed $\sigma \equiv 1$

$$f(x; \mu, \sigma) = \frac{1}{\sigma\sqrt{2\pi}} \exp\left(-\frac{1}{2} \frac{(x - \mu)^2}{\sigma^2}\right)$$

Confidence intervals



- calculate limits of probability intervals for x : $x_1(\mu)$ and $x_2(\mu)$, for different values of μ
- calculated intervals define the “accepted region” in the (μ, x) plane
- confidence interval for μ is found by drawing line $x = x_m$ in the accepted region

Procedure example


06_gauss_interval.ipynb

 Open in Colab

Let us consider the 90% CL interval (or 95% CL limits) for Gaussian pdf: width fixed $\sigma \equiv 1$

$$f(x; \mu, \sigma) = \frac{1}{\sigma\sqrt{2\pi}} \exp\left(-\frac{1}{2} \frac{(x - \mu)^2}{\sigma^2}\right)$$

Confidence intervals



- calculate limits of probability intervals for x : $x_1(\mu)$ and $x_2(\mu)$, for different values of μ
 - calculated intervals define the “accepted region” in the (μ, x) plane
 - confidence interval for μ is found by drawing line $x = x_m$ in the accepted region
- ⇒ limit on μ for given x_m , $\mu_1(x_m)$, corresponds to the probability limit on x for given μ : $x_m = x_1(\mu_1)$.

Parameter Inference

- 1 Maximum Likelihood Method
- 2 Confidence intervals
- 3 Real life example
- 4 Homework



Contents lists available at [ScienceDirect](https://www.sciencedirect.com)

Physics Letters B

www.elsevier.com/locate/physletb



Limits on the effective quark radius from inclusive ep scattering at HERA



ZEUS Collaboration

H. Abramowicz^{y,31}, I. Abt^t, L. Adamczyk^h, M. Adamus^{ae}, S. Antonelli^b, V. Aushev^q, O. Behnke^j, U. Behrens^j, A. Bertolin^v, S. Bhadra^{ag}, I. Bloch^k, E.G. Boos^o, I. Brock^c, N.H. Brook^{ac}, R. Brugnera^w, A. Bruni^a, P.J. Bussey^l, A. Caldwell^t, M. Capua^e, C.D. Catterall^{ag}, J. Chwastowski^g, J. Ciborowski^{ad,33}, R. Ciesielski^{j,16}, A.M. Cooper-Sarkar^u, M. Corradi^{a,11}, R.K. Dementiev^s, R.C.E. Devenish^u, S. Dusini^v, B. Foster^{m,23}, G. Gach^h, E. Gallo^{m,24}, A. Garfagnini^w, A. Geiser^j, A. Gizhko^j, L.K. Gladilin^s, Yu.A. Golubkov^s, G. Grzelak^{ad}, M. Guzik^h, C. Gwenlan^u, W. Hain^j, O. Hlushchenko^q, D. Hochman^{af}, R. Horiⁿ, Z.A. Ibrahim^f, Y. Iga^x, M. Ishitsuka^z, F. Januschek^{j,17}, N.Z. Jomhari^f, I. Kadenko^q, S. Kananov^y, U. Karshon^{af}, P. Kaur^{d,12}, D. Kisielewska^h, R. Klanner^m, U. Klein^{j,18}

HERA electron(positron)-proton collider at DESY




HERA I 1994–2000

about 100pb^{-1} collected per experiment
mainly e^+p data, unpolarised

HERA II 2002–2007


about 400pb^{-1} per experiment
similar amount of e^-p and e^+p data
with longitudinal polarization of $e^\pm$ beams (30–40%)
and small samples collected at reduced proton beam energy




Deep Inelastic $e^\pm p$ Scattering

Main process studied by H1 and ZEUS


NC DIS



CC DIS



Kinematic variables:



$$Q^2 = -(k - k')^2$$

|virtuality| of the exchanged boson

$$x = \frac{Q^2}{2P \cdot (k - k')}$$


fraction of proton momenta carried by struck quark

$$y = \frac{P \cdot (k - k')}{P \cdot k}$$

fraction of lepton energy transferred in the proton rest frame

SM predictions vs measurements from HERA

H1 and ZEUS



Input data for the study:
combined H1 and ZEUS **NC** and **CC** DIS cross sections.

Cross sections comparable for the highest Q^2 values

$$Q^2 \sim M_Z^2, M_W^2$$

QCD+EW analysis shows good agreement with the SM


Phys. Rev. D 93 (2016) 092002, [arXiv:1603.09628](https://arxiv.org/abs/1603.09628)

However, high precision data could also be used to look for **possible new (BSM) effects**...

Quark form factor

“classical” method to look for possible fermion (sub)structure.

If a quark has **finite size**, the standard model cross-section is expected to decrease at high momentum transfer:



$$\frac{d\sigma}{dQ^2} = \frac{d\sigma^{SM}}{dQ^2} \cdot \left[1 - \frac{R_q^2}{6} Q^2 \right]^2 \cdot \left[1 - \frac{R_e^2}{6} Q^2 \right]^2$$

where R_q is the root mean-square radius of the electroweak charge distribution in the quark.

We do not consider the possibility of finite electron size...

same dependence expected for e^+p and e^-p !


We can try to fit our “extended model” (with one extra parameter) to the data...

ZEUS data

06_zeus_rq_limit.ipynb

 Open in Colab

We count NC DIS events in Q^2 bins, looking at the highest 18 bins:




- ZEUS NC DIS data
- SM expectations

R_q expectations

06_zeus_rq_limit.ipynb

 Open in Colab

If quarks had finite size, we should observe linear dependence of $\sqrt{N_{obs}/N_{exp}}$ on Q^2 :



- MC experiment
for $R_q = 0.001$ fm
- SM expectations

$$\sqrt{\frac{N_{obs}}{N_{exp}}} = \left[1 - \frac{R_q^2}{6} Q^2 \right]$$


note log scale in Q^2

Frequentist approach

06_zeus_rq_limit.ipynb

 Open in Colab

For different R_q values, we generate multiple MC experiments to find out, if our actual measurement is within the probability interval



Fit to ZEUS data:
 $R_q^2 = 2.75 \cdot 10^{-8} fm^2$

Our measurement is
consistent with SM
predictions ($R_q = 0$)


It is outside the range
expected for $R_q = 0.001$ fm
 $\Rightarrow$ can be excluded with
very high confidence!

Frequentist upper limit

06_zeus_rq_limit.ipynb

 Open in Colab

For different R_q values, we calculate probability of obtaining the R_q^2 values from the fit higher and lower than the actual ZEUS measurement:



Upper limit on the quark radius is defined as R_q value for which:

$$p(R_{fit}^2 > R_{data}^2) = CL$$

$$p(R_{fit}^2 < R_{data}^2) = 1 - CL$$

For $CL = 0.95$:

$$R_q < 0.00055 \text{ fm}$$

$$R_q < 0.55 \cdot 10^{-18} \text{ m}$$

Actual HERA analysis much more parameters to consider in the fit!

Likelihood function taking into account the possible BSM contributions

$$-2 \ell(\mathbf{p}, \mathbf{s}, \eta) = \sum_i \frac{\left[m^i(\mathbf{p}, \eta) + \sum_j s_j \gamma_j^i m^i(\mathbf{p}, \eta) - \mu_0^i \right]^2}{\left(\delta_{i,\text{stat}}^2 + \delta_{i,\text{uncor}}^2 \right) (\mu_0^i)^2} + \sum_j s_j^2$$

μ_0^i and $m^i(\mathbf{p}, \eta)$ are measured and predicted (SM+BSM) cross sections, γ_j^i , $\delta_{i,\text{stat}}$ and $\delta_{i,\text{uncor}}$ are correlated systematic, statistical and uncorrelated systematic uncertainties of data point i
 $\mathbf{p}$ and $\mathbf{s}$ are vectors of PDF parameters p_k (describe proton structure) and systematic shifts s_j ,
 η is the parameter describing BSM contribution (for quark radius: $\eta = R_q^2$)

$\Rightarrow$ we look for global likelihood maximum for the combined HERA data

$$R_q^2 \text{ Data} = -0.2 \cdot 10^{-33} \text{ cm}^2$$

This one number summarizes the results of all our measurements...

Limit setting


Based on MC experiments: “MC replicas” of the collected data (frequentist approach!).

Replicas are generated sets of cross-section values that are calculated for given $R_q^2 \text{ True}$ and varied randomly according to the statistical and systematic uncertainties (including correlations) of the input data.

Each replica is then used as an input to SM+BSM fit $\Rightarrow R_q^2 \text{ Fit}$

Many replicas for each considered $R_q^2 \text{ True}$ value $\Rightarrow$ distribution of $R_q^2 \text{ Fit}$

$R_q^2 \text{ True}$ is tested by comparing $R_q^2 \text{ Fit}$ distribution with the value of $R_q^2 \text{ Data}$



Limit setting


The probability of obtaining a $R_q^{2 \text{ Fit}}$ value smaller than that obtained for the actual data

$$\text{Prob}(R_q^{2 \text{ Fit}} < R_q^{2 \text{ Data}})$$

is studied as a function of $R_q^{2 \text{ True}}$

$R_q^{2 \text{ True}}$ values corresponding to the probability smaller than 5% are excluded at the 95% C.L.

limits obtained for fixed SM parameters are too strong by about 10% !




$$R_q < 0.43 \cdot 10^{-18} \text{ m}$$

Results

Phys. Lett. B757 (2016) 468, arXiv:1604.01280

ZEUS



$$-(0.47 \cdot 10^{-18} \text{ m})^2 < R_q^2 < (0.43 \cdot 10^{-18} \text{ m})^2$$

Parameter Inference

- 1 Maximum Likelihood Method
- 2 Confidence intervals
- 3 Real life example
- 4 Homework

Homework

Solutions to be uploaded by November 19.

Consider the example discussed today:

sample of 10 measurements from the normal distributions $G(\mu, \sigma)$.

Assuming that the measured variance of the sample is $\hat{\sigma}^2 = 3$:

- calculate the frequentist 95% CL upper limit on the true variance of the distribution, σ^2 , based on the properties of the Gamma distribution.

Please note that the frequentist limit definition considers always distribution of the random variable ($\hat{\sigma}^2$) not of the parameter (σ^2). SciPy library can be used.

- verify the result by the means of the Monte Carlo simulation.