

# **Title of the article**

John Smith,<sup>1</sup> James Smith,<sup>1,2</sup>, and Jane Smith<sup>2</sup>

---

Corresponding author: Jane Smith, Department of Geography, Ohio State University, Columbus, Ohio, USA. (j.smith@ohio.edu)

<sup>1</sup>Department of Hydrology and Water Resources, University of Arizona, Tucson, Arizona, USA.

<sup>2</sup>Department of Geography, Ohio State University, Columbus, Ohio, USA.

<sup>2</sup> **Abstract.**

<sup>3</sup>     The abstract goes here.

## 1. Introduction

As the horizontal resolution of general circulation models (GCMs) has increased over the last decade, the simulated climatology of tropical cyclones has improved greatly [*Oouchi et al.*, 2006; *Bengtsson et al.*, 2007; *Zhao et al.*, 2009; *Murakami et al.*, 2012; *Manganello et al.*, 2012; *Satoh et al.*, 2012; *Strachan et al.*, 2013; *Zarzycki and Jablonowski*, 2014; *Wehner et al.*, 2014].

Kevin, add some references/bulk to the next sentence?

While the climatology of simulated TCs across different climate models is large *Camargo* [2013]; *Walsh et al.* [2015], it has also been shown that within individual models, storms also show sensitivity to both the dynamical core (e.g. *Reed and Jablonowski* [2012]; *Zhao et al.* [2012]) as well as the choice of subgrid parameterizations (e.g., *Reed and Jablonowski* [2011]; *Lim et al.* [2014])

In this manuscript we describe another mechanism through which simulated TC properties are influenced by model design choices, in particular, the manner in which the ocean and atmosphere are coupled within the climate system. Section 2 provides an introduction to the modeling system used in this study and how coupling between the atmosphere and ocean is treated. Section 3 details the sensitivity of TCs to the ocean grid using a deterministic forecast framework while Section 3.1 investigates the impact on multi-year climate simulations. Section 4 assesses the results and offers further insight into their implications.

## 2. Model description

### 2.1. Community Atmosphere Model

In this paper, we utilize the Community Atmosphere Model (CAM), version 5 [Neale et al., 2010]. The Spectral Element (SE) dynamical core option in CAM is used for all simulations. SE is the newest dynamical core available in CAM and is based upon continuous Galerkin spectral finite elements which are applied on a cubed-sphere grid [Taylor et al., 1997; Thomas and Loft, 2005; Taylor and Fournier, 2010; Dennis et al., 2012].

CAM-SE is coupled to land, ocean, and ice model components within the Community Earth System Modelling (CESM) framework. The land model is the Community Land Model (CLM) version 4.0 run in satellite phenology (SP) mode on a 0.23 by 0.31° latitude-longitude grid [Oleson et al., 2010]. While CESM also allows for coupling to dynamic ocean and ice models, all of the simulations in this project utilize prescribed SSTs and ice cover concentrations.

## 2.2. Coupling within CESM

Because earth system model components generally are not integrated on the same spatial grid, CPL7 is used to couple various systems within the CESM framework [Craig et al., 2012]. The coupler utilizes conservative remapping weights to regrid quantities between model components.

(Describe coupling procedure in CESM; draw schematic; confirm with T. Craig or Mariana).

During coupling between the atmosphere and ocean, state variables from the atmospheric grid are conservatively remapped to the ocean grid. Surface momentum stress ( $\tau$ ) and sensible and latent heat fluxes are calculated on the ocean grid. The quantities

are then passed back to the atmospheric grid, allowing these quantities to be available to both the atmospheric dynamical core and subgrid physical parameterizations.

Generally, if the atmosphere and ocean are not on the same grid, the atmospheric grid consists of finer grid spacing.

Prescribed SSTs and ice are passed to the model on a  $1^\circ \times 1^\circ$  grid and internally interpolated to the ocean and ice grids.

### 3. Deterministic simulations

To assess the differences in simulated TCs in a controlled, deterministic manner, we utilize two nearly identical CAM setups to complete short-term forecast simulations of observed storms. These simulations utilize the new, variable-resolution capability of CAM-SE [Zarzycki *et al.*, 2014].

The model is configured with an atmospheric grid with  $1/8^\circ$  ( $\sim 14\text{km}$ ) resolution over the Atlantic Ocean and are initialized with a digitally-filtered atmospheric analysis from the National Center for Atmospheric Predictions's Global Data Assimilation System (GDAS). Observed SSTs are taken from NOAAOI and provided as input to the model on a  $1^\circ \times 1^\circ$  grid. The land surface is modeled by the Community Land Model (CLM) version 4.0 and is initialized with a state nudged to be in balance with the atmospheric initial conditions. The model setup and initialization are further detailed in Zarzycki and Jablonowski [2015].

The only difference between the two setups is the grid used by the data ocean and ice models. The first set of simulations uses a displaced tripole grid with an equivalent resolution of  $1^\circ$  ( $\text{gx1v6}$ ) while the second uses an ocean grid identical to the atmospheric grid with an equivalent resolution of  $1/8^\circ$ . Since the SST and ice cover data are provided at coarser scales than the model interpolates to, any differences in the results arise due to

the differences in calculating of surface fluxes and momentum drag on the corresponding ocean grids.

Figure 1 shows the 120-hour forecast for Hurricane Leslie in the North Atlantic Ocean from the 2012 hurricane season. The simulation was initialized at 00Z on August 31st, 2012 and is valid at 00Z on September 5th, 2012. The forecast using the  $1^\circ$  ocean grid is on the left (Fig. 1a,c), with the  $1/8^\circ$  ocean grid on the right (Fig. 1b,d). All fields are extracted from the atmospheric model component. The top panels depict instantaneous lowest model level wind (black vectors) as well as the surface frictional stress vector (red). In the Fig. 1a, we note many instances where the vectors are not aligned. This results from the surface stress being calculated on the coarser grid. The atmospheric dynamical core then subsamples this coarser information to provide stress information at the same resolution used by the numerics. In Fig. 1b, the wind and stress vectors are essentially parallel ( $180^\circ$  difference), indicating that the frictional drag is acting in parallel opposition to the wind. The higher resolution ocean grid preserves the resolution of the surface wind field during stress calculations. Additionally, the maximum magnitudes of the stress vectors are larger at the storm's radius of maximum wind in Fig. 1b. Since stress is directly correlated with the surface wind speed, this highlights that maxima in the wind field at the atmospheric grid cell scale are preserved with the higher resolution ocean grid, whereas these maxima are “smoothed” in the calculation where wind is first averaged to the coarser ocean grid (Fig. 1a). This is further evidenced by the fact that the integrated dot product (over a  $5^\circ \times 5^\circ$  domain centered over the TC minimum surface pressure) of the two fields is approximately 10% smaller in the simulations using the  $1^\circ$  ocean grid.

Therefore, the use of the coarser ocean grid results in a weaker frictional force used by the atmospheric dynamics.

The cumulative surface heat flux is shown on the bottom (sensible plus latent) for the two storms at the same forecast time. It is readily apparent that the coarser ocean grid (Fig. 1c) provides information back to the atmosphere with significantly less spatial structure than the  $1/8^\circ$  ocean grid (Fig. 1d). While the difference in  $5^\circ \times 5^\circ$  integrated heat flux is relatively small (approximately 1%), it is clear that the spatial structure of the heat flux field is very different between the two model configurations. This may also play a role in storm dynamics with the  $1^\circ$  ocean grid providing a larger, more diffuse source of surface heating to the TC core than the high-resolution grid.

### 3.1. Climate simulations

**JULIO/KEVIN CLIMATE SIMS**

## 4. Discussion

**Discussion goes here.**

The issues outlined in the manuscript are rather trivial to correct for data ocean models, where specifying SSTs and applying coupled atmosphere-ocean calculations on the same grid is straightforward. More problematic adjustments arise when coupling to a dynamical ocean model. The vast majority of coupled simulations not only utilize differing resolutions between different model components, but also different numerical techniques and grids. Therefore, remapping between components is an absolute necessity.

The obvious recommendation to alleviate coupling inconsistencies when it is not feasible to use identical grids is to calculate these quantities on the finest resolution grid of

the coupled system. In the vast majority of earth system models, this is typically the atmosphere. Performing coupling in this manner ensures that information passed back to a model component has not be interpolated to a resolution coarser than the component's native resolution during the coupling process.

However, it is not clear that technique is fully appropriate for dynamical ocean models, where aspects such as turbulent mixing may be sufficiently non-linear that merely averaging from a higher resolution grid may not be the most appropriate mechanism.

**Acknowledgments.** This work was partially supported by a grant from the Spanish Ministry of Science and Technology.

## References

- Bengtsson, L., K. I. Hodges, and M. Esch (2007), Tropical cyclones in a T159 resolution global climate model: comparison with observations and re-analyses, *Tellus A*, 59(4), 396–416, doi:10.1111/j.1600-0870.2007.00236.x.
- Camargo, S. J. (2013), Global and regional aspects of tropical cyclone activity in the cmip5 models, *Journal of Climate*, 26(24), 9880–9902, doi:10.1175/JCLI-D-12-00549.1.
- Craig, A. P., M. Vertenstein, and R. Jacob (2012), A new flexible coupler for earth system modeling developed for CCSM4 and CESM1, *International Journal of High Performance Computing Applications*, 26(1), 31–42, doi:10.1177/1094342011428141.
- Dennis, J. M., J. Edwards, K. J. Evans, O. Guba, P. H. Lauritzen, A. A. Mirin, A. St-Cyr, M. A. Taylor, and P. H. Worley (2012), CAM-SE: A scalable spectral element dynamical core for the Community Atmosphere Model, *International Journal of High Performance Computing Applications*, 26(1), 74–89, doi:10.1177/1094342011428142.



- 129 Lim, Y.-K., S. D. Schubert, O. Reale, M.-I. Lee, A. M. Molod, and M. J. Suarez (2014),  
130 Sensitivity of tropical cyclones to parameterized convection in the nasa geos-5 model,  
131 *Journal of Climate*, *28*(2), 551–573, doi:10.1175/JCLI-D-14-00104.1.
- 132 Manganello, J. V., K. I. Hodges, J. L. Kinter, B. A. Cash, L. Marx, T. Jung,  
133 D. Achuthavarier, J. M. Adams, E. L. Altshuler, B. Huang, E. K. Jin, C. Stan, P. Tow-  
134 ers, and N. Wedi (2012), Tropical cyclone climatology in a 10-km global atmospheric  
135 GCM: Toward weather-resolving climate modeling, *Journal of Climate*, *25*(11), 3867–  
136 3893, doi:10.1175/JCLI-D-11-00346.1.
- 137 Murakami, H., Y. Wang, H. Yoshimura, R. Mizuta, M. Sugi, E. Shindo, Y. Adachi,  
138 S. Yukimoto, M. Hosaka, S. Kusunoki, T. Ose, and A. Kitoh (2012), Future changes in  
139 tropical cyclone activity projected by the new high-resolution MRI-AGCM, *Journal of*  
140 *Climate*, *25*(9), 3237–3260, doi:10.1175/JCLI-D-11-00415.1.
- 141 Neale, R. B., C.-C. Chen, A. Gettelman, P. H. Lauritzen, S. Park, D. L. Williamson,  
142 A. J. Conley, R. Garcia, D. Kinnison, J.-F. Lamarque, D. Marsh, M. Mills, A. K.  
143 Smith, S. Tilmes, F. Vitt, P. Cameron-Smith, W. D. Collins, M. J. Iacono, , R. C.  
144 Easter, X. Liu, S. J. Ghan, P. J. Rasch, and M. A. Taylor (2010), Description of the  
145 NCAR Community Atmosphere Model (CAM 5.0), *NCAR Technical Note NCAR/TN-*  
146 *486+STR*, National Center for Atmospheric Research, Boulder, Colorado.
- 147 Oleson, K., D. Lawrence, G. Bonan, M. Flanner, E. Kluzek, P. Lawrence, S. Levis,  
148 S. Swenson, P. Thornton, A. Dai, M. Decker, R. Dickinson, J. Feddema, C. Heald,  
149 F. Hoffman, J. Lamarque, N. Mahowald, G. Niu, T. Qian, J. Randerson, S. Running,  
150 K. Sakaguchi, A. Slater, R. Stockli, A. Wang, Z. Yang, X. Zeng, and X. Zeng (2010),  
151 Technical description of version 4.0 of the Community Land Model (CLM), *NCAR Tech-*

*nical Note NCAR/TN-478+STR*, National Center for Atmospheric Research, Boulder,  
Colorado, doi:10.5065/D6FB50WZ.

Oouchi, K., J. Yoshimura, H. Yoshimura, R. Mizuta, S. Kusunoki, and A. Noda (2006),  
Tropical cyclone climatology in a global-warming climate as simulated in a 20 km-  
mesh global atmospheric model: Frequency and wind intensity analyses, *Journal of the  
Meteorological Society of Japan*, *84*(2), 259–276.

Reed, K. A., and C. Jablonowski (2011), Impact of physical parameterizations on idealized  
tropical cyclones in the Community Atmosphere Model, *Geophysical Research Letters*,  
*38*(4), doi:10.1029/2010GL046297.

Reed, K. A., and C. Jablonowski (2012), Idealized tropical cyclone simulations of inter-  
mediate complexity: A test case for AGCMs, *Journal of Advances in Modeling Earth  
Systems*, *4*, M04,001, doi:10.1029/2011MS000099.


Satoh, M., K. Oouchi, T. Nasuno, H. Taniguchi, Y. Yamada, H. Tomita, C. Kodama,  
J. Kinter, D. Achuthavarier, J. Manganello, B. Cash, T. Jung, T. Palmer, and N. Wedi  
(2012), The intra-seasonal oscillation and its control of tropical cyclones simulated by  
high-resolution global atmospheric models, *Climate Dynamics*, *39*(9-10), 2185–2206,  
doi:10.1007/s00382-011-1235-6.

Strachan, J., P. L. Vidale, K. Hodges, M. Roberts, and M.-E. Demory (2013), Investi-  
gating global tropical cyclone activity with a hierarchy of AGCMs: The role of model  
resolution, *Journal of Climate*, *26*(1), 133–152, doi:10.1175/JCLI-D-12-00012.1.

Taylor, M., J. Tribbia, and M. Iskandarani (1997), The spectral element method for the  
shallow water equations on the sphere, *Journal of Computational Physics*, *130*, 92–108.

- 174 Taylor, M. A., and A. Fournier (2010), A compatible and conservative spectral element  
175 method on unstructured grids, *Journal of Computational Physics*, *229*(17), 5879–5895.
- 176 Thomas, S., and R. Loft (2005), The NCAR spectral element climate dynamical core:  
177 Semi-implicit Eulerian formulation, *Journal of Scientific Computing*, *25*(1), 307–322,  
178 doi:10.1007/s10915-004-4646-2.
- 179 Walsh, K. J. E., S. J. Camargo, G. A. Vecchi, A. S. Daloz, J. Elsner, K. Emanuel, M. Horn,  
180 Y.-K. Lim, M. Roberts, C. Patricola, E. Scoccimarro, A. H. Sobel, S. Strazzo, G. Vil-  
181 larini, M. Wehner, M. Zhao, J. P. Kossin, T. LaRow, K. Oouchi, S. Schubert, H. Wang,  
182 J. Bacmeister, P. Chang, F. Chauvin, C. Jablonowski, A. Kumar, H. Murakami, T. Ose,  
183 K. A. Reed, R. Saravanan, Y. Yamada, C. M. Zarzycki, P. L. Vidale, J. A. Jonas, and  
184 N. Henderson (2015), Hurricanes and climate: the U.S. CLIVAR working group on  
185 hurricanes, *Bulletin of the American Meteorological Society*, doi:10.1175/BAMS-D-13-  
186 00242.1.
- 187 Wehner, M. F., K. A. Reed, F. Li, Prabhat, J. Bacmeister, C.-T. Chen, C. Paciorek,  
188 P. J. Gleckler, K. R. Sperber, W. D. Collins, A. Gettelman, and C. Jablonowski  
189 (2014), The effect of horizontal resolution on simulation quality in the Community  
190 Atmospheric Model, CAM5.1, *Journal of Advances in Modeling Earth Systems*, doi:  
191 10.1002/2013MS000276.
- 192 Zarzycki, C. M., and C. Jablonowski (2014), A multidecadal simulation of Atlantic tropical  
193 cyclones using a variable-resolution global atmospheric general circulation model, *Jour-  
194 nal of Advances in Modeling Earth Systems*, *6*(3), 805–828, doi:10.1002/2014MS000352.
- 195 Zarzycki, C. M., and C. Jablonowski (2015), Experimental tropical cyclone forecasts using  
196 a variable-resolution global model, *Monthly Weather Review*, in prep.

- Zarzycki, C. M., M. N. Levy, C. Jablonowski, J. R. Overfelt, M. A. Taylor, and P. A. Ullrich (2014), Aquaplanet experiments using CAM’s variable-resolution dynamical core, *Journal of Climate*, *27*, 5481–5503, doi:10.1175/JCLI-D-14-00004.1.
- Zhao, M., I. M. Held, S. J. Lin, and G. A. Vecchi (2009), Simulations of global hurricane climatology, interannual variability, and response to global warming using a 50-km resolution GCM, *Journal of Climate*, *22*(24), 6653–6678.
- Zhao, M., I. M. Held, and S.-J. Lin (2012), Some counterintuitive dependencies of tropical cyclone frequency on parameters in a GCM, *Journal of the Atmospheric Sciences*, *69*(7), 2272–2283, doi:10.1175/JAS-D-11-0238.1.



**Figure 1.** Figure caption