

TEXT

Colin M. Zarzycki¹, Kevin A. Reed², Julio Bacmeister¹, Susan C. Bates¹, Anthony P. Craig¹, and J. E. Truesdale¹

¹Climate and Global Dynamics, National Center for Atmospheric Research, Boulder, Colorado, USA.

²School of Marine and Atmospheric Sciences, State University of New York at Stony Brook, Stony Brook, New York, USA.

Correspondence to: Colin M. Zarzycki, Climate and Global Dynamics, National Center for Atmospheric Research, Boulder, Colorado, USA. (zarzycki@ucar.edu)

Abstract. The abstract goes here.

1 Introduction

The use of general circulation models (GCMs) to evaluate global tropical cyclone (TC) characteristics in current and future climate has grown considerably over the last decade. It has been shown that GCMs can model TCs at horizontal resolutions of approximately 100 km grid spacing with limitations (e.g., Bengtsson et al., 2007; Knutson et al., 2010; Strachan et al., 2013). As models have advanced to even higher horizontal resolutions (i.e., ≤ 50 km) the simulated climatology of tropical cyclones has improved greatly (e.g., Oouchi et al., 2006; Zhao et al., 2009; Murakami et al., 2012; Manganello et al., 2012; Satoh et al., 2012; Bacmeister et al., 2014; Wehner et al., 2014; Reed et al., 2015). Furthermore, the use of variable-resolution GCMs have shown to be useful for the study of regional TC climatologies at reduced computational cost compared to equivalent global high-resolution simulations, providing further resources capable of pushing climate simulations to finer grid spacings (Zarzycki et al., 2014a; Zarzycki and Jablonowski, 2014).

Recently, intercomparisons have shown that the range of simulated TC climatology across different climate models can be large (Camargo, 2013; Walsh et al., 2015). It has also been shown that within individual GCMs TC characteristics can vary greatly depending on model design choices. Various studies have documented the large uncertainty in TC simulations due to the choice of individual subgrid parameterizations, such as cumulus parameterizations (e.g., Kim et al., 2012; Reed and Jablonowski, 2011; Lim et al., 2014)), while others have focused on differences due to changes in whole parameterization suites (Reed and Jablonowski, 2011; Bacmeister et al., 2014). The dynamical core, the main fluid flow component of a GCM, has also been shown to be an important source of uncertainty for TC simulations, though less widely documented (Reed and Jablonowski, 2012; Zhao et al., 2012; Reed et al., 2015).

In this manuscript we describe another mechanism through which simulated TC properties are influenced by model design choices, in particular, the manner in which the ocean and atmosphere are coupled within the climate system. In particular, we will utilize the Community Atmosphere Model 5 (CAM5), within the Community Earth System Model (CESM), to explore the impact of two different strategies for coupling to a prescribed ocean. CAM5 has shown increasing ability to model tropical cyclones at high horizontal resolutions of 0.25° (Bacmeister et al., 2014; Wehner et al., 2014; Wehner et al., 2015; Reed et al., 2015) and a similar model setup will be used for part of this study.

The remainder of the paper is organized as follows. Section 2 provides an introduction to the modeling system used in this study and how coupling between the atmosphere and ocean is treated. Section 3 investigates the impact on multi-year climate simulations while Section 4 details the sensitivity of TCs to the ocean grid using a deterministic forecast framework while. Section 5 discusses the results and offers further insight into their implications.

2 Model description

2.1 Community Earth System Model

In this paper, we utilize CESM which is a community climate model which allows for atmospheric simulations to be coupled to land, ocean, and ice models (Hurrell et al., 2013). The atmospheric component CAM5 (Neale et al., 2010) is configured with the Spectral Element (SE) dynamical core. SE is the newest dynamical core available in CAM5 and is based upon continuous Galerkin spectral finite elements which are applied on a cubed-sphere grid (Taylor et al., 1997; Thomas and Loft, 2005; Taylor and Fournier, 2010). In addition to attractive conservation properties (Taylor, 2011), CAM-SE has shown appealing scaling properties since atmospheric primitive equations are solved locally on individual elements (Dennis et al., 2012; Evans et al., 2013). The land model is the Community Land Model (CLM) version 4.0 run in satellite phenology (SP) mode (Oleson et al., 2010). While CESM also allows for coupling to dynamic ocean and ice models, all of the simulations here utilize prescribed SSTs and ice cover concentrations. In the default CESM configuration, prescribed SSTs and ice are passed to the model on a $1^\circ \times 1^\circ$ grid and internally interpolated to the ocean and ice grids.

2.2 Coupling within CESM

Because earth system model components generally are not integrated on the same spatial grid, CPL7 is used to couple these components to one another within the CESM framework (Craig et al., 2012). The coupler utilizes conservative remapping weights to regrid quantities which are needed across model components. Figure 1 provides a schematic of the coupling process when differences exist between the resolution of the atmosphere and ocean grids in CESM. In this case, the atmospheric grid (red) is of finer resolution, which is a frequently used configuration of coupled climate mod-




Figure 1. Coupling procedure in CESM. Red (blue) boxes indicate atmospheric (ocean) grid cells. Black (Gray) solid (dashed) vectors show surface wind (wind stress) vectors.

els. Atmospheric state variables, such as winds (black vectors; taken here to approximate the flow associated with a tropical cyclone), are computed on the atmospheric grid (Fig. 1a). When coupling is required, these values are then conservatively remapped to the ocean grid (blue) (Fig. 1b). Surface momentum stress (τ , gray vectors) and sensible and latent heat fluxes are calculated on the ocean grid using these remapped values (Fig. 1c). The calculated quantities are then conservatively remapped back to the atmospheric grid (Fig. 1d), where they are used by the atmospheric component of the model for integration (Fig. 1e).

Probably should work in that the coupling is more straightforward when calculated on the same high-res grid - though it may be obvious

3 Climate simulations

We first compare TC statistics in two multi-decadal climate simulations using 0.25° (denoted as ne120 on the spectral element cubed sphere grid) resolution for the atmosphere. Both simulations follow Atmospheric Model Intercomparison Project (AMIP) protocols (Gates, 1992) and are coupled to CLM with an equivalent grid resolution of 0.25° . The first simulation is coupled to a prescribed ocean/ice model using a displaced tripole grid at approximately 1° horizontal resolution (ne120_gx1v6), which is coarser than both the atmosphere and land models. The second simulation is identical to the first except the prescribed ocean/ice model operates on the same 0.25° grid as the atmosphere and land (ne120_ne120). It is worth nothing that while the ocean/ice model operates

on the 0.25° grid, the data is interpolated from the same 1.0° observational dataset used in the first simulation (this is correct, right?).

Both simulations are integrated from 1980 to 2005. Taylor statistics for the 1980-1999 global-mean quantities for sea-level pressure (PSL), total precipitable water (TMQ), total precipitation rate (PRECT), 200 hPa zonal wind (U200), 850 hPa zonal wind (U850), 600 hPa relative humidity (RH600) and 500 hPa temperature (T500) are shown in Fig. 2. The two simulations are compared to observational datasets including NCEP (Kalnay et al., 1996) (PSL, U200, U850, RH600, T500), MERRA (Rienecker et al., 2011) (TMQ), and TRMM (Huffman et al., 2007) (PRECT). The absolute distance from the origin (lower left) represents the magnitude of the spatial variability within the domain (as measured by normalized standard deviation) while the spatial correlation is plotted as the radial angle between the model marker and the origin. A comprehensive discussion of Taylor diagram analysis can be found in Taylor (2001). Red dots highlight the climatology of the ne120_ne120 simulation while blue dots show the same for the ne120_gx1v6 simulation. This analysis is concerned with the relative difference between the two simulations and, therefore, whether or not mean climatology is impacted by choice of coupling grid. A thorough analysis understanding why each parameter is modeled with their particular skill in CAM5 itself is beyond the scope of this paper. We do note, however, that the results are generally consistent with skill scores reported in previous CAM5 modeling studies, such as Bacmeister et al. (2014) (their Fig. 2 and Fig. 3) and Zarzycki et al. (2015) (their Fig. 9).

The most notable result from assessing this skill as the two simulations are highly similar in a global climatological sense. All markers in the ne120_gx1v6 simulation overlap with their corresponding variable from the ne120_ne120 simulation. The occurrence of this overlap highlights that the mean climate state is not impacted by choice of coupling strategy in the climate simulations.

While the mean climatology of the two simulations are essentially identical, notable differences arise when comparing TC statistics between the two simulations. Table 1 displays storm counts for all TCs, storms that reach hurricane strength and storms that reach major hurricane strength on the Saffir-Simpson scale (Simpson, 1974) (i.e., categories 4 and 5) for each simulation for 1980 through 2005. Observations from the International Best Track Archive for Climate Stewardship (IBTrACS, Knapp et al. (2010)) for the same time period are provided as a reference. Tropical cyclones are objectively tracked in model output using the method first outlined in Vitart et al. (1997) and updated by Knutson et al. (2007). The TC tracker uses 3-hourly model output and is described in detail in Zhao et al. (2009). Previous work using this technique to find TCs in CAM/CESM output have produced a realistic storm climatology both spatially and in terms of storm intensity Reed et al. (2015). For the tracker, the surface winds (commonly taken to be at a height of 10 m) are approximated from the lowermost model level winds (≈ 60 m) and a logarithmic law similar to the approach described in Zarzycki and Jablonowski (2014).




Figure 2. Taylor diagram for globally- and annually-averaged climate statistics. Blue circles represent the results from AMIP simulation coupled to 1° ocean grid (ne120_gx1v6) and red circles represent the same for simulation using 0.25° ocean grid (ne120_ne120). See text for description of the diagram and explanation of acronyms.

Table 1. Annual frequency of global tropical cyclones that reach tropical storm (cat. 0-5), hurricane (cat. 1-5) and major hurricane (cat. 4-5) strength the CAM5 simulations and IBTrACS observations for the time period of 1980 to 2005.

Simulation	Total Storms	Hurricanes	Major Hurricanes
IBTrACS	91.6 ± 8.5	47.1 ± 5.5	10.7 ± 3.3
ne120_gx1v6	70.1 ± 9.0	55.5 ± 7.7	12.5 ± 3.4
ne120_ne120	73.2 ± 10.5	50.3 ± 8.2	4.2 ± 1.9




Figure 3. Storm minimum surface pressure vs. maximum wind speed relationship with quadratic least squares fit (solid lines) for the CAM5 simulations and IBTrACS observations from 1980 to 2005. Note that 3-hourly output is used for the model simulations, while the IBTrACS data is 6-hourly.

Both simulations produce roughly the same frequency of total storms. However, the simulation coupled to the lower resolution ocean produces 10% more hurricanes and nearly three times the amount of major hurricanes when compared to the simulation with the higher resolution coupling.

115 This signifies a shift towards high intensities in the ne120_gx1v6 configuration.

To explore this further, Figure 3 displays the minimum surface pressure vs. maximum wind speed relationship for tropical cyclones for each simulation with a quadratic least squares fit shown as a solid line. IBTrACS observations are again included as a reference and to be consistent with the TC tracker only storms that reach tropical storm strength in their lifetime are used. At low wind speeds
 120 (i.e., <40 m/s) the relationship between the minimum surface pressure and maximum wind speed for the two model simulations and observations compare well. However, at larger wind speeds the relationship between the two simulations diverges consistent to the differences in TC counts in Table 1. In particular, the ne120_gx1v6 simulation produces greater winds speeds at a given minimum pressure than ne120_ne120 simulation, suggesting an impact of the ocean coupling resolution on
 125 tropical cyclone intensity.




Figure 4. Number of TC surface wind instances exceeding category 4 and category 5 wind thresholds for AMIP simulation coupled to 1° ocean grid (ne120_gx1v6) and 0.25° ocean grid (ne120_ne120). Instances are calculated using 3-hourly data.

Figure 4 shows the number of annual 60-m wind exceedances in the 6-hourly model output for both category 4 and category 5 storm thresholds. These represent the two most intense classifications of tropical cyclones, with maximum sustained winds surpassing 59 m s^{-1} and 70 m s^{-1} , respectively. While these are wind thresholds associated with intense TCs they are calculated over the entire ocean domain from 30°S to 30°N and are therefore not necessarily only representative of TCs in the model (is this the correct description?). The bolder, black curve indicates the number of data points surpassing each threshold for the simulation using the 1° ocean/ice grid (ne120_gx1v6) while the red curve marked by crosses represents the same for the simulation with the 0.25° ocean/ice grid (ne120_ne120). From the left panel, we see that for all years (except 1985), the simulations coupled to the coarser ocean grid produces a significantly greater frequency of category 4 level winds. This behavior is even more pronounced in the right panel, where the ne120_gx1v6 simulations averages approximately 5 instances of category 5 level winds per year. However, this threshold is not exceeded at any point during the 26-year sample in the ne120_ne120 simulation.

4 Deterministic simulations

Since all aspects of the model configurations in the climate simulations are identical except for the grid on which the prescribed SSTs and ice concentrations are passed to the other model components, we hypothesize that the marked difference in TC climatology is induced by the coupling strategy and difference in grid resolutions. To assess the differences in simulated TCs in a controlled, deterministic manner, we utilize two identical CAM setups to complete short-term forecast simulations of observed storms. These simulations utilize the new, variable-resolution capability of CAM-SE (Zarzycki et al., 2014b).

The setup is similar to that used in the previous section, but the model is configured with a variable-resolution atmospheric grid with $1/8^\circ$ ($\sim 14\text{km}$) resolution over the Atlantic Ocean. Forecast simulations are initialized with a digitally-filtered atmospheric analysis from the National Center for Atmospheric Predictions's Global Data Assimilation System (GDAS). Observed SSTs are taken from NOAAOI and provided as input to the model on a $1^\circ \times 1^\circ$ grid. The land surface is modeled by the Community Land Model (CLM) version 4.0 and is initialized with a state nudged to be in balance with the atmospheric initial conditions. The model setup and initialization are both further detailed in Zarzycki and Jablonowski (2015).

As in the climate simulations, the only difference between the two setups is the grid used by the data ocean and ice models. The first set of simulations uses the aforementioned displaced tripole grid with an equivalent resolution of 1° (gx1v6) while the second uses an ocean grid identical to the atmospheric grid with an equivalent resolution of $1/8^\circ$ (ne240). Since the SST and ice cover data are provided at coarser scales than the model interpolates to, any differences in the results arise due to the differences in calculating of surface fluxes and momentum drag on the corresponding ocean grids.

After initialization, each configuration is integrated for 8 days. Figure 5 shows the 120-hour forecasts for Hurricane Leslie in the North Atlantic Ocean from the 2012 hurricane season. The simulation was initialized at 00Z on August 31st, 2012 making Fig. 5 valid at 00Z on September 5th, 2012. The forecast using the 1° ocean grid is on the left (Fig. 5a,c), with the $1/8^\circ$ ocean grid on the right (Fig. 5b,d). All fields are extracted from the atmospheric model component. The top panels depict instantaneous lowest model level wind (black vectors) as well as the surface frictional stress vector (red). In the Fig. 5a, it is readily apparent that many instances exist where the vectors are not aligned. This results from the surface stress being calculated on the coarser grid. The atmospheric dynamical core then subsamples this coarser information to provide stress information at the same resolution used by the numerics (as in Fig. 1). In Fig. 5b, the wind and stress vectors are parallel (180° difference), indicating that the frictional drag is acting in direct opposition to the wind within the atmospheric dynamical core. Therefore, the higher resolution ocean grid preserves the resolution of the surface wind field during stress calculations. Because of this, not only are the stress vectors properly aligned with the high-resolution ocean grid, the maximum magnitudes of the stress vectors are larger at the storm's radius of maximum wind in Fig. 5b. This highlights that maxima in the stress at the atmospheric grid cell scale are preserved with the higher resolution ocean grid, whereas these maxima are "smoothed" in the calculation where wind is first averaged to the coarser ocean grid (Fig. 5a), leading to disparate forcing on wind speeds when passed back to the atmospheric model. This is further evidenced by the fact that the integrated dot product (over a $5^\circ \times 5^\circ$ domain centered over the TC minimum surface pressure) of the two fields is approximately 10% smaller in the simulations using the 1° ocean grid. Therefore, the use of the coarser ocean grid results in a universal weaker frictional force used by the atmospheric dynamics, which enhances extreme wind speeds.




Figure 5. 120-hour CAM-SE forecast for Hurricane Leslie, valid at 00Z on September 5th, 2012. Left panels (a,c) are results from forecast using 1° ocean grid. Right panels (b,d) show results using 0.125° ocean grid. Top panels (a-b) display instantaneous wind in the lowest model (black vectors) with corresponding lowest model level wind stress (red vectors). Lower panels (c-d) show total surface flux (latent plus sensible heat).

The cumulative surface heat flux is shown on the bottom (sensible plus latent) for the two storms
185 at the same forecast time. It is readily apparent that the coarser ocean grid (Fig. 5c) provides information back to the atmosphere with significantly less spatial structure than the $1/8^\circ$ ocean grid (Fig. 5d). While the difference in $5^\circ \times 5^\circ$ integrated heat flux is relatively small (approximately 1%), it is clear that the spatial structure of the heat flux field is very different between the two model configurations. This may further play a role in storm dynamics, with the 1° ocean grid providing a larger,
190 more diffuse source of surface heating to the TC core than the high-resolution grid.

5 Conclusions

This manuscript describes biases in extreme climatology which arise from choice of ocean grid and coupling strategy in CESM. Since surface stress and flux calculations are carried out on the ocean

grid, running the model with a coarser ocean than atmosphere presents problems with respect to
195 tropical cyclone (TC) climatology. In particular, surface stress vectors which are passed back to the
atmospheric dynamical core following coupling are not aligned with the surface wind due to being
computed on a coarser grid. This allows winds near the core of TCs to become stronger than if
the stresses were computed at the same resolution of the atmosphere. Additionally, when surface
fluxes are calculated on a different grid, the influx of heat and moisture to the lowest levels of the
200 atmosphere underneath the TC are structurally different, with these quantities being more diffuse
and misaligned with the maximum surface wind, in contradiction to bulk aerodynamical flux theory.

The issues outlined in the manuscript are rather trivial to correct for data ocean models, where
specifying SSTs and applying coupled atmosphere-ocean calculations on the same grid is straight-
forward and computationally ‘cheap.’ More problematic adjustments arise when coupling to a dy-
205 namical ocean model. The vast majority of coupled simulations not only utilize differing resolutions
between different model components, but also different numerical techniques and grids. Therefore,
remapping between components is, in many cases, an absolute necessity.

The obvious recommendation to alleviate coupling inconsistencies when it is not feasible to use
identical grids is to calculate these quantities on the finest resolution grid of the coupled system. In
210 the vast majority of earth system models, this is typically the atmosphere. Performing coupling in
this manner ensures that information passed back to a model component has not be interpolated to
a resolution coarser than the component’s native resolution during the coupling process. However,
in addition to increasing computational cost, it is not clear that technique is fully appropriate for
dynamical ocean models, where aspects such as turbulent mixing may be sufficiently non-linear that
215 merely averaging from a higher resolution grid is not the most appropriate mechanism. Further work
is required to determine whether or not this is the case.

Our results demonstrate that the mean climatology of the simulations presented here are essen-
tially identical regardless of coupling strategy, highlighting that this impact only becomes readily
apparent in the tail of the distributions of interest. However, with climate models being used more
220 and more frequently for direct analysis of TCs, as well as other extreme events, both in present
climate and under future scenarios, this impact on model-derived extremes may become more preva-
lent. This is especially relevant as models continue to march forward with respect to horizontal
resolution, and therefore, their ability to dynamically resolve atmospheric phenomena at smaller and
smaller spatial scales. Consideration of these impacts when utilizing high-resolution climate data for
225 analysis is required and modifications to how the current generation of atmospheric models treats
coupling between various earth system components may be necessary.

Acknowledgements. The National Center for Atmospheric Research is sponsored by the National Science Foun-
dation. Resources of the Argonne Leadership Computing Facility at Argonne National Laboratory, which is
supported by the Office of Science of the U.S. Department of Energy under contract DE-AC02-06CH11357,

230 were used for this research. Furthermore, this work utilizes part of the "Using Petascale Computing Capabilities
to Address Climate Change Uncertainties" PRAC allocation support by the National Science Foundation ACI-
1036146. This work is also part of the Blue Waters sustained-petascale computing project, which is supported
by the National Science Foundation (awards OCI-0725070 and ACI-1238993) and the state of Illinois. Blue
Waters is a joint effort of the University of Illinois at Urbana-Champaign and its National Center for Supercom-
235 puting Applications. Bacmeister was partially supported through the Scientific Discovery through Advanced
Computing (SciDAC) program funded by U.S. Department of Energy, Office of Science, Advanced Scientific
Computing Research. Bates was supported by the Regional and Global Climate Modeling Program (RGCM) of
the US. Department of Energy, Office of Science (BER), Cooperative Agreement DE-FC02-97ER62402.

References

- 240 Bacmeister, J. T., Wehner, M. F., Neale, R. B., Gettelman, A., Hannay, C., Lauritzen, P. H., Caron, J. M.,
and Truesdale, J. E.: Exploratory High-Resolution Climate Simulations using the Community Atmosphere
Model (CAM), *J. Climate*, 27, 3073–3099, doi:10.1175/JCLI-D-13-00387.1, 2014.
- Bengtsson, L., Hodges, K. I., and Esch, M.: Tropical cyclones in a T159 resolution global climate model: com-
parison with observations and re-analyses, *Tellus A*, 59, 396–416, doi:10.1111/j.1600-0870.2007.00236.x,
245 2007.
- Camargo, S. J.: Global and Regional Aspects of Tropical Cyclone Activity in the CMIP5 Models, *Journal of
Climate*, 26, 9880–9902, doi:10.1175/JCLI-D-12-00549.1, 2013.
- Craig, A. P., Vertenstein, M., and Jacob, R.: A new flexible coupler for earth system modeling developed
for CCSM4 and CESM1, *International Journal of High Performance Computing Applications*, 26, 31–42,
250 doi:10.1177/1094342011428141, 2012.
- Dennis, J. M., Edwards, J., Evans, K. J., Guba, O., Lauritzen, P. H., Mirin, A. A., St-Cyr, A., Tay-
lor, M. A., and Worley, P. H.: CAM-SE: A scalable spectral element dynamical core for the Commu-
nity Atmosphere Model, *International Journal of High Performance Computing Applications*, 26, 74–89,
doi:10.1177/1094342011428142, 2012.
- 255 Evans, K. J., Lauritzen, P. H., Mishra, S. K., Neale, R. B., Taylor, M. A., and Tribbia, J. J.: AMIP Simulation
with the CAM4 Spectral Element Dynamical Core, *Journal of Climate*, 26, 689–709, doi:10.1175/JCLI-D-
11-00448.1, 2013.
- Gates, W. L.: AMIP: The Atmospheric Model Intercomparison Project, *Bulletin of the American Meteorologi-
cal Society*, 73, 1962–1970, 1992.
- 260 Huffman, G. J., Adler, R. F., Bolvin, D. T., Gu, G., Nelkin, E. J., Bowman, K. P., Hong, Y., Stocker, E. F., and
Wolff, D. B.: The TRMM Multisatellite Precipitation Analysis (TMPA): Quasi-global, multiyear, combined-
sensor precipitation estimates at fine scales., *Journal of Hydrometeorology*, 8, 2007.
- Hurrell, J. W., Holland, M. M., Gent, P. R., Ghan, S., Kay, J. E., Kushner, P. J., Lamarque, J. F., Large, W. G.,
Lawrence, D., Lindsay, K., Lipscomb, W. H., Long, M. C., Mahowald, N., Marsh, D. R., Neale, R. B.,
265 Rasch, P., Vavrus, S., Vertenstein, M., Bader, D., Collins, W. D., Hack, J. J., Kiehl, J., and Marshall, S.:
The Community Earth System Model: A Framework for Collaborative Research, *Bulletin of the American
Meteorological Society*, 94, 1339–1360, doi:10.1175/BAMS-D-12-00121.1, 2013.
- Kalnay, E., Kanamitsu, M., Kistler, R., Collins, W., Deaven, D., Gandin, L., Iredell, M., Saha, S., White,
G., Woollen, J., Zhu, Y., Leetmaa, A., Reynolds, R., Chelliah, M., Ebisuzaki, W., Higgins, W., Janowiak,
270 J., Mo, K. C., Ropelewski, C., Wang, J., Jenne, R., and Joseph, D.: The NCEP/NCAR 40-Year Re-
analysis Project, *Bulletin of the American Meteorological Society*, 77, 437–471, doi:10.1175/1520-
0477(1996)077<0437:TNYRP>2.0.CO;2, 1996.
- Kim, D., Sobel, A. H., Del Genio, A. D., Chen, Y., Camargo, S. J., Yao, M.-S., Kelley, M., and Nazarenko, L.:
The Tropical Subseasonal Variability Simulated in the NASA GISS General Circulation Model, *J. Climate*,
275 25, 4641–4659, doi:10.1175/JCLI-D-11-00447.1, 2012.
- Knapp, K. R., Kruk, M. C., Levinson, D. H., Diamond, H. J., and Neumann, C. J.: The Interna-
tional Best Track Archive for Climate Stewardship (IBTrACS), *Bull. Amer. Meteor. Soc.*, 91, 363–376,
doi:10.1175/2009BAMS2755.1, 2010.

- Knutson, T. R., Sirutis, J. J., Garner, S. T., Held, I. M., and Tuleya, R. E.: Simulation of the Recent Multidecadal
 280 Increase of Atlantic Hurricane Activity Using an 18-km-Grid Regional Model, *Bulletin of the American Meteorological Society*, 88, 1549, doi:10.1175/BAMS-88-10-1549, 2007.
- Knutson, T. R., McBride, J. L., Chan, J., Emanuel, K., Holland, G., Landsea, C., Held, I., Kossin, J. P.,
 Srivastava, A. K., and Sugi, M.: Tropical cyclones and climate change, *Nature Geosci.*, 3, 157–163,
 doi:10.1038/ngeo779, 2010.
- 285 Lim, Y.-K., Schubert, S. D., Reale, O., Lee, M.-I., Molod, A. M., and Suarez, M. J.: Sensitivity of Tropical
 Cyclones to Parameterized Convection in the NASA GEOS-5 Model, *Journal of Climate*, 28, 551–573,
 doi:10.1175/JCLI-D-14-00104.1, 2014.
- Manganello, J. V., Hodges, K. I., Kinter, J. L., Cash, B. A., Marx, L., Jung, T., Achuthavarier, D., Adams, J. M.,
 Altshuler, E. L., Huang, B., Jin, E. K., Stan, C., Towers, P., and Wedi, N.: Tropical Cyclone Climatology in
 290 a 10-km Global Atmospheric GCM: Toward Weather-Resolving Climate Modeling, *Journal of Climate*, 25,
 3867–3893, doi:10.1175/JCLI-D-11-00346.1, 2012.
- Murakami, H., Wang, Y., Yoshimura, H., Mizuta, R., Sugi, M., Shindo, E., Adachi, Y., Yukimoto, S., Hosaka,
 M., Kusunoki, S., Ose, T., and Kitoh, A.: Future Changes in Tropical Cyclone Activity Projected by the New
 High-Resolution MRI-AGCM, *Journal of Climate*, 25, 3237–3260, doi:10.1175/JCLI-D-11-00415.1, 2012.
- 295 Neale, R. B., Chen, C.-C., Gettelman, A., Lauritzen, P. H., Park, S., Williamson, D. L., Conley, A. J., Garcia, R.,
 Kinnison, D., Lamarque, J.-F., Marsh, D., Mills, M., Smith, A. K., Tilmes, S., Vitt, F., Cameron-Smith, P.,
 Collins, W. D., Iacono, M. J., , Easter, R. C., Liu, X., Ghan, S. J., Rasch, P. J., and Taylor, M. A.: Description
 of the NCAR Community Atmosphere Model (CAM 5.0), NCAR Technical Note NCAR/TN-486+STR,
 National Center for Atmospheric Research, Boulder, Colorado, 2010.
- 300 Oleson, K., Lawrence, D., Bonan, G., Flanner, M., Kluzek, E., Lawrence, P., Levis, S., Swenson, S., Thorn-
 ton, P., Dai, A., Decker, M., Dickinson, R., Feddema, J., Heald, C., Hoffman, F., Lamarque, J., Mahowald,
 N., Niu, G., Qian, T., Randerson, J., Running, S., Sakaguchi, K., Slater, A., Stockli, R., Wang, A., Yang,
 Z., Zeng, X., and Zeng, X.: Technical Description of version 4.0 of the Community Land Model (CLM),
 NCAR Technical Note NCAR/TN-478+STR, National Center for Atmospheric Research, Boulder, Colorado,
 305 doi:10.5065/D6FB50WZ, 2010.
- Oouchi, K., Yoshimura, J., Yoshimura, H., Mizuta, R., Kusunoki, S., and Noda, A.: Tropical Cyclone Clima-
 tology in a Global-Warming Climate as Simulated in a 20 km-Mesh Global Atmospheric Model: Frequency
 and Wind Intensity Analyses, *Journal of the Meteorological Society of Japan*, 84, 259–276, 2006.
- Reed, K. A. and Jablonowski, C.: Impact of physical parameterizations on idealized tropical cyclones in the
 310 Community Atmosphere Model, *Geophysical Research Letters*, 38, doi:10.1029/2010GL046297, 2011.
- Reed, K. A. and Jablonowski, C.: Assessing the Uncertainty of Tropical Cyclone Simulations in NCAR’s Com-
 munity Atmosphere Model, *J. Adv. Model. Earth Syst.*, 3, M08 002, doi:10.1029/2011MS000076, 2011.
- Reed, K. A. and Jablonowski, C.: Idealized tropical cyclone simulations of intermediate complexity: A test case
 for AGCMs, *Journal of Advances in Modeling Earth Systems*, 4, M04 001, doi:10.1029/2011MS000099,
 315 2012.
- Reed, K. A., Bacmeister, J. T., Wehner, Rosenbloom, N. A., F., M., Bates, S. C., Lauritzen, P. H., Truesdale,
 J. T., and Hannay, C.: Impact of the dynamical core on the direct simulation of tropical cyclones in a high-
 resolution global model, *Geophys. Res. Lett.*, 42, doi:10.1002/2015GL063974, in press, 2015.

- Rienecker, M. M., Suarez, M. J., Gelaro, R., Todling, R., Bacmeister, J., Liu, E., Bosilovich, M. G., Schubert, S. D., Takacs, L., Kim, G.-K., Bloom, S., Chen, J., Collins, D., Conaty, A., da Silva, A., Gu, W., Joiner, J., Koster, R. D., Lucchesi, R., Molod, A., Owens, T., Pawson, S., Pegion, P., Redder, C. R., Reichle, R., Robertson, F. R., Ruddick, A. G., Sienkiewicz, M., and Woollen, J.: MERRA: NASA's Modern-Era Retrospective Analysis for Research and Applications, *Journal of Climate*, 24, 3624–3648, doi:10.1175/JCLI-D-11-00015.1, 2011.
- Satoh, M., Oouchi, K., Nasuno, T., Taniguchi, H., Yamada, Y., Tomita, H., Kodama, C., Kinter, J., Achuthavari, D., Manganello, J., Cash, B., Jung, T., Palmer, T., and Wedi, N.: The Intra-Seasonal Oscillation and its control of tropical cyclones simulated by high-resolution global atmospheric models, *Climate Dynamics*, 39, 2185–2206, doi:10.1007/s00382-011-1235-6, 2012.
- Simpson, R. H.: The Hurricane Disaster – Potential Scale, *Weatherwise*, 27, 169–186, doi:10.1080/00431672.1974.9931702, 1974.
- Strachan, J., Vidale, P. L., Hodges, K., Roberts, M., and Demory, M.-E.: Investigating Global Tropical Cyclone Activity with a Hierarchy of AGCMs: The Role of Model Resolution, *Journal of Climate*, 26, 133–152, doi:10.1175/JCLI-D-12-00012.1, 2013.
- Taylor, K. E.: Summarizing multiple aspects of model performance in a single diagram, *Journal of Geophysical Research: Atmospheres*, 106, 7183–7192, doi:10.1029/2000JD900719, 2001.
- Taylor, M., Tribbia, J., and Iskandarani, M.: The Spectral Element Method for the Shallow Water Equations on the Sphere, *Journal of Computational Physics*, 130, 92–108, 1997.
- Taylor, M. A.: Conservation of Mass and Energy for the Moist Atmospheric Primitive Equations on Unstructured Grids, in: *Numerical Techniques for Global Atmospheric Models*, edited by Lauritzen, P. H., Jablonowski, C., Taylor, M. A., and Nair, R. D., vol. 80 of *Lecture Notes in Computational Science and Engineering*, pp. 357–380, Springer, 2011.
- Taylor, M. A. and Fournier, A.: A compatible and conservative spectral element method on unstructured grids, *Journal of Computational Physics*, 229, 5879–5895, 2010.
- Thomas, S. and Loft, R.: The NCAR Spectral Element Climate Dynamical Core: Semi-Implicit Eulerian Formulation, *Journal of Scientific Computing*, 25, 307–322, doi:10.1007/s10915-004-4646-2, 2005.
- Vitart, F., Anderson, J. L., and Stern, W. F.: Simulation of interannual variability of tropical storm frequency in an ensemble of GCM integrations, *Journal of Climate*, 10, 745–760, 1997.
- Walsh, K. J. E., Camargo, S. J., Vecchi, G. A., Daloz, A. S., Elsner, J., Emanuel, K., Horn, M., Lim, Y.-K., Roberts, M., Patricola, C., Scoccimarro, E., Sobel, A. H., Strazzo, S., Villarini, G., Wehner, M., Zhao, M., Kossin, J. P., LaRow, T., Oouchi, K., Schubert, S., Wang, H., Bacmeister, J., Chang, P., Chauvin, F., Jablonowski, C., Kumar, A., Murakami, H., Ose, T., Reed, K. A., Saravanan, R., Yamada, Y., Zarzycki, C. M., Vidale, P. L., Jonas, J. A., and Henderson, N.: Hurricanes and climate: the U.S. CLIVAR working group on hurricanes, *Bulletin of the American Meteorological Society*, doi:10.1175/BAMS-D-13-00242.1, 2015.
- Wehner, M. F., Reed, K. A., Li, F., Prabhat, Bacmeister, J., Chen, C.-T., Paciorek, C., Gleckler, P. J., Sperber, K. R., Collins, W. D., Gettelman, A., and Jablonowski, C.: The effect of horizontal resolution on simulation quality in the Community Atmospheric Model, CAM5.1, *Journal of Advances in Modeling Earth Systems*, doi:10.1002/2013MS000276, 2014.

- Wehner, M. F., Prabhat, Reed, K. A., Stone, D., Collins, W. D., and Bacmeister, J. T.: Resolution dependence
 360 of future tropical cyclone projections of CAM5.1 in the US CLIVAR Hurricane Working Group idealized
 configurations, *J. Climate*, doi:10.1175/JCLI-D-14-00311.1, in press, 2015.
- Zarzycki, C. M. and Jablonowski, C.: A multidecadal simulation of Atlantic tropical cyclones using a variable-
 resolution global atmospheric general circulation model, *Journal of Advances in Modeling Earth Systems*,
 6, 805–828, doi:10.1002/2014MS000352, 2014.
- 365 Zarzycki, C. M. and Jablonowski, C.: Experimental tropical cyclone forecasts using a variable-resolution global
 model, *Monthly Weather Review*, in prep., 2015.
- Zarzycki, C. M., Jablonowski, C., and Taylor, M. A.: Using Variable Resolution Meshes to Model
 Tropical Cyclones in the Community Atmosphere Model, *Monthly Weather Review*, 142, 1221–1239,
 doi:10.1175/MWR-D-13-00179.1, 2014a.
- 370 Zarzycki, C. M., Levy, M. N., Jablonowski, C., Overfelt, J. R., Taylor, M. A., and Ullrich, P. A.: Aqua-
 planet Experiments Using CAM’s Variable-Resolution Dynamical Core, *Journal of Climate*, 27, 5481–5503,
 doi:10.1175/JCLI-D-14-00004.1, 2014b.
- Zarzycki, C. M., Jablonowski, C., Thatcher, D. R., and Taylor, M. A.: Effects of localized grid refinement
 on the general circulation and climatology in the Community Atmosphere Model, *Journal of Climate*, 28,
 375 2777–2803, doi:10.1175/JCLI-D-14-00599.1, 2015.
- Zhao, M., Held, I. M., Lin, S. J., and Vecchi, G. A.: Simulations of Global Hurricane Climatology, Interannual
 Variability, and Response to Global Warming Using a 50-km Resolution GCM, *Journal of Climate*, 22,
 6653–6678, 2009.
- Zhao, M., Held, I. M., and Lin, S.-J.: Some Counterintuitive Dependencies of Tropical Cyclone Frequency on
 380 Parameters in a GCM, *Journal of the Atmospheric Sciences*, 69, 2272–2283, doi:10.1175/JAS-D-11-0238.1,
 2012.