

Title of the article

John Smith,¹ James Smith,^{1,2}, and Jane Smith²

Corresponding author: Jane Smith, Department of Geography, Ohio State University, Columbus, Ohio, USA. (j.smith@ohio.edu)

¹Department of Hydrology and Water
Resources, University of Arizona, Tucson,
Arizona, USA.

²Department of Geography, Ohio State
University, Columbus, Ohio, USA.

² **Abstract.**

³ The abstract goes here.

1. Introduction

As horizontal resolution in general circulation models (GCMs) increases, the representation of tropical cyclones has improved greatly [*Oouchi et al.*, 2006; *Bengtsson et al.*, 2007; *Zhao et al.*, 2009; *Murakami et al.*, 2012; *Manganello et al.*, 2012; *Satoh et al.*, 2012; *Strachan et al.*, 2013; *Zarzycki and Jablonowski*, 2014; *Wehner et al.*, 2014].

2. Model description

2.1. Community Atmosphere Model

The Community Atmosphere Model (CAM)

In this paper, all simulations utilize the Spectral Element (SE) dynamical core option within CAM.

2.2. Coupling within CESM

(Describe coupling procedure in CESM; draw schematic; confirm with T. Craig or Mariana).

Prescribed SSTs and ice are passed to the model on a $1^\circ \times 1^\circ$ grid and internally interpolated to the ocean and ice grids.

3. Results

3.1. Deterministic simulations

To assess the differences in simulated TCs in a controlled manner, we utilize two nearly identical CAM setups to forecast observed storms. These simulations utilize the new, variable-resolution capability of CAM-SE [*Zarzycki et al.*, 2014].

grid with $1/8^\circ$ ($\sim 14\text{km}$) grid spacing over the Atlantic Ocean and are initialized with atmospheric analysis from the.

Observed SSTs are taken from NOAAOI and provided as input to the model on a $1^\circ \times 1^\circ$ grid.

Further details about model setup and initialization are described in ?.

The only differences between the two model setups is the grid used by the data ocean and ice models. The first set of simulations uses a displaced tripole grid with an equivalent resolution of 1° (gx1v6) while the second uses an ocean grid identical to the atmospheric grid with an equivalent resolution of $1/8^\circ$.

COLIN FORECAST STUFF AT 14 KM

3.2. Climate simulations

JULIO/KEVIN CLIMATE SIMS

4. Discussion

Discussion goes here.

Acknowledgments. This work was partially supported by a grant from the Spanish Ministry of Science and Technology.

References

Bengtsson, L., K. I. Hodges, and M. Esch (2007), Tropical cyclones in a T159 resolution global climate model: comparison with observations and re-analyses, *Tellus A*, 59(4), 396–416, doi:10.1111/j.1600-0870.2007.00236.x.

Manganello, J. V., K. I. Hodges, J. L. Kinter, B. A. Cash, L. Marx, T. Jung,
D. Achuthavarier, J. M. Adams, E. L. Altshuler, B. Huang, E. K. Jin, C. Stan, P. Tow-
ers, and N. Wedi (2012), Tropical cyclone climatology in a 10-km global atmospheric
GCM: Toward weather-resolving climate modeling, *Journal of Climate*, *25*(11), 3867–
3893, doi:10.1175/JCLI-D-11-00346.1.

Murakami, H., Y. Wang, H. Yoshimura, R. Mizuta, M. Sugi, E. Shindo, Y. Adachi,
S. Yukimoto, M. Hosaka, S. Kusunoki, T. Ose, and A. Kitoh (2012), Future changes in
tropical cyclone activity projected by the new high-resolution MRI-AGCM, *Journal of*
Climate, *25*(9), 3237–3260, doi:10.1175/JCLI-D-11-00415.1.

Oouchi, K., J. Yoshimura, H. Yoshimura, R. Mizuta, S. Kusunoki, and A. Noda (2006),
Tropical cyclone climatology in a global-warming climate as simulated in a 20 km-
mesh global atmospheric model: Frequency and wind intensity analyses, *Journal of the*
Meteorological Society of Japan, *84*(2), 259–276.

Satoh, M., K. Oouchi, T. Nasuno, H. Taniguchi, Y. Yamada, H. Tomita, C. Kodama,
J. Kinter, D. Achuthavarier, J. Manganello, B. Cash, T. Jung, T. Palmer, and N. Wedi
(2012), The intra-seasonal oscillation and its control of tropical cyclones simulated by
high-resolution global atmospheric models, *Climate Dynamics*, *39*(9-10), 2185–2206,
doi:10.1007/s00382-011-1235-6.

Strachan, J., P. L. Vidale, K. Hodges, M. Roberts, and M.-E. Demory (2013), Investi-
gating global tropical cyclone activity with a hierarchy of AGCMs: The role of model
resolution, *Journal of Climate*, *26*(1), 133–152, doi:10.1175/JCLI-D-12-00012.1.

Wehner, M. F., K. A. Reed, F. Li, Prabhat, J. Bacmeister, C.-T. Chen, C. Paciorek,
P. J. Gleckler, K. R. Sperber, W. D. Collins, A. Gettelman, and C. Jablonowski

(2014), The effect of horizontal resolution on simulation quality in the Community Atmospheric Model, CAM5.1, *Journal of Advances in Modeling Earth Systems*, doi:10.1002/2013MS000276.

Zarzycki, C. M., and C. Jablonowski (2014), A multidecadal simulation of Atlantic tropical cyclones using a variable-resolution global atmospheric general circulation model, *Journal of Advances in Modeling Earth Systems*, 6(3), 805–828, doi:10.1002/2014MS000352.

Zarzycki, C. M., M. N. Levy, C. Jablonowski, J. R. Overfelt, M. A. Taylor, and P. A. Ullrich (2014), Aquaplanet experiments using CAM’s variable-resolution dynamical core, *Journal of Climate*, 27, 5481–5503, doi:10.1175/JCLI-D-14-00004.1.

Zhao, M., I. M. Held, S. J. Lin, and G. A. Vecchi (2009), Simulations of global hurricane climatology, interannual variability, and response to global warming using a 50-km resolution GCM, *Journal of Climate*, 22(24), 6653–6678.



Figure 1. Figure caption

Table 1. Table caption

| Treatments | Response 1 | Response 2 |
|-------------------|-------------------|-------------------|
| Treatment 1 | 0.0003262 | 0.562 |
| Treatment 2 | 0.0015681 | 0.910 |
| Treatment 3 | 0.0009271 | 0.296 |