

Table S2. GBD risk hierarchy with levels	
Risk	level
All risk factors	0
Environmental/occupational risks	1
Unsafe water, sanitation, and handwashing	2
Unsafe water source	3
Unsafe sanitation	3
No access to handwashing facility	3
Air pollution	2
Particulate matter pollution	3
Ambient particulate matter pollution	4
Household air pollution from solid fuels	4
Ambient ozone pollution	3
Non-optimal temperature	2
High temperature	3
Low temperature	3
Other environmental risks	2
Residential radon	3
Lead exposure	3
Occupational risks	2
Occupational carcinogens	3
Occupational exposure to asbestos	4
Occupational exposure to arsenic	4
Occupational exposure to benzene	4
Occupational exposure to beryllium	4
Occupational exposure to cadmium	4
Occupational exposure to chromium	4
Occupational exposure to diesel engine exhaust	4
Occupational exposure to formaldehyde	4
Occupational exposure to nickel	4
Occupational exposure to polycyclic aromatic hydrocarbons	4
Occupational exposure to silica	4
Occupational exposure to sulfuric acid	4
Occupational exposure to trichloroethylene	4
Occupational asthmagens	3
Occupational particulate matter, gases, and fumes	3
Occupational noise	3
Occupational injuries	3
Occupational ergonomic factors	3
Behavioral risks	1
Child and maternal malnutrition	2
Suboptimal breastfeeding	3
Non-exclusive breastfeeding	4
Discontinued breastfeeding	4
Child growth failure	3

Child underweight	4
Child wasting	4
Child stunting	4
Low birthweight and short gestation	3
Short gestation	4
Low birthweight	4
Iron deficiency	3
Vitamin A deficiency	3
Zinc deficiency	3
Tobacco	2
Smoking	3
Chewing tobacco	3
Secondhand smoke	3
Alcohol use	2
Drug use	2
Dietary risks	2
Diet low in fruits	3
Diet low in vegetables	3
Diet low in legumes	3
Diet low in whole grains	3
Diet low in nuts and seeds	3
Diet low in milk	3
Diet high in red meat	3
Diet high in processed meat	3
Diet high in sugar-sweetened beverages	3
Diet low in fiber	3
Diet low in calcium	3
Diet low in seafood omega-3 fatty acids	3
Diet low in polyunsaturated fatty acids	3
Diet high in trans fatty acids	3
Diet high in sodium	3
Intimate partner violence	2
Childhood sexual abuse and bullying	2
Childhood sexual abuse	3
Bullying victimisation	3
Unsafe sex	2
Low physical activity	2
Metabolic risks	1
High fasting plasma glucose	2
High LDL cholesterol	2
High systolic blood pressure	2
High body-mass index	2
Low bone mineral density	2
Kidney dysfunction	2