Representations of Compact Groups

Lecture notes

(based on the *Representations of Finite and Compact Groups*)

Zayn

Contents

Conventions and prerequisites **3**

1 Topological groups and their representations **3**
1.1 Topological groups . . . 3
1.2 Representations . . . 4
1.3 Schur's lemma . . . 5

2 Haar measure and complete reducibility **5**
2.1 Existence and uniqueness of Haar measure . . . 5
2.2 Unitarity and complete reducibility . . . 7

3 Schur orthogonality and characters **8**
3.1 Matrix coefficients and the orthogonality relations . . . 8
3.2 Characters . . . 9
3.3 First example: the representations of SU(2) . . . 9

4 The Peter–Weyl theorem **11**
4.1 Compact operators and the regular representation . . . 11
4.2 The Peter–Weyl theorem . . . 12
4.3 Consequences . . . 13

5 Lie groups and Lie algebras **14**
5.1 The matrix exponential . . . 14
5.2 The Lie algebra of a matrix Lie group . . . 15
5.3 Homomorphisms and derived representations . . . 16
5.4 The representations of sl(2,C); SU(2) and SO(3) . . . 17
5.5 Compact Lie groups are matrix groups . . . 18

6 Compact Lie algebras and the exponential map **18**
6.1 Invariant inner products . . . 18
6.2 The differential of the exponential map . . . 19

7 Maximal tori **19**
7.1 Tori . . . 19
7.2 Maximal tori and the Weyl group . . . 20
7.3 The torus theorem . . . 21
7.4 Centralizers of tori . . . 23

8 Roots and weights **23**
8.1 Weights of a representation . . . 23
8.2 Roots . . . 24
8.3 The SU(2)-subalgebras attached to roots . . . 24
8.4 Reflections and the root system . . . 26
8.5 Positive roots, simple roots and dominant weights . . . 26

9 The Weyl group **27**
9.1 W as a reflection group . . . 27
9.2 Alternating sums and the Weyl denominator . . . 28
9.3 Regular elements . . . 29
9.4 The Weyl integration formula . . . 29

10 The Weyl character formula **31**

11 Representations of the classical groups **33**
11.1 Root data of the classical groups 33
11.2 The unitary groups 35
11.3 The special unitary groups 37
11.4 The orthogonal groups and spherical harmonics 38
11.5 The compact symplectic groups 39

12 Applications to physics **39**
12.1 Symmetries in quantum mechanics 39
12.2 Angular momentum 39
12.3 The hydrogen atom and SO(4) 40
12.4 Isospin and the eightfold way 41

Conventions and prerequisites

Conventions. All vector spaces are complex unless stated otherwise. Inner products $\langle \cdot, \cdot \rangle$ are linear in the first and antilinear in the second argument (Simon uses the opposite convention). Topological groups are Hausdorff, and representations are continuous. "Representation" means finite-dimensional representation unless stated otherwise. We write e or 1 for the neutral element of a group, $\mathbb{T} := \{z \in \mathbb{C} \mid |z| = 1\}$, and $\mathbb{N}_0 := \mathbb{N} \cup \{0\}$.

Prerequisites. Proofs of the main mathematical results are included. The following prerequisites are used freely; physical inputs and historical remarks are not proved.

(i) *Analysis and topology:* Heine–Borel; Arzelà–Ascoli; Urysohn's lemma and Tietze's extension theorem for compact Hausdorff spaces; the Riesz–Markov representation theorem (positive linear functionals on $C(X)$, X compact Hausdorff, are given by unique regular Borel measures); density of $C(X)$ in $L^2(X, \mu)$ for regular Borel measures; Fubini–Tonelli and monotone convergence; completeness of $\{\sin(m\theta)\}_{m\in\mathbb{N}}$ in $L^2([0, \pi])$; the lifting of continuous maps $\mathbb{R} \to \mathbb{T}$ along $t \mapsto e^{2\pi i t}$; the spectral theorem for unitary, Hermitian and real orthogonal matrices; Schur triangularization; subgroups of $\mathbb{Z}^m$ are free abelian of rank $\leq m$; a holomorphic function on $\mathbb{C}^N$ vanishing on $\mathbb{R}^N$ vanishes identically.

(ii) *Functional analysis:* the spectral theorem for compact self-adjoint operators (Theorem 4.1.1); Hilbert–Schmidt integral operators are compact; norm limits of finite-rank operators are compact; Riesz' lemma for bounded sesquilinear forms.

(iii) *Differential calculus and geometry:* inverse function theorem and submersion theorem (a smooth map whose differential is surjective at a point maps every neighborhood of that point onto a neighborhood of its image); the change of variables formula for diffeomorphisms; volume forms and the associated measures on manifolds, partitions of unity; the image of a manifold of dimension $< n$ under a smooth map into an n-manifold has measure zero; the surface measure of S^{n-1} is invariant under $\mathrm{O}(n)$; for a closed subgroup H of a Lie group G, the quotient G/H is a manifold such that $G \to G/H$ is a smooth submersion and $G \times G/H \to G/H$ is smooth; every Lie group has an exponential map which is a diffeomorphism from a neighborhood of 0 onto a neighborhood of e (only used in Section 5.5).

1 Topological groups and their representations

1.1 Topological groups

Definition 1.1.1: A *topological group* is a group G equipped with a Hausdorff topology such that the maps $G \times G \to G$, $(g, h) \mapsto gh$, and $G \to G$, $g \mapsto g^{-1}$, are continuous. A topological group is *compact* if it is compact as a topological space.

Remark 1.1.2: In a topological group, left translations $g \mapsto hg$, right translations $g \mapsto gh$ and conjugations $g \mapsto hgh^{-1}$ are homeomorphisms (their inverses are of the same type). Hence if U is a neighborhood of e, then hU is a neighborhood of h.

Example 1.1.3: The following are compact groups.

(i) Every finite group with the discrete topology.

(ii) The circle $\mathbb{T} = \mathrm{U}(1)$ and the n-torus $\mathbb{T}^n$.

(iii) The *unitary group* $\mathrm{U}(n) := \{g \in M_n(\mathbb{C}) \mid g^* g = 1\}$ and $\mathrm{SU}(n) := \{g \in \mathrm{U}(n) \mid \det g = 1\}$.

(iv) The *orthogonal group* $\mathrm{O}(n) := \{g \in M_n(\mathbb{R}) \mid g^T g = 1\}$ and $\mathrm{SO}(n) := \{g \in \mathrm{O}(n) \mid \det g = 1\}$.

(v) The *compact symplectic group* $\mathrm{Sp}(n) := \{g \in \mathrm{U}(2n) \mid g^T J g = J\}$, where $J = \left(\begin{smallmatrix} 0 & 1_n \\ -1_n & 0 \end{smallmatrix}\right)$.

(vi) The infinite product $\prod_{n\in\mathbb{N}} \mathbb{Z}/2\mathbb{Z}$ with the product topology (compact by Tikhonov's theorem).

Lemma 1.1.4: The groups $\mathrm{U}(n)$, $\mathrm{SU}(n)$, $\mathrm{O}(n)$, $\mathrm{SO}(n)$ and $\mathrm{Sp}(n)$ are compact topological groups.

Proof. They are subgroups of $\mathrm{GL}(n,\mathbb{C})$ (resp. $\mathrm{GL}(2n,\mathbb{C})$): e.g. for $\mathrm{Sp}(n)$, if $g^T J g = J = h^T J h$ then $(gh)^T J (gh) = h^T J h = J$ and $(g^{-1})^T J g^{-1} = (g^{-1})^T (g^T J g) g^{-1} = J$. They are closed in $M_n(\mathbb{C}) \cong \mathbb{C}^{n^2}$, being defined by polynomial equations in the entries and their conjugates, and bounded, since the columns of a unitary matrix are unit vectors, so $|g_{jk}| \le 1$. By Heine–Borel they are compact. Multiplication is polynomial and inversion $g \mapsto g^{-1} = g^*$ is continuous. □

Lemma 1.1.5 (Uniform continuity): Let G be a compact group and $f \in C(G)$. For every $\varepsilon > 0$ there is a neighborhood U of e such that

$$|f(hg) - f(g)| < \varepsilon \quad \text{and} \quad |f(gh) - f(g)| < \varepsilon \qquad \text{for all } g \in G,\ h \in U.$$

Proof. We prove the first inequality; the second follows by applying the first to $\tilde{f}(g) := f(g^{-1})$ and replacing U by $U \cap U^{-1}$. For each $g \in G$, continuity of $(h, g') \mapsto f(hg')$ at (e, g) gives open neighborhoods U_g of e and V_g of g with $|f(hg') - f(g)| < \varepsilon/2$ for $h \in U_g$, $g' \in V_g$. Cover G by finitely many $V_{g_1}, \dots, V_{g_m}$ and set $U := \bigcap_j U_{g_j}$. If $g \in V_{g_j}$ and $h \in U$, then $|f(hg) - f(g)| \le |f(hg) - f(g_j)| + |f(g_j) - f(g)| < \varepsilon$. □

1.2 Representations

Definition 1.2.1: Let G be a topological group. A *representation* of G is a pair (π, V) of a finite-dimensional complex vector space V and a continuous homomorphism $\pi : G \to \mathrm{GL}(V)$; $d_\pi := \dim V$ is its *dimension*. If V carries an inner product for which all $\pi(g)$ are unitary, π is a *unitary representation*. For a Hilbert space $\mathcal{H}$ of arbitrary dimension, a *unitary representation* is a homomorphism $\pi : G \to \mathrm{U}(\mathcal{H})$ which is *strongly continuous*, i.e., $g \mapsto \pi(g)v$ is continuous for every $v \in \mathcal{H}$.

Definition 1.2.2: Let (π, V) and (σ, W) be representations of G.

(i) A subspace $V' \subset V$ is *invariant* if $\pi(g)V' \subset V'$ for all $g \in G$; then $\pi|_{V'}$ is a *subrepresentation*. (For Hilbert space representations we require invariant subspaces to be closed.)

(ii) π is *irreducible* if $V \neq \{0\}$ and $\{0\}$ and V are the only invariant subspaces.

(iii) A linear map $A : V \to W$ is an *intertwiner* if $A\pi(g) = \sigma(g)A$ for all $g \in G$. The intertwiners form a vector space $\mathrm{Hom}_G(V, W)$.

(iv) π and σ are *equivalent*, $\pi \cong \sigma$, if there is an invertible intertwiner.

We denote by $\widehat{G}$ the set of equivalence classes of irreducible representations of G.

Definition 1.2.3 (Constructions): For representations (π, V), (σ, W) of G we have the *direct sum* $(\pi \oplus \sigma)(g) := \pi(g) \oplus \sigma(g)$, the *tensor product* $(\pi \otimes \sigma)(g) := \pi(g) \otimes \sigma(g)$, the *contragredient representation* π^* on V^* given by $(\pi^*(g)\ell)(v) := \ell(\pi(g)^{-1}v)$, and the representation $g \cdot A := \sigma(g) A \pi(g)^{-1}$ on $\mathrm{Hom}(V, W)$, whose fixed vectors are exactly the intertwiners.

Example 1.2.4: (i) The *trivial representation* $\pi(g) = 1$ on $\mathbb{C}$.

(ii) For $k \in \mathbb{Z}$, $\chi_k(z) := z^k$ is a one-dimensional representation of $\mathbb{T}$.

(iii) The *defining representation* of a closed subgroup $G \subset \mathrm{GL}(n, \mathbb{C})$ on $\mathbb{C}^n$.

(iv) Let V_n be the space of homogeneous polynomials of degree n in $z = (z_1, z_2)$, so $\dim V_n = n + 1$. For $g \in \mathrm{SU}(2)$ put $(\pi_n(g)P)(z) := P(g^{-1}z)$. Then

$$(\pi_n(g)\pi_n(h)P)(z) = (\pi_n(h)P)(g^{-1}z) = P(h^{-1}g^{-1}z) = (\pi_n(gh)P)(z),$$

and continuity is clear since the coefficients of $\pi_n(g)P$ are polynomials in the entries of $g^{-1} = g^*$. So π_n is a representation of $\mathrm{SU}(2)$.

1.3 Schur's lemma

Lemma 1.3.1 (Schur's lemma): Let (π, V) and (σ, W) be irreducible representations of G.

(i) Every $A \in \mathrm{Hom}_G(V, W)$ is either zero or invertible.

(ii) $\mathrm{Hom}_G(V, V) = \mathbb{C} \cdot 1$.

Consequently, $\dim \mathrm{Hom}_G(V, W) = 1$ if $\pi \cong \sigma$, and $\mathrm{Hom}_G(V, W) = 0$ otherwise.

Proof. (i) $\ker A \subset V$ and $\operatorname{im} A \subset W$ are invariant. If $A \neq 0$, then $\ker A \neq V$, so $\ker A = 0$, and $\operatorname{im} A \neq 0$, so $\operatorname{im} A = W$. (ii) $A \in \mathrm{Hom}_G(V, V)$ has an eigenvalue $\lambda \in \mathbb{C}$, and $A - \lambda$ is a non-invertible intertwiner, hence 0 by (i). For the last claim, if $A_0 : V \to W$ is an invertible intertwiner, then $A \mapsto A_0^{-1}A$ identifies $\mathrm{Hom}_G(V, W)$ with $\mathrm{Hom}_G(V, V) = \mathbb{C}$. □

Corollary 1.3.2: Every irreducible representation of an abelian group is one-dimensional.

Proof. Every $\pi(h)$ is an intertwiner, hence a scalar, so every subspace is invariant, and irreducibility forces $\dim V = 1$. □

Proposition 1.3.3: The irreducible representations of $\mathbb{T}^n$ are exactly the characters

$$\chi_k(z_1, \dots, z_n) = z_1^{k_1} \cdots z_n^{k_n}, \qquad k \in \mathbb{Z}^n,$$

and they are pairwise inequivalent.

Proof. By Corollary 1.3.2, an irreducible representation is a continuous homomorphism $\chi : \mathbb{T}^n \to \mathbb{C}^\times$; its image is a compact subgroup of $\mathbb{C}^\times$, hence lies in $\mathbb{T}$ (if $|\chi(z)| \neq 1$, the powers $\chi(z)^{\pm m}$ would be unbounded). Since $\chi(z) = \prod_j \chi(1, \dots, z_j, \dots, 1)$, it suffices to treat $n = 1$. The map $\psi(t) := \chi(e^{2\pi i t})$ is a continuous homomorphism $\mathbb{R} \to \mathbb{T}$; let $\tilde{\psi} : \mathbb{R} \to \mathbb{R}$ be the continuous lift with $\psi(t) = e^{2\pi i \tilde{\psi}(t)}$, $\tilde{\psi}(0) = 0$. The function $(s, t) \mapsto \tilde{\psi}(s + t) - \tilde{\psi}(s) - \tilde{\psi}(t)$ is continuous, integer-valued on the connected set $\mathbb{R}^2$, and vanishes at $(0, 0)$; hence $\tilde{\psi}$ is additive. A continuous additive function is linear, $\tilde{\psi}(t) = kt$, and $1 = \psi(1) = e^{2\pi i k}$ gives $k \in \mathbb{Z}$. Characters with different k are different functions, hence inequivalent. □

2 Haar measure and complete reducibility

2.1 Existence and uniqueness of Haar measure

Throughout this subsection, G is a compact group, and $C(G)$ carries the sup norm $\| \cdot \|_\infty$. For $f \in C(G)$ and $h \in G$ let

$$(\lambda_h f)(g) := f(hg), \qquad (\rho_h f)(g) := f(gh).$$

Note $\lambda_k\lambda_h = \lambda_{hk}$ and $\rho_k\rho_h = \rho_{kh}$, so the translates of a translate of f are translates of f. For real-valued $f \in C(G)$ let K_f^λ (resp. K_f^ρ) be the closure in $C(G)$ of the convex hull of $\{\lambda_h f \mid h \in G\}$ (resp. of $\{\rho_h f \mid h \in G\}$). Since λ_h is a linear isometry of $C(G)$, $\lambda_h(K_f^\lambda) \subset K_f^\lambda$, and $K_\xi^\lambda \subset K_f^\lambda$ for every $\xi \in K_f^\lambda$; similarly for ρ.

Lemma 2.1.1: K_f^λ and K_f^ρ are compact, and every element of them takes values in $[\min f, \max f]$.

Proof. The set $\{\lambda_h f\}_{h\in G}$ is bounded by $\|f\|_\infty$ and equicontinuous: given $\varepsilon > 0$, choose U as in Lemma 1.1.5; for $g' = gu$ with $u \in U$, $|\lambda_h f(g') - \lambda_h f(g)| = |f(hgu) - f(hg)| < \varepsilon$ for all h. Convex combinations and uniform limits preserve the bound, the estimate (with $\le \varepsilon$) and the range $[\min f, \max f]$. By Arzelà–Ascoli, the closed set K_f^λ is compact. For K_f^ρ argue in the same way with $g' = ug$, using $|\rho_h f(ug) - \rho_h f(g)| = |f(ugh) - f(gh)| < \varepsilon$ (first inequality of Lemma 1.1.5). □

Lemma 2.1.2: K_f^λ and K_f^ρ contain constant functions.

Proof. The oscillation $\omega(\varphi) := \max\varphi - \min\varphi$ is continuous on $C(G)$, so it attains its minimum on the compact set K_f^λ at some φ_0. Suppose $M := \max\varphi_0 > m := \min\varphi_0$. The set $U := \{g \mid \varphi_0(g) < \frac{M+m}{2}\}$ is open and nonempty. The translates kU, $k \in G$, cover G (as $x \in xu^{-1}U$ for $u \in U$), so $G = k_1U \cup \cdots \cup k_nU$. Let $\psi := \frac{1}{n}\sum_i \lambda_{k_i^{-1}}\varphi_0 \in K_f^\lambda$. For every g there is i with $k_i^{-1}g \in U$, so

$$\psi(g) \le \frac{1}{n}\Big((n-1)M + \frac{M+m}{2}\Big) < M.$$

Since ψ is continuous on the compact space G, $\max\psi < M$, while $\min\psi \ge m$. Thus $\omega(\psi) < \omega(\varphi_0)$, a contradiction. Hence φ_0 is constant. The same proof works for K_f^ρ. □

Lemma 2.1.3: If $c \in K_f^\lambda$ and $c' \in K_f^\rho$ are constants, then $c = c'$. Consequently, K_f^λ and K_f^ρ contain exactly one constant, and it is the same; we denote it by $I(f)$.

Proof. Let $\varepsilon > 0$ and choose convex combinations with $|\sum_i a_i f(h_i g) - c| \le \varepsilon$ and $|\sum_j b_j f(gk_j) - c'| \le \varepsilon$ for all g. Then the number $S := \sum_{i,j} a_i b_j f(h_i k_j)$ satisfies $|S - c| \le \varepsilon$ (put $g = k_j$ in the first estimate and average with weights b_j) and $|S - c'| \le \varepsilon$ (put $g = h_i$ in the second and average with weights a_i). So $|c - c'| \le 2\varepsilon$ for all ε. □

Lemma 2.1.4: For real $f, f_1, f_2 \in C(G)$, $a \in \mathbb{R}$ and $h \in G$:

(i) $I(1) = 1$, $I(af) = aI(f)$ and $\min f \le I(f) \le \max f$;

(ii) $I(\lambda_h f) = I(\rho_h f) = I(f)$, and $I(f^\vee) = I(f)$ for $f^\vee(g) := f(g^{-1})$;

(iii) if $\xi \in K_f^\lambda$ and $\|\xi - c\|_\infty \le \delta$ for a constant c, then $|I(f) - c| \le \delta$;

(iv) $I(f_1 + f_2) = I(f_1) + I(f_2)$;

(v) if $f \ge 0$ and $f \ne 0$, then $I(f) > 0$.

Proof. (i) $K_{af}^\lambda = aK_f^\lambda$, and Lemma 2.1.1. (ii) The left translates of $\lambda_h f$ are exactly the left translates of f, so $K_{\lambda_h f}^\lambda = K_f^\lambda$ and hence $I(\lambda_h f) = I(f)$. Similarly $K_{\rho_h f}^\rho = K_f^\rho$; since $I(\rho_h f)$ is the constant in $K_{\rho_h f}^\rho$ and $I(f)$ the constant in K_f^ρ (Lemma 2.1.3), $I(\rho_h f) = I(f)$. For the inversion, $\lambda_h(f^\vee) = (\rho_{h^{-1}} f)^\vee$, so the isometry $\varphi \mapsto \varphi^\vee$ maps K_f^ρ onto $K_{f^\vee}^\lambda$ and fixes constants. (iii) $K_\xi^\lambda \subset K_f^\lambda$, so the unique constant of K_ξ^λ is $I(f)$. All translates of ξ, and hence all elements of K_ξ^λ, are within δ of c. (iv) Let $\varepsilon > 0$ and choose $\varphi = \sum_i a_i \lambda_{h_i} f_1$ with $\|\varphi - I(f_1)\|_\infty \le \varepsilon$.

Put $f_2' := \sum_i a_i \lambda_{h_i} f_2 \in K_{f_2}^\lambda$, and since $K_{f_2'}^\lambda \subset K_{f_2}^\lambda$, the unique constant of $K_{f_2'}^\lambda$ is $I(f_2)$, i.e., $I(f_2') = I(f_2)$. Choose $\psi = \sum_j b_j \lambda_{k_j} f_2'$ with $\|\psi - I(f_2)\|_\infty \le \varepsilon$. Then

$$\xi := \sum_j b_j \lambda_{k_j} \sum_i a_i \lambda_{h_i}(f_1 + f_2) = \sum_j b_j \lambda_{k_j} \varphi + \psi \in K_{f_1+f_2}^\lambda$$

satisfies $\|\xi - I(f_1) - I(f_2)\|_\infty \le 2\varepsilon$, so $|I(f_1 + f_2) - I(f_1) - I(f_2)| \le 2\varepsilon$ by (iii). (v) Let $c := \frac{1}{2} \max f > 0$ and $U := \{f > c\}$. As in the proof of Lemma 2.1.2, $G = k_1 U \cup \cdots \cup k_n U$, and then $\sum_i \lambda_{k_i^{-1}} f \ge c$ on G. By (i), (ii), (iv), $nI(f) = I(\sum_i \lambda_{k_i^{-1}} f) \ge c$. □

We extend I to complex-valued $f = u + iv$ by $I(f) := I(u) + iI(v)$. By Lemma 2.1.4, I is a positive normalized linear functional on $C(G)$ which is invariant under left and right translations and under inversion.

Theorem 2.1.1: Haar measure

Let G be a compact group. There is a unique regular Borel probability measure μ on G which is left invariant, i.e., $\mu(hA) = \mu(A)$ for all Borel sets A and $h \in G$. It is also right invariant and invariant under $g \mapsto g^{-1}$, and $\mu(U) > 0$ for every nonempty open $U \subset G$.

Proof. By Riesz–Markov there is a unique regular Borel measure μ with $\int f\,d\mu = I(f)$ for $f \in C(G)$, and $\mu(G) = I(1) = 1$. For $h \in G$, $\mu_h(A) := \mu(hA)$ is again a regular Borel measure with $\int f\,d\mu_h = \int f(h^{-1}g)\,d\mu(g) = I(\lambda_{h^{-1}} f) = I(f)$, so $\mu_h = \mu$ by the uniqueness part of Riesz–Markov. Right invariance and inversion invariance follow in the same way. If $U \ne \emptyset$ is open, Urysohn gives $f \in C(G)$, $0 \le f \le 1_U$, $f \ne 0$, so $\mu(U) \ge I(f) > 0$.

Uniqueness: let ν be a left-invariant regular Borel probability measure and $J(f) := \int f\,d\nu$; then $J(\lambda_h f) = J(f)$. For real f and $\varepsilon > 0$ choose a convex combination $\varphi = \sum_i a_i \lambda_{h_i} f$ with $\|\varphi - I(f)\|_\infty \le \varepsilon$. Then $J(\varphi) = J(f)$ and $|J(\varphi) - I(f)| \le \varepsilon$ since J is positive with $J(1) = 1$. Hence $J = I$ and $\nu = \mu$ by Riesz–Markov. □

We write $\int_G f(g)\,dg := \int f\,d\mu$. Thus for $f \in L^1(G)$ and $h \in G$,

$$\int_G f(hg)\,dg = \int_G f(gh)\,dg = \int_G f(g^{-1})\,dg = \int_G f(g)\,dg.$$

Example 2.1.5: (i) For finite G, $\int_G f = \frac{1}{|G|} \sum_{g \in G} f(g)$, since this measure is left invariant.

(ii) For $\mathbb{T}^n$, $\int f = (2\pi)^{-n} \int_{[0,2\pi]^n} f(e^{i\theta_1}, \ldots, e^{i\theta_n})\,d\theta$, by translation invariance of Lebesgue measure and periodicity.

(iii) Writing $g = \left(\begin{smallmatrix} a & -\bar{b} \\ b & \bar{a} \end{smallmatrix}\right)$ with $|a|^2 + |b|^2 = 1$ identifies SU(2) with $S^3 \subset \mathbb{C}^2 \cong \mathbb{R}^4$. (Indeed, if $g = \left(\begin{smallmatrix} a & c \\ b & d \end{smallmatrix}\right) \in \mathrm{SU}(2)$, comparing $g^{-1} = g^*$ with $g^{-1} = \left(\begin{smallmatrix} d & -c \\ -b & a \end{smallmatrix}\right)$ gives $d = \bar{a}$, $c = -\bar{b}$; conversely such matrices are unitary of determinant 1.) Left multiplication $x \mapsto gx$ by $g \in \mathrm{SU}(2)$ is a linear isometry of $\mathbb{C}^2 = \mathbb{R}^4$ preserving S^3, so the normalized surface measure of S^3, which is O(4)-invariant, is left invariant, hence it is the Haar measure of SU(2).

2.2 Unitarity and complete reducibility

Theorem 2.2.1: Weyl's unitarian trick

Let (π, V) be a representation of a compact group G. Then V has an inner product for which all $\pi(g)$ are unitary.

Proof. Let $(\cdot,\cdot)$ be any inner product on V and $\langle v,w\rangle := \int_G (\pi(g)v,\pi(g)w)\,dg$. This is a Hermitian form. For $v\neq 0$ the integrand $g\mapsto(\pi(g)v,\pi(g)v)$ is continuous and strictly positive, hence bounded below by a positive constant on the compact G, so $\langle v,v\rangle>0$. By right invariance, $\langle\pi(h)v,\pi(h)w\rangle = \int_G(\pi(gh)v,\pi(gh)w)\,dg = \langle v,w\rangle$. □

Corollary 2.2.1 (Complete reducibility): Every representation of a compact group is a direct sum of irreducible representations.

Proof. By Theorem 2.2.1 we may assume π unitary. If $V'\subset V$ is invariant, so is $V'^{\perp}$: for $w\in V'^{\perp}$, $v\in V'$, $\langle\pi(g)w,v\rangle = \langle w,\pi(g)^{-1}v\rangle = 0$. Now induct on $\dim V$. □

Remark 2.2.2: Compactness is essential: the representation $t\mapsto \left(\begin{smallmatrix}1 & t\\ 0 & 1\end{smallmatrix}\right)$ of $(\mathbb{R},+)$ on $\mathbb{C}^2$ has the invariant line $\mathbb{C}e_1$, and no other invariant line (an eigenvector of $\left(\begin{smallmatrix}1 & 1\\ 0 & 1\end{smallmatrix}\right)$ lies in $\mathbb{C}e_1$), so it has no invariant complement.

Proposition 2.2.3 (Multiplicities): Let $V = V_1\oplus\cdots\oplus V_m$ be a decomposition of a representation of a compact group into irreducibles, and (σ,W) irreducible. Then the number of V_j equivalent to σ equals $\dim\mathrm{Hom}_G(W,V)$. In particular this number, the *multiplicity* $m_\sigma(V)$, does not depend on the decomposition, and two representations with the same multiplicities are equivalent.

Proof. $\mathrm{Hom}_G(W,V) = \bigoplus_j \mathrm{Hom}_G(W,V_j)$, and $\dim\mathrm{Hom}_G(W,V_j)\in\{0,1\}$ according to whether $V_j\cong W$ (Lemma 1.3.1). The last statement follows since both representations are then equivalent to $\bigoplus_\sigma m_\sigma\,\sigma$. □

3 Schur orthogonality and characters

3.1 Matrix coefficients and the orthogonality relations

Definition 3.1.1: Let (π,V) be a unitary representation. For $u,v\in V$, the function $f^\pi_{u,v}(g) := \langle\pi(g)u,v\rangle$ is a *matrix coefficient* of π. If $e_1,\dots,e_d$ is an orthonormal basis, $\pi_{ij}(g) := \langle\pi(g)e_j,e_i\rangle$ are the entries of the matrix of $\pi(g)$.

On $L^2(G) := L^2(G,dg)$ we use $\langle f_1,f_2\rangle_{L^2} := \int_G f_1(g)\overline{f_2(g)}\,dg$.

Theorem 3.1.1: Schur orthogonality relations

Let (π,V), (σ,W) be irreducible unitary representations of a compact group G.

(i) If $\pi\not\cong\sigma$, then $\langle f^\pi_{u,v}, f^\sigma_{u',v'}\rangle_{L^2} = 0$ for all $u,v\in V$, $u',v'\in W$.

(ii) For $u_1,v_1,u_2,v_2\in V$: $\displaystyle\int_G \langle\pi(g)u_1,v_1\rangle\overline{\langle\pi(g)u_2,v_2\rangle}\,dg = \frac{1}{d_\pi}\langle u_1,u_2\rangle\overline{\langle v_1,v_2\rangle}$.

In matrix form: $\displaystyle\int_G \pi_{ij}(g)\overline{\sigma_{kl}(g)}\,dg = \frac{\delta_{\pi\sigma}\delta_{ik}\delta_{jl}}{d_\pi}$, where $\delta_{\pi\sigma}=1$ if $\pi=\sigma$ and 0 if $\pi\not\cong\sigma$.

Proof. For a linear map $A: W\to V$ let $\tilde A := \int_G \pi(g)A\sigma(g)^{-1}\,dg$ (entrywise integral). By left invariance, $\pi(h)\tilde A\sigma(h)^{-1} = \int_G\pi(hg)A\sigma(hg)^{-1}\,dg = \tilde A$, so $\tilde A\in\mathrm{Hom}_G(W,V)$. By Schur's lemma, $\tilde A = 0$ if $\pi\not\cong\sigma$, while for $\sigma=\pi$, $\tilde A = c\cdot 1$ with $cd_\pi = \mathrm{tr}\,\tilde A = \mathrm{tr}\,A$.

Take $Ax := \langle x,u'\rangle u$ with $u\in V$, $u'\in W$. For $w\in V$, $w'\in W$, using unitarity of σ,

$$\langle\tilde A w',w\rangle = \int_G\langle\sigma(g)^{-1}w',u'\rangle\langle\pi(g)u,w\rangle\,dg = \int_G\langle\pi(g)u,w\rangle\overline{\langle\sigma(g)u',w'\rangle}\,dg.$$

If $\pi \not\cong \sigma$ this vanishes, which is (i). If $\sigma = \pi$, then $\operatorname{tr} A = \langle u, u' \rangle$ and $\langle \tilde{A} w', w \rangle = \frac{1}{d_\pi} \langle u, u' \rangle \langle w', w \rangle$; renaming $u = u_1$, $w = v_1$, $u' = u_2$, $w' = v_2$ gives (ii). The matrix form follows with $u_1 = e_j$, $v_1 = e_i$, $u_2 = e_l$, $v_2 = e_k$. □

3.2 Characters

Definition 3.2.1: The *character* of a representation (π, V) is $\chi_\pi(g) := \operatorname{tr} \pi(g)$. A function f on G is a *class function* if $f(hgh^{-1}) = f(g)$ for all g, h.

Lemma 3.2.2: For representations π, σ of a compact group:

(i) $\chi_\pi(e) = d_\pi$, and χ_π is a continuous class function;

(ii) $\pi \cong \sigma$ implies $\chi_\pi = \chi_\sigma$;

(iii) $\chi_{\pi \oplus \sigma} = \chi_\pi + \chi_\sigma$ and $\chi_{\pi \otimes \sigma} = \chi_\pi \chi_\sigma$;

(iv) $\chi_{\pi^*}(g) = \chi_\pi(g^{-1}) = \overline{\chi_\pi(g)}$.

Proof. (i)–(iii) follow from $\operatorname{tr}(ABA^{-1}) = \operatorname{tr} B$, $\operatorname{tr}(A \oplus B) = \operatorname{tr} A + \operatorname{tr} B$ and $\operatorname{tr}(A \otimes B) = \operatorname{tr} A \operatorname{tr} B$. (iv): the matrix of $\pi^*(g)$ in the dual basis is the transpose of that of $\pi(g^{-1})$; with an invariant inner product (Theorem 2.2.1), $\pi(g^{-1}) = \pi(g)^*$ has trace $\overline{\operatorname{tr} \pi(g)}$. □

Theorem 3.2.1: Orthogonality of characters

For irreducible representations π, σ of a compact group, $\langle \chi_\pi, \chi_\sigma \rangle_{L^2} = 1$ if $\pi \cong \sigma$ and $= 0$ otherwise.

Proof. By Lemma 3.2.2(ii) and Theorem 2.2.1 we may assume both unitary and $\sigma = \pi$ in the equivalent case. Then $\chi_\pi = \sum_i \pi_{ii}$, and Theorem 3.1.1 gives $\langle \chi_\pi, \chi_\sigma \rangle = \sum_{i,k} \delta_{\pi\sigma} \delta_{ik} / d_\pi = \delta_{\pi\sigma}$. □

Corollary 3.2.3: Let π be a representation of a compact group G.

(i) $m_\sigma(\pi) = \langle \chi_\pi, \chi_\sigma \rangle_{L^2}$ for every irreducible σ.

(ii) π is determined up to equivalence by χ_π.

(iii) $\|\chi_\pi\|_{L^2}^2 = \sum_\sigma m_\sigma(\pi)^2$; in particular π is irreducible if and only if $\|\chi_\pi\|_{L^2} = 1$.

(iv) $\dim V^G = \int_G \chi_\pi(g)\, dg$, where V^G is the space of fixed vectors.

Proof. Write $\pi \cong \bigoplus_\sigma m_\sigma(\pi) \sigma$ (Corollary 2.2.1), so $\chi_\pi = \sum_\sigma m_\sigma(\pi) \chi_\sigma$, and apply Theorem 3.2.1; (ii) then follows from Proposition 2.2.3. For (iv), V^G is the isotypic component of the trivial representation, whose character is 1, so $\dim V^G = m_1(\pi) = \langle \chi_\pi, 1 \rangle$. □

3.3 First example: the representations of SU(2)

For $\theta \in \mathbb{R}$ put $t_\theta := \operatorname{diag}(e^{i\theta}, e^{-i\theta}) \in \mathrm{SU}(2)$.

Lemma 3.3.1: Every $g \in \mathrm{SU}(2)$ is conjugate in $\mathrm{SU}(2)$ to t_θ for a unique $\theta \in [0, \pi]$, determined by $\operatorname{tr} g = 2 \cos\theta$.

Proof. By the spectral theorem there is $k \in \mathrm{U}(2)$ with $k^{-1} g k$ diagonal; its entries λ, λ^{-1} have modulus 1. Replacing k by $k \cdot (\det k)^{-1/2}$ (any square root) gives $k \in \mathrm{SU}(2)$. So g is conjugate to some t_θ, and $t_{-\theta} = w t_\theta w^{-1}$ with $w = \left(\begin{smallmatrix} 0 & 1 \\ -1 & 0 \end{smallmatrix}\right) \in \mathrm{SU}(2)$. Uniqueness: $\operatorname{tr} t_\theta = 2\cos\theta$ is injective on $[0, \pi]$. □

Lemma 3.3.2 (Integration formula for SU(2)): For every continuous class function f on SU(2),

$$\int_{\mathrm{SU}(2)} f(g)\,dg = \frac{2}{\pi}\int_0^\pi f(t_\theta)\sin^2\theta\,d\theta.$$

Proof. By Example 2.1.5(iii), Haar measure is the normalized surface measure of S^3. In the coordinates $x = (x_0,\dots,x_3) = (\operatorname{Re}a, \operatorname{Im}a, \operatorname{Re}b, \operatorname{Im}b)$, $\operatorname{tr} g = 2x_0$, so by Lemma 3.3.1, $f(g) = F(x_0)$ with $F(\cos\theta) := f(t_\theta)$. The map $\Phi(\theta,\omega) := (\cos\theta, \sin\theta\,\omega)$, $(\theta,\omega) \in (0,\pi)\times S^2$, is a diffeomorphism onto S^3 minus two points. Its derivative maps ∂_θ to a unit vector orthogonal to $\Phi_*(T_\omega S^2)$, and scales $T_\omega S^2$ by $\sin\theta$; hence the surface measure is $\sin^2\theta\,d\theta\,d\sigma_{S^2}(\omega)$. Thus $\operatorname{vol}(S^3) = 4\pi\int_0^\pi \sin^2\theta\,d\theta = 2\pi^2$ and

$$\int_{\mathrm{SU}(2)} f = \frac{4\pi}{2\pi^2}\int_0^\pi F(\cos\theta)\sin^2\theta\,d\theta = \frac{2}{\pi}\int_0^\pi f(t_\theta)\sin^2\theta\,d\theta. \qquad \square$$

Theorem 3.3.1: Irreducible representations of SU(2)

The representations π_n, $n \in \mathbb{N}_0$, of Example 1.2.4 are irreducible and pairwise inequivalent, with

$$\chi_n(t_\theta) = \sum_{k=0}^n e^{i(n-2k)\theta} = \frac{\sin((n+1)\theta)}{\sin\theta} \quad (\theta \notin \pi\mathbb{Z}).$$

Every irreducible representation of SU(2) is equivalent to some π_n.

Proof. Since $t_\theta^{-1} z = (e^{-i\theta}z_1, e^{i\theta}z_2)$, the monomial $z_1^k z_2^{n-k}$ is an eigenvector of $\pi_n(t_\theta)$ with eigenvalue $e^{i(n-2k)\theta}$; summing the geometric series gives χ_n. By Lemma 3.3.2,

$$\langle \chi_n, \chi_m\rangle_{L^2} = \frac{2}{\pi}\int_0^\pi \sin((n+1)\theta)\sin((m+1)\theta)\,d\theta = \delta_{nm},$$

so by Corollary 3.2.3 the π_n are irreducible and pairwise inequivalent. Let σ be irreducible and not equivalent to any π_n. Then for all n, using that χ_n is real,

$$0 = \langle \chi_\sigma, \chi_n\rangle_{L^2} = \frac{2}{\pi}\int_0^\pi \big(\chi_\sigma(t_\theta)\sin\theta\big)\sin((n+1)\theta)\,d\theta.$$

By completeness of $\{\sin(m\theta)\}_{m\ge1}$ in $L^2([0,\pi])$, the continuous function $\chi_\sigma(t_\theta)\sin\theta$ vanishes on $[0,\pi]$, so $\chi_\sigma(t_\theta) = 0$ for $\theta \in (0,\pi)$ and by continuity for $\theta = 0$. This contradicts $\chi_\sigma(e) = d_\sigma \ge 1$. $\square$

Corollary 3.3.3 (Clebsch–Gordan): For $m \ge n \ge 0$: $\pi_m \otimes \pi_n \cong \bigoplus_{k=0}^n \pi_{m+n-2k}$.

Proof. By Corollary 3.2.3(ii) and Lemma 3.2.2(iii), it suffices to compare characters on the conjugacy class representatives t_θ, $\theta \notin \pi\mathbb{Z}$ (and then everywhere by continuity). With $x = e^{i\theta}$,

$$\chi_m\chi_n = \sum_{j=0}^n \chi_m(t_\theta)x^{n-2j} = \sum_{j=0}^n \frac{x^{m+n-2j+1} - x^{-(m-n+2j+1)}}{x - x^{-1}}.$$

Substituting $j \mapsto n-j$ in the second part, this becomes $\sum_{k=0}^n \frac{x^{m+n-2k+1} - x^{-(m+n-2k+1)}}{x-x^{-1}} = \sum_{k=0}^n \chi_{m+n-2k}$. $\square$

Remark 3.3.4: In physics one writes $j = n/2$ (*spin*) and $D^{(j)} := \pi_{2j}$; then $D^{(j_1)} \otimes D^{(j_2)} \cong \bigoplus_{J=|j_1-j_2|}^{j_1+j_2} D^{(J)}$.

4 The Peter–Weyl theorem

4.1 Compact operators and the regular representation

Theorem 4.1.1: Spectral theorem for compact self-adjoint operators

Let K be a compact self-adjoint operator on a Hilbert space $\mathcal{H}$. Then $\mathcal{H} = \ker K \oplus \widehat{\bigoplus}_\lambda \mathcal{H}_\lambda$ (orthogonal Hilbert sum), where λ runs through the nonzero eigenvalues of K, which are real, and each $\mathcal{H}_\lambda := \ker(K - \lambda)$ is finite-dimensional. If $K \neq 0$, there is at least one nonzero eigenvalue.

This is a prerequisite from functional analysis.

Definition 4.1.1: The *left* and *right regular representations* of a compact group G on $L^2(G)$ are $(L(h)f)(g) := f(h^{-1}g)$ and $(R(h)f)(g) := f(gh)$.

Lemma 4.1.2: L and R are unitary representations, and they commute.

Proof. Unitarity and the homomorphism property follow from invariance of Haar measure, and $L(h)R(k)f(g) = f(h^{-1}gk) = R(k)L(h)f(g)$. For $f \in C(G)$, Lemma 1.1.5 gives $\|L(h)f - f\|_2 \leq \|L(h)f - f\|_\infty \to 0$ as $h \to e$. For $f \in L^2(G)$ and $\varepsilon > 0$ choose $f_0 \in C(G)$ with $\|f - f_0\|_2 < \varepsilon$; then $\|L(h)f - f\|_2 \leq 2\varepsilon + \|L(h)f_0 - f_0\|_2$. Continuity at h_0 follows from $\|L(h)f - L(h_0)f\|_2 = \|L(h_0^{-1}h)f - f\|_2$. The same applies to R. □

Definition 4.1.3: For $\varphi \in C(G)$ and $f \in L^2(G)$ let $(\varphi * f)(g) := \int_G \varphi(h) f(h^{-1}g)\, dh = \int_G \varphi(gh^{-1}) f(h)\, dh$ (the second form follows by substituting $h \mapsto gh^{-1}$, using invariance under right translation and inversion).

Lemma 4.1.4: Let $\varphi \in C(G)$ and $T_\varphi f := \varphi * f$.

(i) T_φ is a Hilbert–Schmidt operator with kernel $K(g,h) = \varphi(gh^{-1})$, hence compact. If $\varphi(g^{-1}) = \overline{\varphi(g)}$, then T_φ is self-adjoint.

(ii) $T_\varphi R(k) = R(k) T_\varphi$ for all $k \in G$.

(iii) $\varphi * f$ is continuous for $f \in L^2(G)$, and $\|\varphi * f\|_\infty \leq \|\varphi\|_2 \|f\|_2$.

(iv) $\varphi * f = \int_G \varphi(h) L(h) f\, dh$ in the sense that $\langle \varphi * f, u \rangle = \int_G \varphi(h) \langle L(h)f, u \rangle\, dh$ for $u \in L^2(G)$.

Proof. (i) The kernel is continuous on $G \times G$, hence square integrable; $\overline{K(h,g)} = \overline{\varphi(hg^{-1})} = \varphi(gh^{-1}) = K(g,h)$. (ii) $(\varphi * R(k)f)(g) = \int \varphi(h) f(h^{-1}gk)\, dh = (\varphi * f)(gk)$. (iii) By Cauchy–Schwarz and invariance, $|(\varphi * f)(g) - (\varphi * f)(g')| \leq \big(\int_G |\varphi(gh^{-1}) - \varphi(g'h^{-1})|^2 dh\big)^{1/2} \|f\|_2$, which tends to 0 as $g' \to g$ by Lemma 1.1.5; and $|(\varphi * f)(g)| \leq \|\varphi\|_2 \|f\|_2$. (iv) Fubini. □

Lemma 4.1.5 (Approximate identities): For every neighborhood U of e there is $\varphi_U \in C(G)$ with $\varphi_U \geq 0$, $\operatorname{supp} \varphi_U \subset U$, $\int_G \varphi_U = 1$ and $\varphi_U(g^{-1}) = \varphi_U(g)$. For $f \in L^2(G)$, $\|\varphi_U * f - f\|_2 \leq \sup_{h \in U} \|L(h)f - f\|_2$, and for $f \in C(G)$, $\|\varphi_U * f - f\|_\infty \leq \sup_{h \in U,\, g \in G} |f(h^{-1}g) - f(g)|$. In particular $\varphi_U * f \to f$ in L^2 (resp. uniformly for continuous f) as U shrinks.

Proof. By Urysohn choose $\psi \in C(G)$, $0 \leq \psi \leq 1$, $\psi(e) = 1$, $\operatorname{supp} \psi \subset U \cap U^{-1}$. Then $\psi + \psi^\vee \geq 0$ is nonzero, so its integral is positive (Theorem 2.1.1), and $\varphi_U := (\psi + \psi^\vee)/\int (\psi + \psi^\vee)$ works. Since $\int \varphi_U = 1$, $\varphi_U * f - f = \int \varphi_U(h)(L(h)f - f)\, dh$ (weakly, Lemma 4.1.4(iv)), so $|\langle \varphi_U * f - f, u \rangle| \leq \sup_{h \in U} \|L(h)f - f\|_2 \|u\|_2$. The uniform estimate is immediate, and convergence follows from Lemmas 4.1.2 and 1.1.5. □

4.2 The Peter–Weyl theorem

Let $\mathcal{E}(G) \subset C(G)$ be the span of all matrix coefficients of (finite-dimensional) representations. By Theorem 2.2.1 and Corollary 2.2.1, $\mathcal{E}(G)$ is spanned by the functions π_{ij} for unitary irreducible π and orthonormal bases. For a representation π, the space $\mathcal{E}_\pi := \operatorname{span}\{f^\pi_{u,v}\}$ is finite-dimensional and invariant under L and R, since $L(h)f^\pi_{u,v} = f^\pi_{u,\pi(h)v}$ and $R(h)f^\pi_{u,v} = f^\pi_{\pi(h)u,v}$.

Theorem 4.2.1: Peter–Weyl

Let G be a compact group, and fix a unitary representative π of each class in $\widehat{G}$ together with an orthonormal basis.

(i) $\mathcal{E}(G)$ is dense in $C(G)$ with respect to $\|\cdot\|_\infty$.

(ii) $\{\sqrt{d_\pi}\,\pi_{ij} \mid \pi \in \widehat{G},\ 1 \le i,j \le d_\pi\}$ is an orthonormal basis of $L^2(G)$.

(iii) The maps $A \mapsto f_A$, $f_A(x) := \sqrt{d_\pi}\operatorname{tr}(A\pi(x)^{-1})$, define a unitary equivalence of $G \times G$-representations

$$\widehat{\bigoplus_{\pi\in\widehat{G}}} \operatorname{End}(V_\pi) \xrightarrow{\cong} L^2(G),$$

where $(g,h)\cdot A := \pi(g)A\pi(h)^{-1}$ and $(g,h)\cdot f := L(g)R(h)f$, and $\operatorname{End}(V_\pi)$ carries the Hilbert–Schmidt inner product $\langle A, B\rangle = \operatorname{tr}(AB^*)$.

(iv) $\{\chi_\pi\}_{\pi\in\widehat{G}}$ is an orthonormal basis of the closed subspace $L^2(G)^{\text{class}}$ of class functions.

Proof. Step 1: $\mathcal{E}(G)$ is dense in $L^2(G)$. Suppose $0 \ne f \perp \mathcal{E}(G)$. By Lemma 4.1.5 there is a symmetric approximate identity φ with $\varphi * f \ne 0$. By Lemma 4.1.4, T_φ is compact, self-adjoint and commutes with R. Let $\lambda \ne 0$ be an eigenvalue and $\mathcal{H}_\lambda$ its finite-dimensional eigenspace. It is R-invariant, and its elements are continuous since $\psi = \lambda^{-1}\varphi * \psi$. Choose a basis $\psi_1, \dots, \psi_m$ of $\mathcal{H}_\lambda$ and write $R(h)\psi_j = \sum_i \rho_{ij}(h)\psi_i$; then $\rho = R|_{\mathcal{H}_\lambda}$ is a representation of G (continuity follows from Lemma 4.1.2). Both sides are continuous functions, so evaluating at e gives

$$\psi_j(h) = \sum_i \rho_{ij}(h)\psi_i(e),$$

so $\mathcal{H}_\lambda \subset \mathcal{E}(G)$ and $f \perp \mathcal{H}_\lambda$. This holds for every $\lambda \ne 0$, so $f \in \ker T_\varphi$ by Theorem 4.1.1, contradicting $\varphi * f \ne 0$.

Step 2: (i). Let $f \in C(G)$, $\varepsilon > 0$. Choose φ with $\|\varphi * f - f\|_\infty < \varepsilon$ (Lemma 4.1.5) and, by Step 1, $e' \in \mathcal{E}(G)$ with $\|f - e'\|_2 < \varepsilon/(1+\|\varphi\|_2)$. If $e' \in \mathcal{E}_{\pi_1} + \cdots + \mathcal{E}_{\pi_r}$, a finite-dimensional L-invariant space E, then $\varphi * e' = \int \varphi(h)L(h)e'\,dh \in E$ (a weak integral of a continuous E-valued function lies in the closed subspace E). By Lemma 4.1.4(iii), $\|f - \varphi*e'\|_\infty \le \|f - \varphi*f\|_\infty + \|\varphi*(f-e')\|_\infty < 2\varepsilon$.

Step 3: (ii) and (iii). Orthonormality is Theorem 3.1.1, completeness is Step 1. For (iii), $L(g)R(h)f_A(x) = \sqrt{d_\pi}\operatorname{tr}(A\pi(h)^{-1}\pi(x)^{-1}\pi(g)) = f_{\pi(g)A\pi(h)^{-1}}(x)$. Since $\operatorname{tr}(A\pi(x)^*) = \sum_{ij} A_{ij}\overline{\pi_{ij}(x)}$, the map sends the orthonormal basis $\{E_{ij}\}$ of $\operatorname{End}(V_\pi)$ to $\{\sqrt{d_\pi}\,\overline{\pi_{ij}}\}$. Now $\overline{\pi_{ij}}$ are the matrix coefficients of π^* in the dual basis (Lemma 3.2.2(iv) and its proof), and $\pi \mapsto \pi^*$ permutes $\widehat{G}$ (since $\pi^{**} \cong \pi$ and π^* is irreducible as $\chi_{\pi^*} = \overline{\chi_\pi}$ has norm 1). So (iii) follows from (ii).

Step 4: (iv). The characters are orthonormal class functions. Let P_π be the orthogonal projection onto $\mathcal{E}_\pi$. As $\mathcal{E}_\pi$ and $\mathcal{E}_\pi^\perp$ are invariant under the unitary operators $C(h) := L(h)R(h)$, P_π commutes with $C(h)$; class functions are the fixed vectors of all $C(h)$, so P_π maps class functions to class functions. A class function in $\mathcal{E}_\pi$ has the form $x \mapsto \operatorname{tr}(A\pi(x))$ with $\operatorname{tr}(A\pi(hxh^{-1})) = \operatorname{tr}(A\pi(x))$; averaging over h gives $\operatorname{tr}(A\pi(x)) = \operatorname{tr}(\tilde{A}\pi(x))$ with $\tilde{A} = \int_G \pi(h)^{-1}A\pi(h)\,dh$, an in-

tertwiner, hence $\tilde{A} = \frac{\operatorname{tr} A}{d_\pi} 1$ by Schur. So $P_\pi f \in \mathbb{C}\chi_\pi$, and by (ii), $f = \sum_\pi P_\pi f = \sum_\pi \langle f, \chi_\pi\rangle \chi_\pi$ for every class function $f \in L^2(G)$. □

Example 4.2.1 (Fourier series): For $G = \mathbb{T}$, $\widehat{G} = \{\chi_k\}_{k\in\mathbb{Z}}$ (Proposition 1.3.3), and Theorem 4.2.1 states completeness of $\{e^{ik\theta}\}$ in $L^2(\mathbb{T})$ and uniform density of trigonometric polynomials in $C(\mathbb{T})$. In general, $f = \sum_\pi d_\pi \operatorname{tr}(\hat{f}(\pi)\pi(\cdot))$ in $L^2(G)$, with $\hat{f}(\pi) := \int_G f(g)\pi(g)^{-1}\,dg$, and $\|f\|_2^2 = \sum_\pi d_\pi \|\hat{f}(\pi)\|_{\mathrm{HS}}^2$ (Plancherel). Indeed, $\langle f, \pi_{ij}\rangle = \hat{f}(\pi)_{ji}$, so the component of f in $\mathcal{E}_\pi$ is $\sum_{ij} d_\pi \hat{f}(\pi)_{ji}\pi_{ij} = d_\pi \operatorname{tr}(\hat{f}(\pi)\pi(\cdot))$, of norm squared $d_\pi^2 \sum_{ij} |\hat{f}(\pi)_{ji}|^2/d_\pi$.

4.3 Consequences

Corollary 4.3.1: Let G be a compact group.

(i) If G is finite, then $\sum_{\pi\in\widehat{G}} d_\pi^2 = |G|$, and $|\widehat{G}|$ equals the number of conjugacy classes.

(ii) For $g \neq e$ there is $\pi \in \widehat{G}$ with $\pi(g) \neq 1$.

Proof. (i) Compare the dimension $|G|$ of $L^2(G)$, resp. the dimension of the space of class functions (the number of conjugacy classes), with Theorem 4.2.1(ii) and (iv). (ii) By Urysohn there is $f \in C(G)$ with $f(e) = 0$, $f(g) = 1$; by Theorem 4.2.1(i), some $e' \in \mathcal{E}(G)$ satisfies $e'(g) \neq e'(e)$. If all $\pi(g) = 1$, all matrix coefficients would take the same value at g and e. □

Theorem 4.3.1: Unitary representations of compact groups

Let $(\pi, \mathcal{H})$ be a unitary representation of a compact group G on a Hilbert space. Then $\mathcal{H}$ is an orthogonal Hilbert sum of finite-dimensional irreducible invariant subspaces. In particular every irreducible unitary representation is finite-dimensional.

Proof. Claim: every nonzero closed invariant subspace $\mathcal{H}'$ contains a finite-dimensional irreducible invariant subspace. Replacing $\mathcal{H}$ by $\mathcal{H}'$, we prove this for $\mathcal{H}$. Let $v \neq 0$ and let $R_g := \langle \cdot, \pi(g)v\rangle \pi(g)v$ (rank one). The sesquilinear form $(x, y) \mapsto \int_G \langle R_g x, y\rangle\, dg$ is bounded by $\|v\|^2\|x\|\|y\|$, so it defines a bounded operator K with $\langle Kx, y\rangle = \int_G \langle x, \pi(g)v\rangle\langle \pi(g)v, y\rangle\, dg$. It is self-adjoint and positive, and by left invariance of Haar measure $\pi(h)K\pi(h)^{-1} = K$. Moreover $\langle Kv, v\rangle = \int_G |\langle v, \pi(g)v\rangle|^2\, dg > 0$, since the integrand is continuous, nonnegative and positive at e (Theorem 2.1.1). K is compact: $\|R_g - R_{g'}\| \leq 2\|v\|\|\pi(g)v - \pi(g')v\|$, so $g \mapsto R_g$ is norm-continuous. Given $\varepsilon > 0$, cover G by finitely many open sets $U_i \ni g_i$ with $\|R_g - R_{g_i}\| < \varepsilon$ on U_i, refine to a Borel partition $B_i \subset U_i$, and let $K_\varepsilon := \sum_i \mu(B_i) R_{g_i}$, a finite-rank operator with $|\langle (K - K_\varepsilon)x, y\rangle| \leq \varepsilon\|x\|\|y\|$. So K is a norm limit of finite-rank operators. By Theorem 4.1.1, K has a nonzero eigenvalue with finite-dimensional, nonzero, π-invariant eigenspace, and an invariant subspace of minimal positive dimension in it is irreducible.

Decomposition. By Zorn's lemma choose a maximal family $\{\mathcal{H}_\alpha\}$ of mutually orthogonal finite-dimensional irreducible invariant subspaces. The orthogonal complement of $\widehat{\bigoplus}_\alpha \mathcal{H}_\alpha$ is closed and invariant (π is unitary); if it were nonzero, the claim would contradict maximality. Finally, if π is irreducible, the claim shows that $\mathcal{H}$ itself is finite-dimensional. □

Proposition 4.3.2 (Homogeneous spaces and Frobenius reciprocity): Let $K \subset G$ be a closed subgroup with Haar measure dk, and define $L^2(G/K) := \{f \in L^2(G) \mid R(k)f = f \text{ for all } k \in K\}$, a closed L-invariant subspace. Then, as representations of G (acting by L),

$$L^2(G/K) \cong \widehat{\bigoplus_{\pi\in\widehat{G}}} \left(\dim V_\pi^K\right) \cdot \pi,$$

i.e., π occurs in $L^2(G/K)$ with multiplicity $\dim V_\pi^K$, where V_π^K is the space of K-fixed vectors.

Proof. Under the isomorphism of Theorem 4.2.1(iii), $R(k)$ acts on $\mathrm{End}(V_\pi)$ by $A \mapsto A\pi(k)^{-1}$, componentwise, so $L^2(G/K)$ corresponds to $\widehat{\bigoplus}_\pi \{A \mid A\pi(k) = A\ \forall k \in K\}$. Let $P := \int_K \pi(k)\,dk$. By invariance of dk, $\pi(k)P = P\pi(k) = P$, $P^2 = P$ and $P^* = P$; so P is the orthogonal projection onto V_π^K (its image is fixed by K, and $Pv = v$ for $v \in V_\pi^K$). If $A\pi(k) = A$ for all k, then $A = AP$; conversely $AP\pi(k) = AP$. Hence the K-invariant part of $\mathrm{End}(V_\pi)$ is $\{A \mid A = AP\} \cong \mathrm{Hom}(V_\pi^K, V_\pi)$, on which $L(g)$ acts by $A \mapsto \pi(g)A$; choosing a basis of V_π^K identifies this with $\dim V_\pi^K$ copies of π. □

5 Lie groups and Lie algebras

We mostly work with *matrix Lie groups*, i.e., closed subgroups of $\mathrm{GL}(n,\mathbb{C})$. By Theorem 5.5.1, every compact Lie group is of this form. On $M_n(\mathbb{C})$ we use the operator norm.

5.1 The matrix exponential

Lemma 5.1.1: For $X, Y \in M_n(\mathbb{C})$ and $g \in \mathrm{GL}(n,\mathbb{C})$:

(i) $\exp X := e^X := \sum_{k\geq 0} X^k/k!$ converges absolutely and locally uniformly, and exp is smooth (indeed real analytic).

(ii) If $XY = YX$, then $e^{X+Y} = e^X e^Y$. In particular $(e^X)^{-1} = e^{-X}$ and $t \mapsto e^{tX}$ is a homomorphism $\mathbb{R} \to \mathrm{GL}(n,\mathbb{C})$.

(iii) $ge^X g^{-1} = e^{gXg^{-1}}$, $(e^X)^* = e^{X^*}$, $(e^X)^T = e^{X^T}$ and $\det e^X = e^{\mathrm{tr}\, X}$.

(iv) $\frac{d}{dt} e^{tX} = Xe^{tX} = e^{tX}X$.

(v) There are $r > 0$ and an open neighborhood V of 1 such that $\exp : B_{2r}(0) \to V$ is a diffeomorphism. We denote its inverse by $\log : V \to B_{2r}(0)$.

Proof. (i) $\|X^k/k!\| \leq \|X\|^k/k!$; the entries of e^X are power series in the $2n^2$ real coordinates converging everywhere, hence real analytic. (ii) By absolute convergence we may form the Cauchy product: $e^X e^Y = \sum_m \sum_{k+l=m} \frac{X^k Y^l}{k!l!} = \sum_m \frac{(X+Y)^m}{m!}$ by the binomial theorem, which needs $XY = YX$. (iii) Apply $A \mapsto gAg^{-1}$, $A \mapsto A^*$, $A \mapsto A^T$ termwise. For the determinant, write $X = UTU^{-1}$ with T upper triangular (Schur); e^T is upper triangular with diagonal entries $e^{T_{jj}}$, so $\det e^X = \det e^T = \prod_j e^{T_{jj}} = e^{\mathrm{tr}\, T} = e^{\mathrm{tr}\, X}$. (iv) Differentiate the power series in t termwise. (v) By (iv), $d\exp_0 = \mathrm{id}$; apply the inverse function theorem. □

Lemma 5.1.2 (Lie product formula): $e^{X+Y} = \lim_{k\to\infty} (e^{X/k} e^{Y/k})^k$.

Proof. By the power series, $A_k := e^{X/k}e^{Y/k} - 1 = \frac{X+Y}{k} + R_k$ with $\|R_k\| \leq C/k^2$. Since log is smooth near 1 with $d\log_1 = \mathrm{id}$, Taylor's theorem gives $\|\log(1+A) - A\| \leq C'\|A\|^2$ for small A. Hence for large k, $Z_k := \log(e^{X/k}e^{Y/k}) = \frac{X+Y}{k} + O(k^{-2})$, and $(e^{X/k}e^{Y/k})^k = e^{kZ_k} \to e^{X+Y}$ by continuity of exp, as $kZ_k \to X + Y$. □

Proposition 5.1.3 (One-parameter subgroups): Every continuous homomorphism $\gamma : \mathbb{R} \to \mathrm{GL}(n,\mathbb{C})$ is of the form $\gamma(t) = e^{tX}$ for a unique $X \in M_n(\mathbb{C})$.

Proof. Uniqueness: $X = \frac{d}{dt}|_{t=0} e^{tX}$. Existence: let r, V be as in Lemma 5.1.1(v), and choose $\varepsilon > 0$ with $\gamma([-\varepsilon,\varepsilon]) \subset \exp(B_r(0))$. Let $Y := \log\gamma(\varepsilon)$ and $Z := \log\gamma(\varepsilon/2)$, both in $B_r(0)$. Then $e^{2Z} = \gamma(\varepsilon/2)^2 = \gamma(\varepsilon) = e^Y$ with $2Z, Y \in B_{2r}(0)$, so $2Z = Y$. Inductively $\gamma(\varepsilon/2^m) = e^{Y/2^m}$, hence $\gamma(q\varepsilon/2^m) = e^{qY/2^m}$ for $q \in \mathbb{Z}$, $m \in \mathbb{N}$. Both sides of $\gamma(t) = e^{tY/\varepsilon}$ are continuous in t and agree on a dense set. □

5.2 The Lie algebra of a matrix Lie group

Definition 5.2.1: A real or complex *Lie algebra* is a vector space $\mathfrak{g}$ with a bilinear, antisymmetric map $[\cdot,\cdot]$ satisfying the Jacobi identity $[X,[Y,Z]]+[Y,[Z,X]]+[Z,[X,Y]]=0$. Homomorphisms preserve brackets, and $\operatorname{ad} X(Y) := [X,Y]$. A *representation* of $\mathfrak{g}$ on V is a homomorphism $\mathfrak{g} \to \mathfrak{gl}(V) = \operatorname{End}(V)$ (with the commutator).

Definition 5.2.2: For a closed subgroup $G \subset \mathrm{GL}(n,\mathbb{C})$, its *Lie algebra* is $\mathfrak{g} := \{X \in M_n(\mathbb{C}) \mid e^{tX} \in G \ \forall t \in \mathbb{R}\}$, and $\operatorname{Ad}(g)X := gXg^{-1}$ for $g \in G$.

If $X \in \mathfrak{g}$ and $g \in G$, then $e^{t \operatorname{Ad}(g)X} = ge^{tX}g^{-1} \in G$, so $\operatorname{Ad}(g)\mathfrak{g} \subset \mathfrak{g}$.

Theorem 5.2.1: Lie algebra of a closed subgroup

$\mathfrak{g}$ is a real Lie subalgebra of $M_n(\mathbb{C})$ with $[X,Y] = XY - YX$.

Proof. $\mathbb{R}\mathfrak{g} \subset \mathfrak{g}$ is clear. For $X, Y \in \mathfrak{g}$, $e^{t(X+Y)} = \lim_k (e^{tX/k}e^{tY/k})^k \in G$ by Lemma 5.1.2 and closedness, so $X+Y \in \mathfrak{g}$. Next, $t \mapsto \operatorname{Ad}(e^{tX})Y$ is a smooth curve in the closed subspace $\mathfrak{g}$, so its derivative at 0, which is $XY - YX$, lies in $\mathfrak{g}$. The Jacobi identity holds for commutators of matrices. □

Example 5.2.3:

G	$\mathfrak{g}$	$\dim_{\mathbb{R}} \mathfrak{g}$
$\mathrm{SL}(n,\mathbb{C}) = \{\det = 1\}$	$\mathfrak{sl}(n,\mathbb{C}) = \{X \mid \operatorname{tr} X = 0\}$	$2n^2 - 2$
$\mathrm{U}(n)$	$\mathfrak{u}(n) = \{X \mid X^* = -X\}$	n^2
$\mathrm{SU}(n)$	$\mathfrak{su}(n) = \{X \in \mathfrak{u}(n) \mid \operatorname{tr} X = 0\}$	$n^2 - 1$
$\mathrm{SO}(n)$	$\mathfrak{so}(n) = \{X \in M_n(\mathbb{R}) \mid X^T = -X\}$	$n(n-1)/2$
$\mathrm{Sp}(n)$	$\mathfrak{sp}(n) = \{X \in \mathfrak{u}(2n) \mid X^T J + JX = 0\}$	$n(2n+1)$
$\mathbb{T}^n$ (diagonal)	$\{\operatorname{diag}(i\theta_1, \ldots, i\theta_n) \mid \theta \in \mathbb{R}^n\}$	n

Proof. $\mathrm{U}(n)$: if $e^{tX} \in \mathrm{U}(n)$ for all t, differentiating $e^{tX^*}e^{tX} = 1$ at 0 gives $X^* + X = 0$; conversely if $X^* = -X$ then $(e^{tX})^* = e^{-tX} = (e^{tX})^{-1}$. The real dimension of skew-Hermitian matrices is n (imaginary diagonal) $+2\binom{n}{2}$ (upper triangle) $= n^2$. SL, SU: $\det e^{tX} = e^{t \operatorname{tr} X} = 1$ for all t iff $\operatorname{tr} X = 0$. $\mathrm{SO}(n)$: e^{tX} real for all t iff X real (derivative at 0), then argue as for $\mathrm{U}(n)$; $\det e^{tX} = e^{t \operatorname{tr} X} = 1$ automatically. $\mathrm{Sp}(n)$: if $(e^{tX})^T J e^{tX} = J$ for all t, differentiation gives $X^T J + JX = 0$; conversely $X^T J = -JX$ implies $(X^T)^k J = J(-X)^k$, hence $(e^{tX})^T J = Je^{-tX}$. Writing $X = \left(\begin{smallmatrix} A & B \\ C & D \end{smallmatrix}\right)$, one computes that $X^T J + JX = 0$ means $D = -A^T$, $B = B^T$, $C = C^T$, and $X^* = -X$ then means $A^* = -A$ and $C = -B^*$. So $\mathfrak{sp}(n) \cong \mathfrak{u}(n) \oplus \operatorname{Sym}_n(\mathbb{C})$, of real dimension $n^2 + n(n+1)$. The torus case is clear. □

Theorem 5.2.2: Closed subgroup theorem

Let $G \subset \mathrm{GL}(n,\mathbb{C})$ be closed with Lie algebra $\mathfrak{g}$. There are a neighborhood U of 0 in $\mathfrak{g}$ and an open neighborhood V of 1 in G such that $\exp : U \to V$ is a homeomorphism. Consequently G is an embedded submanifold of $\mathrm{GL}(n,\mathbb{C})$ of dimension $\dim_{\mathbb{R}} \mathfrak{g}$ with $T_1 G = \mathfrak{g}$, multiplication and inversion are smooth, and smooth maps into $\mathrm{GL}(n,\mathbb{C})$ with values in G are smooth into G.

Proof. Choose a real complement $\mathfrak{m}$ of $\mathfrak{g}$ in $M_n(\mathbb{C})$. The map $\Phi(X+Y) := e^X e^Y$ $(X \in \mathfrak{g}, Y \in \mathfrak{m})$ has $d\Phi_0 = \mathrm{id}$, so it restricts to a diffeomorphism from a neighborhood $U_1 \times W_1$ of 0 onto an open neighborhood O of 1. We claim that there is a neighborhood $W \subset W_1$ of 0 in $\mathfrak{m}$ such that $Y \in W$, $e^Y \in G$ imply $Y = 0$. Otherwise there are $0 \neq Y_k \to 0$ in $\mathfrak{m}$ with $e^{Y_k} \in G$; passing to a subsequence, $Y_k/\|Y_k\| \to Y \in \mathfrak{m}$ with $\|Y\| = 1$. For $t \in \mathbb{R}$ let $n_k := \lfloor t/\|Y_k\| \rfloor$; then $n_k Y_k \to tY$, so $e^{tY} = \lim (e^{Y_k})^{n_k} \in G$. Thus $Y \in \mathfrak{g} \cap \mathfrak{m} = \{0\}$, a contradiction.

Now shrink U_1, W_1 so that $W_1 \subset W$ and set $U := U_1$, $V := \Phi(U \times W) \cap G$, which is open in G. For $X \in U$, $Y \in W$: $e^X e^Y \in G \iff e^Y \in G \iff Y = 0$. Hence $V = \exp(U)$, and Φ is a chart of $\mathrm{GL}(n,\mathbb{C})$ near 1 in which G corresponds to the linear subspace $\mathfrak{g}$. Left translation by $g \in G$ transports this to a neighborhood of g. The remaining statements follow since G is an embedded submanifold. □

Proposition 5.2.4: Let G be a matrix Lie group. The subgroup G_0 generated by $\exp(\mathfrak{g})$ is open and closed in G, and it is the connected component of 1. If G is compact, G/G_0 is finite. If G is connected, G is generated by $\exp(\mathfrak{g})$.

Proof. G_0 contains the neighborhood V of 1 from Theorem 5.2.2, hence $hV \subset G_0$ for $h \in G_0$, so G_0 is open. Its complement is a union of cosets gG_0, which are open, so G_0 is closed. G_0 is path-connected: $t \mapsto e^{tX_1} \cdots e^{tX_k}$ connects 1 to any product of exponentials. Hence G_0 is the component of 1. If G is compact, the open cosets gG_0 cover G, so there are finitely many. □

5.3 Homomorphisms and derived representations

Theorem 5.3.1: Lie correspondence for homomorphisms

Let G, H be matrix Lie groups and $\Phi : G \to H$ a continuous homomorphism. There is a unique map $\phi = d\Phi : \mathfrak{g} \to \mathfrak{h}$ with $\Phi(e^X) = e^{\phi(X)}$ for all $X \in \mathfrak{g}$. It is a real-linear Lie algebra homomorphism, $\phi(X) = \frac{d}{dt}|_{t=0}\Phi(e^{tX})$, and $\phi(\mathrm{Ad}(g)X) = \mathrm{Ad}(\Phi(g))\phi(X)$.

Proof. $t \mapsto \Phi(e^{tX})$ is a continuous one-parameter subgroup, so $\Phi(e^{tX}) = e^{t\phi(X)}$ for a unique $\phi(X)$ (Proposition 5.1.3), which lies in $\mathfrak{h}$, and $\phi(X) = \frac{d}{dt}|_0\Phi(e^{tX})$. Clearly $\phi(sX) = s\phi(X)$. By Lemma 5.1.2, $\Phi(e^{t(X+Y)}) = \lim_k (e^{t\phi(X)/k}e^{t\phi(Y)/k})^k = e^{t(\phi(X)+\phi(Y))}$, so ϕ is additive. From $\Phi(ge^{tX}g^{-1}) = \Phi(g)e^{t\phi(X)}\Phi(g)^{-1}$ we get the equivariance. Applying it with $g = e^{sX}$ and differentiating at $s = 0$ (using linearity of ϕ) gives $\phi([X,Y]) = [\phi(X), \phi(Y)]$. □

Corollary 5.3.1: $\mathrm{Ad} : G \to \mathrm{GL}(\mathfrak{g})$ is a continuous homomorphism with $d\,\mathrm{Ad} = \mathrm{ad}$, i.e., $\mathrm{Ad}(e^X) = e^{\mathrm{ad}\,X}$.

Proof. $\frac{d}{dt}|_0 e^{tX}Ye^{-tX} = XY - YX$, and Theorem 5.3.1. □

Definition 5.3.2: For a representation (π, V) of a matrix Lie group G (viewing $\mathrm{GL}(V) \cong \mathrm{GL}(d,\mathbb{C})$), $d\pi : \mathfrak{g} \to \mathfrak{gl}(V)$ is the *derived representation*. Its complex-linear extension to the *complexification* $\mathfrak{g}_{\mathbb{C}} := \mathfrak{g} \otimes_{\mathbb{R}} \mathbb{C} = \mathfrak{g} \oplus i\mathfrak{g}$ (with the $\mathbb{C}$-bilinear extension of the bracket) is again denoted $d\pi$.

Proposition 5.3.3: Let G be a connected matrix Lie group and (π, V), (σ, W) representations. A subspace of V is G-invariant if and only if it is invariant under $d\pi(\mathfrak{g})$ (equivalently, under $d\pi(\mathfrak{g}_{\mathbb{C}})$), and $\mathrm{Hom}_G(V,W) = \{A \mid A\,d\pi(X) = d\sigma(X)A \ \forall X \in \mathfrak{g}\}$. In particular, π is irreducible iff $d\pi$ is.

Proof. If V' is $d\pi(\mathfrak{g})$-invariant, it is invariant under $\pi(e^X) = e^{d\pi(X)}$, hence under the group generated by $\exp(\mathfrak{g})$, which is G (Proposition 5.2.4). Conversely, if V' is G-invariant, it contains $d\pi(X)v = \lim_{t\to 0}(\pi(e^{tX})v - v)/t$ for $v \in V'$. The statement on intertwiners is proved in the same way. □

5.4 The representations of sl(2,C); SU(2) and SO(3)

We have $\mathfrak{su}(2)_{\mathbb{C}} = \mathfrak{sl}(2,\mathbb{C})$, since $M_n(\mathbb{C}) = \mathfrak{u}(n) \oplus i\mathfrak{u}(n)$ and the trace condition is preserved. Put

$$H = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad E = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad F = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \qquad [H,E] = 2E,\ [H,F] = -2F,\ [E,F] = H.$$

Note that iH, $E - F$ and $i(E + F)$ form a basis of $\mathfrak{su}(2)$.

Theorem 5.4.1: Representations of sl(2,C)

(i) For every $n \in \mathbb{N}_0$ there is, up to equivalence, exactly one irreducible representation $V^{(n)}$ of $\mathfrak{sl}(2,\mathbb{C})$ of dimension $n+1$. It has a basis $v_0, \dots, v_n$ with

$$Hv_k = (n-2k)v_k, \qquad Fv_k = v_{k+1}\ (v_{n+1} := 0), \qquad Ev_k = k(n-k+1)v_{k-1}.$$

(ii) Let V be a finite-dimensional representation of $\mathfrak{sl}(2,\mathbb{C})$ with an inner product for which $\mathfrak{su}(2)$ acts by skew-Hermitian operators. Then V is a direct sum of irreducible representations; in particular H is diagonalizable with integer eigenvalues.

Proof. (i) Let V be irreducible. From the relations, $Hv = \mu v$ implies $H(Ev) = (\mu+2)Ev$ and $H(Fv) = (\mu-2)Fv$. Starting from an eigenvector of H and applying E repeatedly, since H has finitely many eigenvalues we find $v_0 \neq 0$ with $Hv_0 = \mu v_0$, $Ev_0 = 0$. Let $v_k := F^k v_0$. Then $Hv_k = (\mu - 2k)v_k$, and by induction, using $EF = FE + H$,

$$Ev_k = EFv_{k-1} = F(k-1)(\mu-k+2)v_{k-2} + (\mu-2k+2)v_{k-1} = k(\mu-k+1)v_{k-1}.$$

The nonzero v_k are linearly independent (distinct eigenvalues of H); let n be maximal with $v_n \neq 0$. Then $0 = Ev_{n+1} = (n+1)(\mu-n)v_n$ gives $\mu = n$. The span of $v_0, \dots, v_n$ is invariant, hence equals V, and the formulas show that V is determined by $n = \dim V - 1$ up to equivalence. Conversely, the formulas define a representation: on v_k,

$$[E,F]v_k = (k+1)(n-k)v_k - k(n-k+1)v_k = (n-2k)v_k, \qquad [H,E]v_k = 2k(n-k+1)v_{k-1} = 2Ev_k,$$

and $[H,F]v_k = \big((n-2k-2) - (n-2k)\big)v_{k+1} = -2Fv_k$. It is irreducible: an invariant subspace $\neq 0$ contains an H-eigenvector, i.e., some v_k, and applying E and F yields all v_j.

(ii) The operators iH, $E-F$, $i(E+F)$ are skew-Hermitian, so $H^* = H$, and $E^* - F^* = F - E$, $E^* + F^* = E + F$, i.e., $E^* = F$. If $W \subset V$ is invariant, so is $W^\perp$: e.g. $\langle Ew', w\rangle = \langle w', Fw\rangle = 0$ for $w' \in W^\perp$, $w \in W$. Induction on $\dim V$ and (i) finish the proof. □

Lemma 5.4.1: $\mathrm{U}(n)$, $\mathrm{SU}(n)$ and $\mathrm{SO}(n)$ are connected.

Proof. If $g \in \mathrm{U}(n)$, then $g = k\,\mathrm{diag}(e^{i\theta_1}, \dots, e^{i\theta_n})k^{-1}$ with k unitary, and $s \mapsto k\,\mathrm{diag}(e^{is\theta_j})k^{-1}$ connects 1 to g in $\mathrm{U}(n)$. If $\det g = 1$ we may choose the θ_j with $\sum_j \theta_j = 0$, and the path stays in $\mathrm{SU}(n)$. If $g \in \mathrm{SO}(n)$, the real normal form of orthogonal matrices gives $k \in \mathrm{O}(n)$ with $k^{-1}gk$ block diagonal with blocks $R(\theta_j) = \left(\begin{smallmatrix} \cos\theta_j & -\sin\theta_j \\ \sin\theta_j & \cos\theta_j \end{smallmatrix}\right)$ and ± 1; as $\det g = 1$, the -1's come in pairs, which we write as $R(\pi)$. Then $s \mapsto k\,\mathrm{diag}(R(s\theta_j), 1, \dots, 1)k^{-1}$ is a path in $\mathrm{SO}(n)$ from 1 to g. □

Proposition 5.4.2 (SU(2) covers SO(3)): With the inner product $\langle X, Y\rangle := -2\,\mathrm{tr}(XY)$ on $\mathfrak{su}(2)$, $\mathrm{Ad} : \mathrm{SU}(2) \to \mathrm{SO}(\mathfrak{su}(2)) \cong \mathrm{SO}(3)$ is surjective with kernel $\{\pm 1\}$, and it induces an isomorphism of topological groups $\mathrm{SU}(2)/\{\pm 1\} \cong \mathrm{SO}(3)$. The irreducible representations of $\mathrm{SO}(3)$ correspond to the π_n with n even: writing $n = 2\ell$, they are the representations $D^{(\ell)}$, $\ell \in \mathbb{N}_0$, of dimension $2\ell + 1$.

Proof. $X_a := -\frac{i}{2}\sigma_a$ (σ_a the Pauli matrices) is an orthonormal basis, and $\mathrm{Ad}(g)$ preserves $\mathrm{tr}(XY)$, so $\mathrm{Ad}(g)$ is orthogonal; $\det \mathrm{Ad}(g) \in \{\pm 1\}$ depends continuously on $g \in \mathrm{SU}(2) \cong S^3$, which is connected, so $\mathrm{Ad}(g) \in \mathrm{SO}(3)$. If $\mathrm{Ad}(g) = 1$, then g commutes with $\mathfrak{su}(2)$ and hence with its complex span $M_2(\mathbb{C})$, so $g = \pm 1$. The derivative $\mathrm{ad} : \mathfrak{su}(2) \to \mathfrak{so}(3)$ is injective by the same argument and therefore bijective (both have dimension 3). By Corollary 5.3.1, $\mathrm{Ad}(\mathrm{SU}(2)) \supset \exp(\mathfrak{so}(3))$, which contains a neighborhood of 1 in SO(3) (Theorem 5.2.2); so the image is an open subgroup, hence closed, hence all of the connected group SO(3) (Lemma 5.4.1). The induced continuous bijection $\mathrm{SU}(2)/\{\pm 1\} \to \mathrm{SO}(3)$ from a compact to a Hausdorff space is a homeomorphism. Thus representations of SO(3) are the same as representations of SU(2) trivial on -1, with the same invariant subspaces; and $\pi_n(-1)P(z) = P(-z) = (-1)^n P(z)$. □

5.5 Compact Lie groups are matrix groups

Lemma 5.5.1 (No small subgroups): A Lie group G has a neighborhood V of e containing no subgroup other than $\{e\}$.

Proof. Choose $r > 0$ such that exp is injective on $B_{2r}(0) \subset \mathfrak{g}$ (for some norm on $\mathfrak{g}$) with open image, and put $V := \exp(B_r(0))$. Let $H \subset V$ be a subgroup and $e \neq h = \exp X \in H$ with $0 < \|X\| < r$. Choose $k \in \mathbb{N}$ with $r \leq k\|X\| < 2r$. Then $h^k = \exp(kX) \in H \subset V = \exp(B_r(0))$, which contradicts injectivity of exp on $B_{2r}(0)$. □

Remark 5.5.2: $\prod_{n\in\mathbb{N}} \mathbb{Z}/2\mathbb{Z}$ is a compact group which is not a Lie group: every neighborhood of e contains a subgroup $\{x \mid x_1 = \cdots = x_N = 0\} \neq \{e\}$.

Theorem 5.5.1: Compact Lie groups are linear

Every compact Lie group G has a faithful unitary representation, and is therefore isomorphic as a topological group to a closed subgroup of some $\mathrm{U}(N)$.

Proof. For $\pi \in \widehat{G}$, $K_\pi := \ker \pi$ is a closed subgroup, and $\bigcap_\pi K_\pi = \{e\}$ by Corollary 4.3.1(ii). Let V be an open neighborhood of e as in Lemma 5.5.1. The compact sets $K_\pi \setminus V$ have empty intersection, so finitely many of them, say for $\pi_1, \ldots, \pi_k$, already do. Then $\bigcap_j K_{\pi_j}$ is a subgroup contained in V, hence trivial, and $\pi := \pi_1 \oplus \cdots \oplus \pi_k$ (unitary by Theorem 2.2.1) is injective. A continuous injection of a compact space into a Hausdorff space is a homeomorphism onto its (compact, hence closed) image. □

6 Compact Lie algebras and the exponential map

6.1 Invariant inner products

Proposition 6.1.1: Let G be a compact matrix Lie group. There is an inner product on $\mathfrak{g}$ which is Ad-invariant. For it, every $\mathrm{ad}\, X$ is skew-symmetric, $\langle [X,Y], Z\rangle = -\langle Y, [X,Z]\rangle$, and hence diagonalizable over $\mathbb{C}$ with purely imaginary eigenvalues.

Proof. Average any inner product as in Theorem 2.2.1 (the argument works verbatim for real representations). Differentiating $\langle \mathrm{Ad}(e^{tX})Y, \mathrm{Ad}(e^{tX})Z\rangle = \langle Y, Z\rangle$ at $t = 0$ and using Corollary 5.3.1 gives skew-symmetry. A real skew-symmetric matrix is normal, hence unitarily diagonalizable, with eigenvalues λ satisfying $\bar{\lambda} = -\lambda$. □

Example 6.1.2: For $G \subset \mathrm{U}(n)$, $\langle X, Y\rangle := -\mathrm{tr}(XY) = \mathrm{tr}(XY^*)$ is Ad-invariant and positive definite on $\mathfrak{g}$.

Proposition 6.1.3: Let $\mathfrak{z}$ be the center of $\mathfrak{g}$ and fix an Ad-invariant inner product. Then $\mathfrak{g} = \mathfrak{z} \oplus [\mathfrak{g},\mathfrak{g}]$ orthogonally, $[\mathfrak{g},\mathfrak{g}]$ has trivial center, and the Killing form $B(X,Y) := \operatorname{tr}(\operatorname{ad} X \operatorname{ad} Y)$ is negative semidefinite with kernel $\mathfrak{z}$.

Proof. By skew-symmetry, $\langle Z,[X,Y]\rangle = -\langle [X,Z],Y\rangle$, which vanishes for all X,Y iff $Z \in \mathfrak{z}$; thus $[\mathfrak{g},\mathfrak{g}]^\perp = \mathfrak{z}$. An element of the center of $[\mathfrak{g},\mathfrak{g}]$ commutes with $[\mathfrak{g},\mathfrak{g}]$ and with $\mathfrak{z}$, hence lies in $\mathfrak{z} \cap [\mathfrak{g},\mathfrak{g}] = 0$. Finally $B(X,X) = \operatorname{tr}((\operatorname{ad} X)^2) = -\operatorname{tr}((\operatorname{ad} X)^T \operatorname{ad} X) \le 0$ with equality iff $\operatorname{ad} X = 0$, i.e., $X \in \mathfrak{z}$; since B is symmetric and semidefinite, its kernel is $\{X \mid B(X,X)=0\} = \mathfrak{z}$. □

6.2 The differential of the exponential map

Proposition 6.2.1: For $X, Y \in M_n(\mathbb{C})$,

$$e^{-X}\frac{d}{ds}\Big|_{s=0} e^{X+sY} = \frac{1-e^{-\operatorname{ad} X}}{\operatorname{ad} X} Y := \sum_{k\ge 0} \frac{(-1)^k}{(k+1)!} (\operatorname{ad} X)^k Y.$$

In particular, $d\exp_X$ is invertible if and only if $\operatorname{ad} X$ has no eigenvalue in $2\pi i\mathbb{Z}\setminus\{0\}$.

Proof. Let $Z(s) := X + sY$ and $\Gamma(s,t) := e^{-tZ(s)}\partial_s e^{tZ(s)}$. Then $\partial_t\Gamma = e^{-tZ}(-Z\partial_s e^{tZ} + \partial_s(Ze^{tZ})) = e^{-tZ}Ye^{tZ}$, which at $s=0$ equals $e^{-t\operatorname{ad} X}Y$ by Corollary 5.3.1 (applied in $\mathrm{GL}(n,\mathbb{C})$). Since $\Gamma(s,0)=0$, integrating over $t\in[0,1]$ gives the formula. The operator $\frac{1-e^{-\operatorname{ad} X}}{\operatorname{ad} X}$ has eigenvalues $\frac{1-e^{-z}}{z}$ (:= 1 for $z=0$) for the eigenvalues z of $\operatorname{ad} X$ (triangularize $\operatorname{ad} X$), and these vanish exactly for $z \in 2\pi i\mathbb{Z}\setminus\{0\}$. □

Remark 6.2.2: The exponential map of a connected matrix Lie group need not be surjective. For $A = \left(\begin{smallmatrix} -1 & 1 \\ 0 & -1 \end{smallmatrix}\right) \in \mathrm{SL}(2,\mathbb{R})$ there is no $X \in \mathfrak{sl}(2,\mathbb{R})$ with $e^X = A$: the eigenvalues of X are $\pm\lambda$ with $\lambda^2 = -\det X \in \mathbb{R}$. If $\lambda \in \mathbb{R}\setminus\{0\}$, e^X has the positive eigenvalues $e^{\pm\lambda}$; if $\lambda = 0$, X is nilpotent and $e^X = 1 + X$ has eigenvalue 1; if $\lambda \in i\mathbb{R}\setminus\{0\}$, X is diagonalizable (distinct eigenvalues), so e^X is diagonalizable, and having eigenvalues $-1,-1$ it would equal $-1 \ne A$. For compact connected groups, exp is surjective (Corollary 7.3.2).

7 Maximal tori

From now on, G denotes a compact connected matrix Lie group with Lie algebra $\mathfrak{g}$, equipped with a fixed Ad-invariant inner product $\langle\cdot,\cdot\rangle$ (Proposition 6.1.1).

7.1 Tori

Definition 7.1.1: A *torus* is a compact connected abelian matrix Lie group.

Lemma 7.1.2: Let Λ be a discrete subgroup of a finite-dimensional real vector space E. Then $\Lambda = \mathbb{Z}v_1 \oplus \cdots \oplus \mathbb{Z}v_m$ for linearly independent $v_1,\dots,v_m$.

Proof. Let $m := \dim \operatorname{span}\Lambda$ and choose $w_1,\dots,w_m \in \Lambda$ forming a basis of $\operatorname{span}\Lambda$. The bounded set $\{\sum c_j w_j \mid 0 \le c_j < 1\}$ contains only finitely many points of the discrete closed set Λ, and every coset of $\Lambda_0 := \bigoplus \mathbb{Z}w_j$ in Λ meets it; so $N := [\Lambda:\Lambda_0] < \infty$ and $\Lambda \subset \frac{1}{N}\Lambda_0 \cong \mathbb{Z}^m$. Subgroups of $\mathbb{Z}^m$ are free of rank $\le m$, and Λ spans an m-dimensional space, so a $\mathbb{Z}$-basis of Λ consists of m linearly independent vectors. □

Proposition 7.1.3: Let T be a torus of dimension k with Lie algebra $\mathfrak{t}$. Then $\exp : \mathfrak{t} \to T$ is a surjective homomorphism whose kernel Λ is a lattice of rank k, and $\mathfrak{t}/\Lambda \cong T$ as topological groups. In particular $T \cong \mathbb{R}^k/\mathbb{Z}^k \cong \mathbb{T}^k$.

Proof. T is abelian, so Ad is trivial and $\mathfrak{t}$ is abelian (Corollary 5.3.1); hence exp is a homomorphism (Lemma 5.1.1(ii)). Its image is a subgroup containing $\exp(\mathfrak{t})$, so it is T by Proposition 5.2.4. Since exp is injective near 0, Λ is discrete, so $\Lambda = \bigoplus_{j=1}^{m} \mathbb{Z} v_j$ by Lemma 7.1.2. The induced bijection $\mathfrak{t}/\Lambda \to T$ is a homomorphism and a local homeomorphism (Theorem 5.2.2), hence a homeomorphism. So $\mathfrak{t}/\Lambda \cong \mathbb{T}^m \times \mathbb{R}^{k-m}$ is compact, forcing $m = k$. □

Definition 7.1.4: For a torus $T = \exp(\mathfrak{t})$ with $\Lambda = \ker \exp$, the *weight lattice* is $P(T) := \{\lambda \in \mathfrak{t}^* \mid \lambda(\Lambda) \subset 2\pi\mathbb{Z}\}$. For $\lambda \in P(T)$ and $t = \exp H$ put $t^\lambda := e^{i\lambda(H)}$.

Proposition 7.1.5: $\lambda \mapsto (t \mapsto t^\lambda)$ is a bijection from $P(T)$ onto the set of irreducible representations (characters) of T, and $t^{\lambda+\mu} = t^\lambda t^\mu$. The characters t^λ, $\lambda \in P(T)$, form an orthonormal basis of $L^2(T)$.

Proof. Choose a $\mathbb{Z}$-basis $v_1, \dots, v_k$ of Λ; then $\mathbb{T}^k \to T$, $(e^{2\pi i s_j})_j \mapsto \exp(\sum_j s_j v_j)$ is an isomorphism. Under it, t^λ corresponds to $\prod_j z_j^{m_j}$ with $m_j = \lambda(v_j)/2\pi \in \mathbb{Z}$, and every $m \in \mathbb{Z}^k$ arises from a unique λ. Now apply Proposition 1.3.3 and Theorem 4.2.1(ii). □

Definition 7.1.6: $t \in T$ is a *generator* of T if $\{t^m \mid m \in \mathbb{Z}\}$ is dense in T.

Theorem 7.1.1: Kronecker

The element $(e^{2\pi i\theta_1}, \dots, e^{2\pi i\theta_k}) \in \mathbb{T}^k$ is a generator if and only if $1, \theta_1, \dots, \theta_k$ are linearly independent over $\mathbb{Q}$. Every torus has a generator.

Proof. Let t be the given element and S the closure of $\{t^m\}$, a closed subgroup. If $S \neq \mathbb{T}^k$, then $\mathbb{T}^k/S$ is a nontrivial compact abelian group, so by Corollary 4.3.1(ii) and Corollary 1.3.2 it has a nontrivial character; composing with $\mathbb{T}^k \to \mathbb{T}^k/S$ gives a character z^m, $m \neq 0$, with t^m-value $e^{2\pi i \sum m_j\theta_j} = 1$, i.e., $\sum_j m_j\theta_j \in \mathbb{Z}$, a rational relation. Conversely, a relation $\sum_j m_j\theta_j \in \mathbb{Z}$ with $m \neq 0$ gives a nontrivial character equal to 1 on all t^m, whose kernel is a proper closed subgroup containing $\{t^m\}$. Existence: since the $\mathbb{Q}$-span of finitely many reals is countable, we can choose inductively $\theta_j \notin \operatorname{span}_{\mathbb{Q}}\{1, \theta_1, \dots, \theta_{j-1}\}$. For a general torus use $T \cong \mathbb{T}^k$. □

7.2 Maximal tori and the Weyl group

Definition 7.2.1: A *maximal torus* of G is a torus $T \subset G$ not properly contained in another torus. Its dimension is the *rank* of G.

Lemma 7.2.2: (i) If $\mathfrak{a} \subset \mathfrak{g}$ is an abelian subalgebra, then the closure $\overline{\exp(\mathfrak{a})}$ is a torus with Lie algebra containing $\mathfrak{a}$.

(ii) If $T \subset T'$ are tori of the same dimension, then $T = T'$. In particular, maximal tori exist.

Proof. (i) $\exp(\mathfrak{a})$ is a connected abelian subgroup (Lemma 5.1.1(ii)), so its closure is a compact connected abelian closed subgroup of G. (ii) T is an open subgroup of T' by Theorem 5.2.2, hence closed, hence equal to T' as T' is connected. Thus any increasing chain of tori stabilizes, as dimensions are bounded by $\dim G$. □

Lemma 7.2.3: Let $T \subset G$ be a torus with Lie algebra $\mathfrak{t}$. Then T is a maximal torus if and only if

$$\mathfrak{g}^T := \{X \in \mathfrak{g} \mid \operatorname{Ad}(t)X = X \ \forall t \in T\} = \{X \in \mathfrak{g} \mid [H, X] = 0 \ \forall H \in \mathfrak{t}\}$$

equals $\mathfrak{t}$, i.e., iff $\mathfrak{t}$ is a maximal abelian subalgebra of $\mathfrak{g}$.

Proof. The two descriptions of $\mathfrak{g}^T$ agree: since $T = \exp(\mathfrak{t})$ and $\mathrm{Ad}(e^H) = e^{\mathrm{ad}\, H}$, X is fixed by $\mathrm{Ad}(T)$ iff $e^{s\,\mathrm{ad}\, H}X = X$ for all s, H, iff $[H, X] = 0$ for all $H \in \mathfrak{t}$. If T is maximal and $X \in \mathfrak{g}^T$, then $\mathfrak{t} + \mathbb{R}X$ is abelian, so by Lemma 7.2.2(i) $T \subset \overline{\exp(\mathfrak{t} + \mathbb{R}X)}$, a torus; by maximality they are equal, and $e^{sX} \in T$ for all s, i.e., $X \in \mathfrak{t}$. Conversely, if $\mathfrak{g}^T = \mathfrak{t}$ and $T \subset T'$ with T' a torus, then $\mathrm{Lie}(T')$ is abelian and contains $\mathfrak{t}$, so it lies in $\mathfrak{g}^T = \mathfrak{t}$; by Lemma 7.2.2(ii), $T' = T$. □

Example 7.2.4: The following are maximal tori.

(i) $\mathrm{U}(n)$: $T = \{\mathrm{diag}(e^{i\theta_1}, \dots, e^{i\theta_n})\}$; rank n.

(ii) $\mathrm{SU}(n)$: $T \cap \mathrm{SU}(n)$; rank $n - 1$.

(iii) $\mathrm{SO}(2n)$: $T = \{\mathrm{diag}(R(\theta_1), \dots, R(\theta_n))\}$; $\mathrm{SO}(2n+1)$: $T = \{\mathrm{diag}(R(\theta_1), \dots, R(\theta_n), 1)\}$; rank n.

(iv) $\mathrm{Sp}(n)$: $T = \{\mathrm{diag}(e^{i\theta_1}, \dots, e^{i\theta_n}, e^{-i\theta_1}, \dots, e^{-i\theta_n})\}$; rank n.

Proof. Each T is a compact connected abelian closed subgroup, hence a torus; we verify $\mathfrak{g}^T = \mathfrak{t}$ (Lemma 7.2.3). (i), (ii): $\mathfrak{t}$ contains a diagonal matrix H with distinct entries (with trace 0 for $\mathrm{SU}(n)$, $n \geq 2$); a matrix commuting with H preserves its eigenspaces, the coordinate axes, so it is diagonal. Diagonal skew-Hermitian (traceless) matrices form $\mathfrak{t}$. (iii) Take $H = \mathrm{diag}(\theta_1 J_2, \dots, \theta_n J_2)$ (plus a 0 for $\mathrm{SO}(2n+1)$), $J_2 = \left(\begin{smallmatrix} 0 & -1 \\ 1 & 0 \end{smallmatrix}\right)$, with $0 < \theta_1 < \cdots < \theta_n$. Its eigenspaces in $\mathbb{C}^{2n}$ (resp. $\mathbb{C}^{2n+1}$) for the distinct eigenvalues $\pm i\theta_j$ (and 0) are lines in the coordinate planes (resp. the last axis). A real matrix X commuting with H preserves these, hence preserves each coordinate plane, so X is block diagonal with 2×2 blocks commuting with J_2, i.e., of the form $a + bJ_2$ (and a 1×1 block). If $X^T = -X$, then $a = 0$ and the 1×1 block vanishes, so $X \in \mathfrak{t}$. (iv) A generic element of $\mathfrak{t}$ is diagonal with distinct entries $\pm i\theta_j$, so its centralizer in $\mathfrak{u}(2n)$ consists of diagonal matrices $\mathrm{diag}(a, b)$, and the condition $X^T J + JX = 0$ gives $b = -a$ (cf. the proof of Example 5.2.3); hence $\mathfrak{g}^T = \mathfrak{t}$. □

Definition 7.2.5: Let $T \subset G$ be a maximal torus and $N(T) := \{g \in G \mid gTg^{-1} = T\}$. The *Weyl group* is $W := N(T)/T$. It acts on T by $w \cdot t := ntn^{-1}$ ($w = nT$), on $\mathfrak{t}$ by $\mathrm{Ad}(n)$, and on $\mathfrak{t}^*$ by $(w\lambda)(H) := \lambda(\mathrm{Ad}(n)^{-1}H)$. These actions are well defined since T is abelian, and w preserves Λ and $P(T)$, with $(w \cdot t)^{w\lambda} = t^\lambda$.

Proposition 7.2.6: W is finite.

Proof. $N(T)$ is closed in G, hence compact, so $N(T)/N(T)_0$ is finite (Proposition 5.2.4). For $n \in N(T)$, $\mathrm{Ad}(n)|_{\mathfrak{t}}$ preserves Λ, so it lies in the discrete set $\{A \in \mathrm{GL}(\mathfrak{t}) \mid A\Lambda = \Lambda\}$. The continuous map $n \mapsto \mathrm{Ad}(n)|_{\mathfrak{t}}$ is therefore constant on the connected set $N(T)_0$, so $\mathrm{Ad}(n)|_{\mathfrak{t}} = \mathrm{id}$ for $n \in N(T)_0$. Thus $N(T)_0 \subset Z_G(T)$ and $\mathrm{Lie}(N(T)_0) \subset \mathfrak{g}^T = \mathfrak{t}$ (Lemma 7.2.3), so $N(T)_0 \subset T$ by Proposition 5.2.4. Hence W is a quotient of $N(T)/N(T)_0$. □

Example 7.2.7: For $G = \mathrm{U}(n)$, $W \cong S_n$, acting by permuting the diagonal entries. Indeed, permutation matrices normalize T. Conversely, if $n \in N(T)$ and $t \in T$ has distinct eigenvalues, then n maps the eigenspaces of t (the coordinate axes) to the eigenspaces of $ntn^{-1} \in T$, which are also coordinate axes; so n is a permutation matrix times a diagonal matrix in T.

7.3 The torus theorem

Lemma 7.3.1: Let T be a maximal torus of G. Every $X \in \mathfrak{g}$ is $\mathrm{Ad}(G)$-conjugate to an element of $\mathfrak{t}$.

Proof. Let $t = e^{H_0}$ be a generator of T (Theorem 7.1.1, Proposition 7.1.3). Then $\overline{\exp(\mathbb{R}H_0)} \supset \overline{\{t^m\}} = T$. The continuous function $f(g) := \langle \mathrm{Ad}(g)X, H_0\rangle$ on the compact group G attains its maximum at some g_0. Put $Y := \mathrm{Ad}(g_0)X$. For $Z \in \mathfrak{g}$, by Corollary 5.3.1 and skew-symmetry of ad,

$$0 = \frac{d}{ds}\Big|_{s=0} f(e^{sZ}g_0) = \langle [Z,Y], H_0\rangle = \langle Z, [Y, H_0]\rangle,$$

so $[H_0, Y] = 0$. Hence $\mathrm{Ad}(e^{sH_0})Y = e^{s\,\mathrm{ad}\,H_0}Y = Y$ for all s, so by continuity $\mathrm{Ad}(t')Y = Y$ for all $t' \in \overline{\exp(\mathbb{R}H_0)} = T$, i.e., $Y \in \mathfrak{g}^T = \mathfrak{t}$ (Lemma 7.2.3). □

Theorem 7.3.1: Torus theorem (É. Cartan)

Let G be a compact connected matrix Lie group and T a maximal torus. Then $G = \bigcup_{g\in G} gTg^{-1}$. Moreover, any two maximal tori are conjugate.

Proof. We use induction on $\dim G$; for $\dim G = 0$, $G = \{1\} = T$. Let $S := \bigcup_g gTg^{-1}$, the image of the compact space $G \times T$ under $(g,t) \mapsto gtg^{-1}$; so S is closed and nonempty. We show that S is open; then $S = G$ by connectedness. Since S is invariant under conjugation, it suffices to show that every $t_0 \in T$ has a neighborhood contained in S.

Case 1: t_0 is central in G. By Lemma 7.3.1, every $X \in \mathfrak{g}$ can be written as $X = \mathrm{Ad}(g)Y$ with $Y \in \mathfrak{t}$, so $t_0 e^X = g(t_0 e^Y)g^{-1} \in gTg^{-1}$. Hence S contains $t_0 \exp(\mathfrak{g})$, which is a neighborhood of t_0 (Theorem 5.2.2).

Case 2: t_0 is not central. Let $H := Z_G(t_0)_0$, a compact connected matrix Lie group with Lie algebra $\mathfrak{h} = \{X \in \mathfrak{g} \mid \mathrm{Ad}(t_0)X = X\}$. Since T is connected and commutes with t_0, $T \subset H$, and T is a maximal torus of H. If $\dim H = \dim G$, then H would be an open subgroup of G (Theorem 5.2.2), hence closed, hence equal to G, and t_0 would be central. So $\dim H < \dim G$, and by induction $H = \bigcup_{h\in H} hTh^{-1}$. Consider the smooth map $\mu : G\times H \to G$, $\mu(g,h) := ghg^{-1}$, whose image is contained in $\bigcup_g gHg^{-1} \subset S$. For $X \in \mathfrak{g}$, $Y \in \mathfrak{h}$,

$$t_0^{-1}\mu(e^{sX}, t_0 e^{sY}) = e^{s\,\mathrm{Ad}(t_0^{-1})X} e^{sY} e^{-sX}, \qquad \frac{d}{ds}\Big|_{s=0}(\ldots) = (\mathrm{Ad}(t_0^{-1}) - 1)X + Y.$$

Since $\mathrm{Ad}(t_0^{-1})$ is orthogonal, the image of $\mathrm{Ad}(t_0^{-1}) - 1$ is the orthogonal complement of $\ker(\mathrm{Ad}(t_0^{-1}) - 1) = \mathfrak{h}$. Hence the differential of $(g,h) \mapsto t_0^{-1}\mu(g,h)$ at $(1,t_0)$ is onto $\mathfrak{h}^\perp + \mathfrak{h} = \mathfrak{g}$. By the submersion theorem, $\mu(G\times H)$, and hence S, contains a neighborhood of t_0.

Finally, let T' be another maximal torus and t' a generator of T'. Then $t' = gtg^{-1}$ for some $g \in G$, $t \in T$, so $T' = \overline{\{t'^m\}} \subset gTg^{-1}$, and by maximality $T' = gTg^{-1}$. □

Corollary 7.3.2: Let G be compact connected with maximal torus T.

(i) $\exp : \mathfrak{g} \to G$ is surjective, and every element of G lies in a maximal torus.

(ii) Every $X \in \mathfrak{g}$ is $\mathrm{Ad}(G)$-conjugate to an element of $\mathfrak{t}$.

(iii) Two elements of T are conjugate in G if and only if they are conjugate under W.

(iv) A continuous class function on G is determined by its restriction to T, which is W-invariant.

Proof. (i) $g = hth^{-1} \in hTh^{-1}$, a maximal torus, and $t = e^H$ with $H \in \mathfrak{t}$ gives $g = e^{\mathrm{Ad}(h)H}$. (ii) is Lemma 7.3.1. (iv) follows from the theorem and $f(w\cdot t) = f(ntn^{-1}) = f(t)$. (iii) Let $s,t \in T$ with $gsg^{-1} = t$. Then T and gTg^{-1} are maximal tori of G contained in the compact connected group $H := Z_G(t)_0$ (both are connected and commute with t), hence maximal tori of H. By the theorem for H, there is $h \in H$ with $hgTg^{-1}h^{-1} = T$. So $n := hg \in N(T)$ and $nsn^{-1} = hth^{-1} = t$. □

7.4 Centralizers of tori

Lemma 7.4.1: Let A be a compact abelian matrix Lie group whose identity component A_0 satisfies: A/A_0 is cyclic. Then A has a generator, i.e., an element a with $\overline{\{a^m\}} = A$.

Proof. A_0 is a torus; let t_0 be a generator of A_0. Let xA_0 generate A/A_0, of order m, so $x^m \in A_0$. Since $\exp : \mathrm{Lie}(A_0) \to A_0$ is a surjective homomorphism, A_0 is divisible: there is $y \in A_0$ with $y^m = x^{-m}t_0$. Put $a := xy$. Then $a^m = t_0$, so $\overline{\{a^k\}} \supset \overline{\{t_0^k\}} = A_0$; and $a^k \in x^k A_0$ meets every coset. Hence $\overline{\{a^k\}} = A$. □

Proposition 7.4.2: Let $S \subset G$ be a torus. Then the centralizer $Z_G(S)$ is connected, with Lie algebra $\{X \in \mathfrak{g} \mid \mathrm{Ad}(s)X = X \ \forall s \in S\}$.

Proof. The Lie algebra statement is the definition of $\mathrm{Lie}(Z_G(S))$ together with $e^{tX}s = se^{tX} \iff e^{t\,\mathrm{Ad}(s^{-1})X} = e^{tX}$ for all $t \iff \mathrm{Ad}(s^{-1})X = X$. Let $x \in Z_G(S)$ and let A be the closure of the subgroup generated by x and S, a compact abelian group with $S \subset A_0$. As A/A_0 is finite, $x^m A_0 = A_0$ for some $m \geq 1$, so $\bigcup_k x^k A_0$ is a finite union of closed cosets, a closed subgroup containing x and S; hence it equals A, and A/A_0 is cyclic. By Lemma 7.4.1 A has a generator a, which by Corollary 7.3.2(i) lies in a maximal torus T'. Then $A \subset T'$, so T' is a connected subgroup of $Z_G(S)$ containing 1 and x. Hence $x \in Z_G(S)_0$. □

Corollary 7.4.3: (i) $Z_G(T) = T$ for a maximal torus T. Consequently, W acts faithfully on T and on $\mathfrak{t}$.

(ii) If $H \in \mathfrak{t}$ satisfies $\{X \in \mathfrak{g} \mid [H, X] = 0\} = \mathfrak{t}$, then every $g \in G$ with $\mathrm{Ad}(g)H = H$ lies in T.

Proof. (i) $Z_G(T)$ is connected with Lie algebra $\mathfrak{g}^T = \mathfrak{t}$, so it equals T (Proposition 5.2.4). If $n \in N(T)$ acts trivially on T, or on $\mathfrak{t}$ (then also on $T = \exp \mathfrak{t}$), then $n \in Z_G(T) = T$. (ii) $\mathrm{Ad}(g)H = H$ implies that g commutes with e^{sH} for all s, hence with the torus $S := \overline{\exp(\mathbb{R}H)}$. By Proposition 7.4.2, $Z_G(S)$ is connected, and its Lie algebra $\{X \mid \mathrm{Ad}(e^{sH})X = X \ \forall s\} = \{X \mid [H, X] = 0\} = \mathfrak{t}$. Since $T \subset Z_G(S)$, we get $Z_G(S) = T$ and $g \in T$. □

8 Roots and weights

Throughout, $T \subset G$ is a fixed maximal torus with Lie algebra $\mathfrak{t}$ and $\Lambda = \ker(\exp|_{\mathfrak{t}})$.

8.1 Weights of a representation

Definition 8.1.1: Let (π, V) be a representation of G. For $\lambda \in P(T)$ let $V_\lambda := \{v \in V \mid \pi(t)v = t^\lambda v \ \forall t \in T\}$. The λ with $V_\lambda \neq 0$ are the *weights* of π, and $m_\lambda := \dim V_\lambda$ their *multiplicities*.

Proposition 8.1.2: (i) $V = \bigoplus_\lambda V_\lambda$, and $d\pi(H)v = i\lambda(H)v$ for $H \in \mathfrak{t}$, $v \in V_\lambda$ (with $d\pi$ extended $\mathbb{C}$-linearly to $\mathfrak{t}_\mathbb{C}$).

(ii) $\chi_\pi(t) = \sum_\lambda m_\lambda t^\lambda$ for $t \in T$, and π is determined up to equivalence by its weights and multiplicities.

(iii) $m_{w\lambda} = m_\lambda$ for all $w \in W$.

Proof. (i) By Corollary 2.2.1, $\pi|_T$ is a sum of irreducible representations of T, which are characters t^λ (Proposition 7.1.5); grouping them gives $V = \bigoplus V_\lambda$. Differentiating $\pi(e^{sH})v = e^{is\lambda(H)}v$ gives the second claim. (ii) The trace formula follows from (i), and χ_π determines π (Corollary 3.2.3(ii), Corollary 7.3.2(iv)). (iii) For $w = nT$, $\chi_\pi(ntn^{-1}) = \chi_\pi(t)$ and $(ntn^{-1})^\lambda = t^{w^{-1}\lambda}$, so $\sum_\lambda m_\lambda t^{w^{-1}\lambda} = \sum_\lambda m_\lambda t^\lambda$. Since the characters of T are linearly independent (orthonormal), $m_{w\lambda} = m_\lambda$. □

Example 8.1.3: The weights of the defining representation of $\mathrm{U}(n)$ are $e_1, \dots, e_n$, where $e_j(\mathrm{diag}(i\theta_1, \dots, i\theta_n)) := \theta_j$. For $\mathrm{SU}(2)$ with $\mathfrak{t} = \{\mathrm{diag}(i\theta, -i\theta)\}$ and $\varepsilon(\mathrm{diag}(i\theta, -i\theta)) := \theta$, the weights of π_n are $(n-2k)\varepsilon$, $k = 0, \dots, n$ (proof of Theorem 3.3.1).

8.2 Roots

Let B be the complex-bilinear extension of the invariant inner product to $\mathfrak{g}_{\mathbb{C}}$ and $X \mapsto \bar{X}$ the complex conjugation of $\mathfrak{g}_{\mathbb{C}} = \mathfrak{g} \oplus i\mathfrak{g}$. Then $(X, Y)_h := B(X, \bar{Y})$ is a Hermitian inner product on $\mathfrak{g}_{\mathbb{C}}$, and for $Z \in \mathfrak{g}$, $\operatorname{ad} Z$ is skew-Hermitian for $(\cdot,\cdot)_h$ (Proposition 6.1.1). B is ad-invariant: $B([Z, X], Y) = -B(X, [Z, Y])$ for $X, Y, Z \in \mathfrak{g}_{\mathbb{C}}$.

For $\lambda \in \mathfrak{t}^*$ let $H_\lambda \in \mathfrak{t}$ be defined by $\langle H_\lambda, H\rangle = \lambda(H)$ for $H \in \mathfrak{t}$, and put $(\lambda, \mu) := \langle H_\lambda, H_\mu\rangle$, an inner product on $\mathfrak{t}^*$. Since W acts on $\mathfrak{t}$ by the orthogonal maps $\mathrm{Ad}(n)$, it acts on $\mathfrak{t}^*$ by orthogonal maps.

Definition 8.2.1: The nonzero weights of the representation Ad of G on $\mathfrak{g}_{\mathbb{C}}$ are the *roots*; their set is $\Delta \subset P(T)$. For $\alpha \in \Delta$, $\mathfrak{g}_\alpha := \{X \in \mathfrak{g}_{\mathbb{C}} \mid [H, X] = i\alpha(H)X \ \forall H \in \mathfrak{t}\}$ is the *root space*.

Proposition 8.2.2 (Root space decomposition): (i) $\mathfrak{g}_{\mathbb{C}} = \mathfrak{t}_{\mathbb{C}} \oplus \bigoplus_{\alpha\in\Delta} \mathfrak{g}_\alpha$, orthogonally for $(\cdot,\cdot)_h$.

(ii) $\overline{\mathfrak{g}_\alpha} = \mathfrak{g}_{-\alpha}$; in particular $\Delta = -\Delta$.

(iii) $[\mathfrak{g}_\alpha, \mathfrak{g}_\beta] \subset \mathfrak{g}_{\alpha+\beta}$, where $\mathfrak{g}_0 := \mathfrak{t}_{\mathbb{C}}$ and $\mathfrak{g}_\gamma := 0$ for $\gamma \notin \Delta \cup \{0\}$.

(iv) Δ is W-invariant.

Proof. (i) By Proposition 8.1.2(i) applied to Ad, $\mathfrak{g}_{\mathbb{C}}$ is the sum of its weight spaces, which for nonzero weights are the $\mathfrak{g}_\alpha$ (as $\mathrm{Ad}(e^H) = e^{\operatorname{ad} H}$). The zero weight space is the complexification of $\mathfrak{g}^T = \mathfrak{t}$ (Lemma 7.2.3). Orthogonality: the $\operatorname{ad} H$, $H \in \mathfrak{t}$, are commuting skew-Hermitian operators, and the weight spaces are their joint eigenspaces for distinct eigenvalue functions. (ii) Conjugating $[H, X] = i\alpha(H)X$ ($H \in \mathfrak{t}$ is real) gives $[H, \bar{X}] = -i\alpha(H)\bar{X}$. (iii) Jacobi identity. (iv) Proposition 8.1.2(iii). □

Example 8.2.3 (Roots of U(n)): Here $\mathfrak{g}_{\mathbb{C}} = \mathfrak{gl}(n, \mathbb{C})$, and for $H = \mathrm{diag}(i\theta_1, \dots, i\theta_n)$, $[H, E_{jk}] = i(\theta_j - \theta_k)E_{jk}$. Hence $\Delta = \{e_j - e_k \mid j \neq k\}$ and $\mathfrak{g}_{e_j - e_k} = \mathbb{C}E_{jk}$. With $\langle X, Y\rangle = -\operatorname{tr}(XY)$, the e_j are orthonormal in $\mathfrak{t}^*$.

8.3 The SU(2)-subalgebras attached to roots

Definition 8.3.1: For $\alpha \in \Delta$, the *coroot* is $\alpha^\vee := \frac{2}{(\alpha,\alpha)} H_\alpha \in \mathfrak{t}$. Thus $\alpha(\alpha^\vee) = 2$ and $\lambda(\alpha^\vee) = \frac{2(\lambda,\alpha)}{(\alpha,\alpha)}$ for $\lambda \in \mathfrak{t}^*$.

Proposition 8.3.2: Let $\alpha \in \Delta$.

(i) For $X \in \mathfrak{g}_\alpha$ and $Y \in \mathfrak{g}_{-\alpha}$: $[X, Y] = iB(X, Y)H_\alpha$.

(ii) There is $X_\alpha \in \mathfrak{g}_\alpha$ such that, with $H' := -i\alpha^\vee$, $E' := X_\alpha$, $F' := -\bar{X}_\alpha$, the linear map $\varphi_\alpha : \mathfrak{sl}(2, \mathbb{C}) \to \mathfrak{g}_{\mathbb{C}}$, $H \mapsto H'$, $E \mapsto E'$, $F \mapsto F'$, is an injective Lie algebra homomorphism with $\varphi_\alpha(\mathfrak{su}(2)) \subset \mathfrak{g}$. Explicitly, $\varphi_\alpha(iH) = \alpha^\vee$ and $\varphi_\alpha(E - F) = X_\alpha + \bar{X}_\alpha$.

(iii) $\dim_{\mathbb{C}} \mathfrak{g}_\alpha = 1$, and $\mathbb{R}\alpha \cap \Delta = \{\pm\alpha\}$.

Proof. (i) $[X, Y] \in \mathfrak{t}_{\mathbb{C}}$ by Proposition 8.2.2(iii). For $H \in \mathfrak{t}$, by invariance of B,

$$B(H, [X, Y]) = -B([X, H], Y) = B([H, X], Y) = i\alpha(H)B(X, Y) = B\big(H,\, iB(X, Y)H_\alpha\big),$$

and B is nondegenerate on $\mathfrak{t}_{\mathbb{C}}$.

(ii) For $0 \neq X \in \mathfrak{g}_\alpha$, $B(X, \bar{X}) = (X, X)_h > 0$; rescale X to get X_α with $B(X_\alpha, \bar{X}_\alpha) = 2/(\alpha, \alpha)$. Then by (i), using $\bar{X}_\alpha \in \mathfrak{g}_{-\alpha}$,

$$[E', F'] = -[X_\alpha, \bar{X}_\alpha] = -i\tfrac{2}{(\alpha,\alpha)} H_\alpha = H', \qquad [H', E'] = -i[\alpha^\vee, X_\alpha] = -i \cdot i\alpha(\alpha^\vee) X_\alpha = 2E',$$

and, since $[\alpha^\vee, \bar{X}_\alpha] = -i\alpha(\alpha^\vee)\bar{X}_\alpha = -2i\bar{X}_\alpha$, $[H', F'] = i[\alpha^\vee, \bar{X}_\alpha] = 2\bar{X}_\alpha = -2F'$. So φ_α is a homomorphism; it is injective because H', E', F' are linearly independent (they lie in different summands of Proposition 8.2.2(i)). Finally $\varphi_\alpha(iH) = \alpha^\vee$, $\varphi_\alpha(E - F) = X_\alpha + \bar{X}_\alpha$ and $\varphi_\alpha(i(E + F)) = i(X_\alpha - \bar{X}_\alpha)$ are fixed by conjugation, hence lie in $\mathfrak{g}$.

(iii) Let $\mathfrak{s}_\alpha := \varphi_\alpha(\mathfrak{sl}(2, \mathbb{C}))$ and

$$M := \mathbb{C}\alpha^\vee \oplus \bigoplus_{c \in \mathbb{R}\setminus\{0\},\ c\alpha \in \Delta} \mathfrak{g}_{c\alpha}.$$

M is invariant under $\mathrm{ad}(\mathfrak{s}_\alpha)$: $\mathrm{ad}\, H'$ preserves each summand; $\mathrm{ad}\, E'$ maps $\mathfrak{g}_{c\alpha}$ into $\mathfrak{g}_{(c+1)\alpha}$ for $c \neq -1$, maps $\mathfrak{g}_{-\alpha}$ into $\mathbb{C}\alpha^\vee$ by (i), and maps $\alpha^\vee$ into $\mathfrak{g}_\alpha$; similarly for F'. Since $\varphi_\alpha(\mathfrak{su}(2)) \subset \mathfrak{g}$ acts by skew-Hermitian operators, Theorem 5.4.1(ii) applies: M is a direct sum of irreducible representations $V^{(n)}$, and H' acts on $\mathfrak{g}_{c\alpha}$ by $-i \cdot ic\,\alpha(\alpha^\vee) = 2c$ and on $\mathbb{C}\alpha^\vee$ by 0. In particular $2c \in \mathbb{Z}$.

$\mathfrak{s}_\alpha \subset M$ is invariant (it is a subalgebra), with H'-eigenvalues $2, 0, -2$. In a decomposition of $\mathfrak{s}_\alpha$ into irreducibles, the summand containing an eigenvector for 2 is some $V^{(n)}$ with $n \geq 2$, of dimension $\geq 3 = \dim \mathfrak{s}_\alpha$; hence $\mathfrak{s}_\alpha \cong V^{(2)}$ is irreducible. Let M' be an invariant complement, $M = \mathfrak{s}_\alpha \oplus M'$ (orthogonal complement). The zero eigenspace of H' on M is $\mathbb{C}\alpha^\vee \subset \mathfrak{s}_\alpha$, so H' has no eigenvalue 0 on M'. Since $V^{(n)}$ has the eigenvalue 0 iff n is even, M' is a sum of $V^{(n)}$ with n odd, and all H'-eigenvalues on M' are odd. Therefore $M' \cap \mathfrak{g}_{c\alpha} = 0$ for $c \in \mathbb{Z}$. Since H'-eigenspaces are compatible with the decomposition $M = \mathfrak{s}_\alpha \oplus M'$, we get $\mathfrak{g}_\alpha = \mathfrak{s}_\alpha \cap \mathfrak{g}_\alpha = \mathbb{C}X_\alpha$ and $\mathfrak{g}_{2\alpha} = \mathfrak{s}_\alpha \cap \mathfrak{g}_{2\alpha} = 0$. As α was arbitrary: *twice a root is never a root.* If $M' \neq 0$, then (odd n) H' has the eigenvalue 1 on M', so $\mathfrak{g}_{\alpha/2} \neq 0$ and $\alpha/2 \in \Delta$; but then $\alpha = 2 \cdot \frac{\alpha}{2}$ is twice a root, a contradiction. Hence $M = \mathfrak{s}_\alpha$, and the only nonzero real multiples of α in Δ are $\pm\alpha$. □

Corollary 8.3.3: $\dim \mathfrak{g} = \dim \mathfrak{t} + |\Delta|$.

Corollary 8.3.4 (Integrality): For every weight λ of a representation of G and every $\alpha \in \Delta$, $\lambda(\alpha^\vee) = \frac{2(\lambda,\alpha)}{(\alpha,\alpha)} \in \mathbb{Z}$. In particular $\beta(\alpha^\vee) \in \mathbb{Z}$ for $\alpha, \beta \in \Delta$.

Proof. Choose an invariant inner product on V (Theorem 2.2.1); then $d\pi(\mathfrak{g})$ consists of skew-Hermitian operators, so $d\pi \circ \varphi_\alpha$ satisfies the hypothesis of Theorem 5.4.1(ii), and H acts with integer eigenvalues. On V_λ, $d\pi(\varphi_\alpha(H)) = d\pi(-i\alpha^\vee)$ acts by $-i \cdot i\lambda(\alpha^\vee) = \lambda(\alpha^\vee)$. □

Lemma 8.3.5: Every $\lambda \in P(T)$ is a weight of some irreducible representation of G. Consequently $\lambda(\alpha^\vee) \in \mathbb{Z}$ for all $\lambda \in P(T)$ and $\alpha \in \Delta$.

Proof. For a representation (π, V) let $P_\lambda := \int_T \overline{t^\lambda} \pi(t)\, dt$. On V_μ it acts by $\int_T t^\mu \overline{t^\lambda}\, dt = \delta_{\lambda\mu}$ (Proposition 7.1.5), so P_λ is the projection onto V_λ along the other weight spaces. Suppose λ is not a weight of any representation. Then $P_\lambda = 0$ always, so for every matrix coefficient $f = f^\pi_{u,v}$ and every $g \in G$,

$$\int_T f(gt)\overline{t^\lambda}\, dt = \langle \pi(g) P_\lambda u, v \rangle = 0.$$

By Theorem 4.2.1(i) this holds for all $f \in C(G)$. By Tietze's extension theorem there is $f \in C(G)$ with $f|_T = t^\lambda$; for $g = e$ we get $\int_T |t^\lambda|^2\, dt = 1 \neq 0$, a contradiction. The last statement now follows from Corollary 8.3.4. □

8.4 Reflections and the root system

Definition 8.4.1: For $\alpha \in \Delta$, the *reflection* $s_\alpha : \mathfrak{t}^* \to \mathfrak{t}^*$ is $s_\alpha(\lambda) := \lambda - \lambda(\alpha^\vee)\alpha = \lambda - \frac{2(\lambda,\alpha)}{(\alpha,\alpha)}\alpha$, the orthogonal reflection in the hyperplane $\alpha^\perp$.

Proposition 8.4.2: For $\alpha \in \Delta$ let $n_\alpha := \exp\big(\frac{\pi}{2}(X_\alpha + \bar{X}_\alpha)\big) \in G$. Then $n_\alpha \in N(T)$, and $w_\alpha := n_\alpha T \in W$ acts on $\mathfrak{t}^*$ by s_α. In particular $s_\alpha \in W$.

Proof. Let $Z := X_\alpha + \bar{X}_\alpha = \varphi_\alpha(E - F) \in \mathfrak{g}$, so $\mathrm{Ad}(n_\alpha) = e^{\frac{\pi}{2}\operatorname{ad} Z}$ (Corollary 5.3.1). For $H \in \ker\alpha \subset \mathfrak{t}$, $[Z, H] = -i\alpha(H)X_\alpha + i\alpha(H)\bar{X}_\alpha = 0$, so $\mathrm{Ad}(n_\alpha)H = H$. The subalgebra $\mathfrak{s}_\alpha$ is $\operatorname{ad} Z$-invariant, and since φ_α is a homomorphism, $\operatorname{ad} Z \circ \varphi_\alpha = \varphi_\alpha \circ \operatorname{ad}(E - F)$. Hence

$$\mathrm{Ad}(n_\alpha)\alpha^\vee = e^{\frac{\pi}{2}\operatorname{ad} Z}\varphi_\alpha(iH) = \varphi_\alpha\big(e^{\frac{\pi}{2}\operatorname{ad}(E-F)}iH\big) = \varphi_\alpha\big(i\,wHw^{-1}\big) = -\alpha^\vee,$$

where $w := e^{\frac{\pi}{2}(E-F)} = \left(\begin{smallmatrix} 0 & 1 \\ -1 & 0 \end{smallmatrix}\right)$ (as $(E-F)^2 = -1$) and $e^{\frac{\pi}{2}\operatorname{ad}(E-F)} = \mathrm{Ad}(w)$ (Corollary 5.3.1 for $\mathrm{GL}(2,\mathbb{C})$). So $\mathrm{Ad}(n_\alpha)$ preserves $\mathfrak{t} = \ker\alpha \oplus \mathbb{R}\alpha^\vee$, whence $n_\alpha T n_\alpha^{-1} = \exp(\mathrm{Ad}(n_\alpha)\mathfrak{t}) = T$. On $\mathfrak{t}$, $\mathrm{Ad}(n_\alpha)$ is the reflection r fixing $\ker\alpha$ and negating $\alpha^\vee$; thus $(w_\alpha\lambda)(H) = \lambda(rH)$ equals $\lambda(H)$ on $\ker\alpha$ and $-\lambda(\alpha^\vee)$ for $H = \alpha^\vee$, which are also the values of $s_\alpha\lambda$. □

Corollary 8.4.3: Δ is a *reduced root system* in $E := \operatorname{span}_{\mathbb{R}}\Delta \subset \mathfrak{t}^*$, i.e., a finite subset of $E \setminus \{0\}$ such that (i) $s_\alpha\Delta = \Delta$, (ii) $\beta(\alpha^\vee) \in \mathbb{Z}$, and (iii) $\mathbb{R}\alpha \cap \Delta = \{\pm\alpha\}$ for all $\alpha, \beta \in \Delta$.

Proof. Propositions 8.4.2 and 8.2.2(iv), Corollary 8.3.4 and Proposition 8.3.2(iii). □

8.5 Positive roots, simple roots and dominant weights

Fix $H_0 \in \mathfrak{t}$ with $\alpha(H_0) \neq 0$ for all $\alpha \in \Delta$, and with $\mu(H_0) \neq \nu(H_0)$ for all distinct $\mu, \nu \in P(T)$ (possible since $P(T)$ is countable and each condition excludes a hyperplane). Put $\Delta^+ := \{\alpha \in \Delta \mid \alpha(H_0) > 0\}$, the *positive roots*; then $\Delta = \Delta^+ \sqcup (-\Delta^+)$, and we write $\alpha > 0$ for $\alpha \in \Delta^+$. A positive root is *simple* if it is not the sum of two positive roots.

Lemma 8.5.1: (i) If $\alpha, \beta \in \Delta$, $\beta \neq \pm\alpha$ and $(\alpha, \beta) > 0$, then $\alpha - \beta \in \Delta$.

(ii) Every positive root is a sum of simple roots (with repetitions).

(iii) If $\alpha \neq \beta$ are simple, then $(\alpha, \beta) \leq 0$.

(iv) The simple roots $\alpha_1, \ldots, \alpha_r$ are linearly independent; hence they form a basis of E, and every root is $\sum_j c_j\alpha_j$ with all $c_j \in \mathbb{N}_0$ or all $c_j \in -\mathbb{N}_0$.

Proof. (i) $\alpha(\beta^\vee)\beta(\alpha^\vee) = \frac{4(\alpha,\beta)^2}{(\alpha,\alpha)(\beta,\beta)}$ is a product of two positive integers (Corollary 8.3.4) and is < 4 by the strict Cauchy–Schwarz inequality (α, β are not proportional by Proposition 8.3.2(iii)). So one factor is 1. If $\alpha(\beta^\vee) = 1$, then $s_\beta\alpha = \alpha - \beta \in \Delta$; if $\beta(\alpha^\vee) = 1$, then $s_\alpha\beta = \beta - \alpha \in \Delta$, so $\alpha - \beta \in \Delta$. (ii) Induction on the finitely many values $\alpha(H_0) > 0$: a non-simple positive root is $\beta + \gamma$ with $\beta, \gamma > 0$ and $\beta(H_0), \gamma(H_0) < \alpha(H_0)$. (iii) Otherwise $\alpha - \beta \in \Delta$ by (i). If $\alpha - \beta > 0$, then $\alpha = \beta + (\alpha - \beta)$ is not simple; if $\beta - \alpha > 0$, then β is not simple. (iv) Suppose $\sum_j c_j\alpha_j = 0$. Let $v := \sum_{c_j>0} c_j\alpha_j = \sum_{c_k<0}(-c_k)\alpha_k$. Then $(v, v) = \sum_{c_j>0,\, c_k<0} c_j(-c_k)(\alpha_j, \alpha_k) \leq 0$ by (iii), so $v = 0$; evaluating at H_0 gives $\sum_{c_j>0} c_j\alpha_j(H_0) = 0$, so no c_j is positive; similarly none is negative. The last statement follows from (ii) and $\Delta = \Delta^+ \sqcup -\Delta^+$. □

Definition 8.5.2: (i) The *fundamental Weyl chamber* is $C := \{\lambda \in \mathfrak{t}^* \mid (\lambda, \alpha) > 0\ \forall \alpha \in \Delta^+\}$. The *Weyl chambers* are the connected components of $\mathfrak{t}^*_{\mathrm{reg}} := \{\lambda \mid (\lambda, \alpha) \neq 0\ \forall \alpha \in \Delta\}$; $\lambda \in \mathfrak{t}^*_{\mathrm{reg}}$ is called *regular*.

(ii) $\lambda \in P(T)$ is *dominant* if $(\lambda, \alpha) \geq 0$ for all $\alpha > 0$, and *strictly dominant* if $(\lambda, \alpha) > 0$ for all $\alpha > 0$. P_+ denotes the set of dominant weights.

(iii) $\rho := \frac{1}{2}\sum_{\alpha>0}\alpha$ is the *Weyl vector*.

(iv) $Q_+ := \{\sum_j c_j\alpha_j \mid c_j \geq 0\}$, and $\mu \preceq \lambda$ means $\lambda - \mu \in Q_+$.

By Lemma 8.5.1(ii), λ is dominant iff $(\lambda, \alpha_j) \geq 0$ for all simple roots, and $C = \{\lambda \mid (\lambda, \alpha_j) > 0\ \forall j\}$. Each Weyl chamber is the set of λ with a fixed pattern of signs of (λ, α), $\alpha \in \Delta^+$ (such a set is convex, hence connected, and open, and different sign patterns cannot be connected within $\mathfrak{t}^*_{\mathrm{reg}}$). C is a chamber, as H_0 corresponds to an element of it. Moreover $Q_+ \cap (-Q_+) = \{0\}$ and $\mu(H_0) > 0$ for $0 \neq \mu \in Q_+$, by Lemma 8.5.1(iv).

Lemma 8.5.3: Let α_i be simple. Then s_{α_i} permutes $\Delta^+ \setminus \{\alpha_i\}$ and $s_{\alpha_i}\rho = \rho - \alpha_i$. Hence $\rho(\alpha_i^\vee) = 1$, i.e., $(\rho, \alpha_i) = \frac{1}{2}(\alpha_i, \alpha_i)$, and $\rho \in C$.

Proof. Let $\beta = \sum_j c_j\alpha_j \in \Delta^+ \setminus \{\alpha_i\}$. By Proposition 8.3.2(iii), β is not a multiple of α_i, so $c_j > 0$ for some $j \neq i$. The root $s_{\alpha_i}\beta = \beta - \beta(\alpha_i^\vee)\alpha_i$ has the same coefficient $c_j > 0$, so it is positive by Lemma 8.5.1(iv), and $\neq \alpha_i$. As $s_{\alpha_i}\alpha_i = -\alpha_i$, $s_{\alpha_i}\rho = \rho - \alpha_i$; comparing with $s_{\alpha_i}\rho = \rho - \rho(\alpha_i^\vee)\alpha_i$ gives $\rho(\alpha_i^\vee) = 1$. Finally, $(\rho, \alpha_i) > 0$ for all i means $\rho \in C$. □

9 The Weyl group

9.1 W as a reflection group

Let $s_i := s_{\alpha_i}$ for the simple roots $\alpha_1, \ldots, \alpha_r$. Recall that W acts on $\mathfrak{t}^*$ by orthogonal maps permuting Δ (Proposition 8.2.2(iv)); hence it permutes the Weyl chambers.

Theorem 9.1.1: Structure of the Weyl group

(i) W acts simply transitively on the set of Weyl chambers.

(ii) W is generated by $s_1, \ldots, s_r$; in particular it is generated by the reflections s_α, $\alpha \in \Delta$.

(iii) If $\lambda \in \overline{C}$ and $w \in W$, then $\lambda - w\lambda \in Q_+$. If $\lambda \in C$ and $w \neq 1$, then $\lambda - w\lambda \neq 0$, so $(w\lambda)(H_0) < \lambda(H_0)$.

(iv) Every $\lambda \in \mathfrak{t}^*$ is W-conjugate to exactly one element of $\overline{C}$. W acts freely on the regular elements of $\mathfrak{t}^*$.

Proof. Step 1. Let $W' \subset W$ be the subgroup generated by $s_1, \ldots, s_r$ (Proposition 8.4.2). For $\lambda \in \mathfrak{t}^*$ choose $w \in W'$ maximizing $(w\lambda, \rho)$ over the finite set W'. If $(w\lambda, \alpha_i) < 0$ for some i, then by Lemma 8.5.3

$$(s_i w\lambda, \rho) = (w\lambda, s_i\rho) = (w\lambda, \rho) - (w\lambda, \alpha_i) > (w\lambda, \rho),$$

a contradiction. So $w\lambda \in \overline{C}$; if λ is regular, so is $w\lambda$, and $w\lambda \in C$. Hence W' acts transitively on chambers.

Step 2: if $w \in W$ and $wC = C$, then $w = 1$. For $\mu \in C$ and $\alpha > 0$, $(\mu, w\alpha) = (w^{-1}\mu, \alpha) > 0$, so $w\Delta^+ = \Delta^+$ and $w\rho = \rho$. Write $w = nT$; then $\mathrm{Ad}(n)H_\rho = H_\rho$. Since $\alpha(H_\rho) = (\alpha, \rho) \neq 0$ for all $\alpha \in \Delta$, Proposition 8.2.2(i) shows that $\{X \in \mathfrak{g} \mid [H_\rho, X] = 0\} = \mathfrak{t}$. By Corollary 7.4.3(ii), $n \in T$, i.e., $w = 1$.

(i) Transitivity is Step 1. If $wC' = C'$ for a chamber $C' = uC$, $u \in W$, then $u^{-1}wuC = C$, so $w = 1$ by Step 2. (ii) For $w \in W$ choose $w' \in W'$ with $w'wC = C$ (Step 1); then $w'w = 1$ by Step 2, so $w \in W'$.

(iii) For $w \in W$ let $N(w) := |\{\beta \in \Delta^+ \mid w\beta < 0\}|$. If $w \neq 1$, there is a simple root with $w\alpha_i < 0$: otherwise w maps every positive root $\sum c_j\alpha_j$ to a root in Q_+, i.e., to a positive root, so $wC = C$ and $w = 1$. Since s_i permutes $\Delta^+ \setminus \{\alpha_i\}$ and $ws_i\alpha_i = -w\alpha_i > 0$, we get

$N(ws_i) = N(w) - 1$. We prove the claim by induction on $N(w)$; $N(w) = 0$ forces $w = 1$. For $w \neq 1$ choose i as above. Since $s_i\lambda = \lambda - \lambda(\alpha_i^\vee)\alpha_i$,

$$\lambda - w\lambda = (\lambda - ws_i\lambda) - \lambda(\alpha_i^\vee)\, w\alpha_i.$$

The first term is in Q_+ by induction, and $-w\alpha_i \in \Delta^+$ and $\lambda(\alpha_i^\vee) \geq 0$, so $\lambda - w\lambda \in Q_+$. If $\lambda \in C$, then $\lambda(\alpha_i^\vee) > 0$, and a sum $a + b$ with $a \in Q_+$, $0 \neq b \in Q_+$ is nonzero (since $Q_+ \cap -Q_+ = 0$). Finally, $\mu(H_0) > 0$ for $0 \neq \mu \in Q_+$.

(iv) Existence is Step 1. If $\lambda, \mu \in \overline{C}$ and $\mu = w\lambda$, then $\lambda - \mu \in Q_+$ and $\mu - \lambda = \mu - w^{-1}\mu \in Q_+$ by (iii), so $\lambda = \mu$. If λ is regular and $w\lambda = \lambda$, then w fixes the chamber containing λ, so $w = 1$ by (i). □

Definition 9.1.1: The *sign* of $w \in W$ is $\varepsilon(w) := \det(w|_{\mathfrak{t}^*}) \in \{\pm 1\}$. It is a homomorphism with $\varepsilon(s_\alpha) = -1$.

Lemma 9.1.2: For $w \in W$, $\rho - w\rho = \sum_{\beta>0,\ w^{-1}\beta<0} \beta$. In particular $w\rho - \rho \in P(T)$, so $\rho + P(T)$ is W-invariant.

Proof. $w\rho = \frac{1}{2}\sum_{\alpha>0} w\alpha$. As α runs through Δ^+, $w\alpha$ runs through the positive roots β with $w^{-1}\beta > 0$ and the negatives of the positive roots β with $w^{-1}\beta < 0$. Hence $w\rho = \rho - \sum_{\beta>0,\, w^{-1}\beta<0} \beta$. Roots lie in $P(T)$, and W preserves $P(T)$. □

9.2 Alternating sums and the Weyl denominator

For $\mu \in \mathfrak{t}^*$ let $e^\mu : \mathfrak{t} \to \mathbb{C}$, $e^\mu(H) := e^{i\mu(H)}$. For distinct $\mu_1, \ldots, \mu_m$ these functions are linearly independent: choose H with distinct values $\mu_j(H)$; then $s \mapsto e^{is\mu_j(H)}$ are linearly independent functions of $s \in \mathbb{R}$ (consider derivatives at $s = 0$ and a Vandermonde determinant). W acts on functions on $\mathfrak{t}$ by $(w \cdot f)(H) := f(\mathrm{Ad}(n)^{-1}H)$, so that $w \cdot e^\mu = e^{w\mu}$. For $\mu \in \mathfrak{t}^*$ define the *alternating sum*

$$A_\mu := \sum_{w \in W} \varepsilon(w)\, e^{w\mu}.$$

If $\mu \in P(T)$, then $e^\mu(H) = (e^H)^\mu$ is a function on T.

Lemma 9.2.1: Let $L \subset \mathfrak{t}^*$ be a W-invariant subset and $f = \sum_{\mu \in L} c_\mu e^\mu$ (finite sum) with $w \cdot f = \varepsilon(w) f$ for all $w \in W$. Then $f = \sum_{\nu \in L \cap C} c_\nu A_\nu$.

Proof. By linear independence, $w \cdot f = \varepsilon(w) f$ means $c_{w\mu} = \varepsilon(w) c_\mu$. If μ is not regular, then $(\mu, \alpha) = 0$ for some α, so $s_\alpha \mu = \mu$ and $c_\mu = -c_\mu = 0$. By Theorem 9.1.1(iv), the regular $\mu \in L$ are partitioned into orbits $\{w\nu\}_{w \in W}$ with unique $\nu \in C \cap L$ and $|W|$ distinct elements; the corresponding part of f is $c_\nu A_\nu$. □

Lemma 9.2.2: If $\nu, \nu' \in C$ and $\nu - \nu' \in P(T)$ and $w\nu - \nu \in P(T)$ for all w, then $|A_\nu|^2$ and $A_\nu \overline{A_{\nu'}}$ are functions on T, and $\int_T A_\nu \overline{A_{\nu'}}\, dt = |W| \delta_{\nu\nu'}$.

Proof. $A_\nu \overline{A_{\nu'}} = \sum_{w,w'} \varepsilon(w)\varepsilon(w') e^{w\nu - w'\nu'}$ with $w\nu - w'\nu' = (w\nu - \nu) + (\nu - \nu') + (\nu' - w'\nu') \in P(T)$, so each term is a character of T. By Proposition 7.1.5, the integral counts the pairs with $w\nu = w'\nu'$. If $\nu \neq \nu'$ there are none (Theorem 9.1.1(iv)); if $\nu = \nu'$, exactly the pairs $w = w'$ (free action). □

Proposition 9.2.3 (Weyl denominator formula): As functions on $\mathfrak{t}$,

$$A_\rho = \sum_{w \in W} \varepsilon(w)\, e^{w\rho} = e^\rho \prod_{\alpha>0} \left(1 - e^{-\alpha}\right) = \prod_{\alpha>0} \left(e^{\alpha/2} - e^{-\alpha/2}\right).$$

Consequently, for $t = e^H \in T$, $|A_\rho(H)|^2 = \prod_{\alpha>0} |1 - t^\alpha|^2 =: |\Delta(t)|^2$ depends only on t, and $\int_T |\Delta(t)|^2\, dt = |W|$.

Proof. Let $D := \prod_{\alpha>0}(e^{\alpha/2} - e^{-\alpha/2}) = e^{\rho}\prod_{\alpha>0}(1 - e^{-\alpha})$. By Lemma 8.5.3, $s_i \cdot D$ is the same product with the factor for α_i replaced by $e^{-\alpha_i/2} - e^{\alpha_i/2}$; so $s_i \cdot D = -D$. By Theorem 9.1.1(ii), $w \cdot D = \varepsilon(w)D$ for all w. Expanding, $D = \sum_S c_S e^{\rho - \sum_{\alpha\in S}\alpha}$ over subsets $S \subset \Delta^+$, with exponents in the W-invariant set $L := \rho + P(T)$ (Lemma 9.1.2); the exponent ρ occurs only for $S = \emptyset$ (as $\sum_{\alpha\in S}\alpha$ is positive on H_0 for $S \neq \emptyset$), with coefficient 1. By Lemma 9.2.1, $D = \sum_{\nu\in C\cap L} c_\nu A_\nu$, where only ν of the form $\rho - \sum_{\alpha\in S}\alpha$ occur. For such ν, $\nu(\alpha_i^\vee) = 1 - \sum_{\alpha\in S}\alpha(\alpha_i^\vee) \in \mathbb{Z}$ (Corollary 8.3.4) and > 0, hence $\geq 1 = \rho(\alpha_i^\vee)$. So $\eta := \nu - \rho$ satisfies $(\eta, \alpha_i) \geq 0$ for all i, while $-\eta = \sum_{\alpha\in S}\alpha = \sum_i c_i\alpha_i$ with $c_i \geq 0$. Then $(\eta,\eta) = -\sum_i c_i(\eta, \alpha_i) \leq 0$, so $\eta = 0$. Hence $D = A_\rho$. Since $|e^{\alpha/2} - e^{-\alpha/2}| = |1 - e^{\alpha}|$ on $\mathfrak{t}$, $|A_\rho(H)|^2 = \prod_{\alpha>0}|1 - t^\alpha|^2$. Finally apply Lemma 9.2.2 with $\nu = \nu' = \rho$. □

9.3 Regular elements

Definition 9.3.1: $t \in T$ is *regular* if $t^\alpha \neq 1$ for all $\alpha \in \Delta$. Let T^{reg} be the set of regular elements of T and $G^{\mathrm{reg}} := \bigcup_g gT^{\mathrm{reg}}g^{-1}$.

Lemma 9.3.2: (i) $\dim Z_G(t) = \dim T + |\{\alpha \in \Delta \mid t^\alpha = 1\}|$ for $t \in T$. Hence $t \in T$ is regular iff $\dim Z_G(t) = \dim T$, and then $Z_G(t)_0 = T$. In particular, $G^{\mathrm{reg}} \cap T = T^{\mathrm{reg}}$.

(ii) $T \setminus T^{\mathrm{reg}}$ is a null set in T, and $G \setminus G^{\mathrm{reg}}$ is a null set in G.

(iii) If $s, t \in T^{\mathrm{reg}}$ and $nsn^{-1} = t$ with $n \in G$, then $n \in N(T)$.

Proof. (i) $\mathrm{Lie}(Z_G(t)) = \{X \in \mathfrak{g} \mid \mathrm{Ad}(t)X = X\}$, whose complexification is $\mathfrak{t}_{\mathbb{C}} \oplus \bigoplus_{t^\alpha=1}\mathfrak{g}_\alpha$ by Proposition 8.2.2, and $\dim \mathfrak{g}_\alpha = 1$. If t is regular, $T \subset Z_G(t)_0$ have the same dimension, so T is open in the connected group $Z_G(t)_0$, hence equal to it. The dimension of $Z_G(t)$ is invariant under conjugation, which gives the last claim. (ii) $T \setminus T^{\mathrm{reg}} = \bigcup_{\alpha>0} K_\alpha$ with $K_\alpha := \{t \mid t^\alpha = 1\}$, a closed subgroup with Lie algebra $\ker\alpha$, hence a finite union of submanifolds of dimension $\dim T - 1$; such sets are null for Haar measure of T (which is a smooth density in charts). By the torus theorem, $G \setminus G^{\mathrm{reg}}$ is the image of the compact manifold-pieces $G/T \times K_\alpha$, of dimension $\dim G - 1$, under the smooth map $q(gT, t) := gtg^{-1}$; so it is null. (iii) $nZ_G(s)_0n^{-1} = Z_G(t)_0$, i.e., $nTn^{-1} = T$ by (i). □

9.4 The Weyl integration formula

We use the manifold G/T (see the prerequisites) and the smooth map $q : G/T \times T \to G$, $q(gT, t) := gtg^{-1}$, which is well defined as T is abelian. Let $\mathfrak{p} := \mathfrak{t}^\perp \subset \mathfrak{g}$; then $\mathrm{Ad}(t)\mathfrak{p} = \mathfrak{p}$ for $t \in T$, and $\mathfrak{p}_{\mathbb{C}} = \bigoplus_{\alpha\in\Delta}\mathfrak{g}_\alpha$. The differential of $G \to G/T$ at e restricts to an isomorphism $\mathfrak{p} \to T_{eT}(G/T)$ (its kernel is $\mathfrak{t}$).

Lemma 9.4.1 (Invariant measures): Fix Lebesgue measures on $\mathfrak{p}$, $\mathfrak{t}$ and the product measure on $\mathfrak{g} = \mathfrak{p} \oplus \mathfrak{t}$.

(i) There are unique smooth positive measures (densities) ω_G on G and ω_T on T, invariant under left translations, which at e are the given Lebesgue measures on $\mathfrak{g}$ and $\mathfrak{t}$. They are positive multiples of the Haar measures.

(ii) There is a unique smooth positive density $\omega_{G/T}$ on G/T, invariant under the action of G, which at eT is the given Lebesgue measure on $\mathfrak{p} \cong T_{eT}(G/T)$.

(iii) There is $c > 0$ with $\int_{G/T} h\, d\omega_{G/T} = c\int_G h(gT)\, dg$ for all $h \in C(G/T)$.

Proof. (i) Transport the density at e by left translations; the resulting measure is left invariant, so its normalization is Haar measure by uniqueness (Theorem 2.1.1). (ii) For $g \in G$ let $\ell_g(xT) :=$

gxT. Define $\omega_{G/T}$ at gT as the image of the density at eT under $d\ell_g$. This is independent of the representative: for $s \in T$, $d\ell_s$ at eT corresponds to $\mathrm{Ad}(s)|_{\mathfrak{p}}$ (differentiate $se^{uX}T = e^{u\,\mathrm{Ad}(s)X}sT = e^{u\,\mathrm{Ad}(s)X}T$), which has $|\det| = 1$ as $\mathrm{Ad}(s)$ is orthogonal. Smoothness follows from the existence of smooth local sections of $G \to G/T$, and invariance is clear. (iii) For $f \in C(G)$, $g \mapsto \int_T f(gs)\,ds$ is continuous and right T-invariant, so it defines $\tilde{f} \in C(G/T)$. The functional $f \mapsto \int_{G/T} \tilde{f}\,d\omega_{G/T}$ is positive, left invariant and nonzero; by the uniqueness part of Theorem 2.1.1 it is $c\int_G f\,dg$ with $c = \omega_{G/T}(G/T)$. For $f(g) := h(gT)$, $\tilde{f} = h$. □

Lemma 9.4.2 (Jacobian of q): $q^*\omega_G = J(t)\,\omega_{G/T} \otimes \omega_T$ with

$$J(t) = \big|\det\big((\mathrm{Ad}(t^{-1}) - 1)|_{\mathfrak{p}}\big)\big| = \prod_{\alpha\in\Delta}(t^{-\alpha} - 1) = |\Delta(t)|^2.$$

In particular q is a local diffeomorphism near every point of $G/T \times T^{\mathrm{reg}}$.

Proof. Fix (g_0T, t_0). The map $\psi(X,Y) := (g_0e^XT, t_0e^Y)$, $(X,Y) \in \mathfrak{p} \times \mathfrak{t}$ near 0, is a local diffeomorphism whose differential at 0 maps the Lebesgue measure on $\mathfrak{p} \times \mathfrak{t}$ to $\omega_{G/T} \otimes \omega_T$ (by construction). The map $Z \mapsto q(g_0T, t_0)e^Z$, $Z \in \mathfrak{g}$ near 0, has the analogous property for ω_G. Now

$$\begin{aligned} q(g_0T,t_0)^{-1}\,q(\psi(sX,sY)) &= g_0t_0^{-1}e^{sX}t_0e^{sY}e^{-sX}g_0^{-1} \\ &= \exp\Big(s\,\mathrm{Ad}(g_0)\big((\mathrm{Ad}(t_0^{-1}) - 1)X + Y\big) + O(s^2)\Big). \end{aligned}$$

So in these coordinates the differential of q is $(X,Y) \mapsto \mathrm{Ad}(g_0)\big((\mathrm{Ad}(t_0^{-1}) - 1)X + Y\big)$. Its absolute determinant is $|\det \mathrm{Ad}(g_0)| \cdot |\det((\mathrm{Ad}(t_0^{-1}) - 1)|_{\mathfrak{p}})| = J(t_0)$, as $\mathrm{Ad}(g_0)$ is orthogonal and $(\mathrm{Ad}(t_0^{-1}) - 1)$ maps $\mathfrak{p}$ to $\mathfrak{p}$. On $\mathfrak{p}_{\mathbb{C}} = \bigoplus_\alpha \mathfrak{g}_\alpha$, $\mathrm{Ad}(t_0^{-1})$ acts on $\mathfrak{g}_\alpha$ by $t_0^{-\alpha}$, so $\det((\mathrm{Ad}(t_0^{-1}) - 1)|_{\mathfrak{p}}) = \prod_{\alpha\in\Delta}(t_0^{-\alpha} - 1) = \prod_{\alpha>0}(\overline{t_0^{\alpha}} - 1)(t_0^{\alpha} - 1) = \prod_{\alpha>0}|1 - t_0^{\alpha}|^2 \geq 0$, which equals $|\Delta(t_0)|^2$ by Proposition 9.2.3. It is nonzero iff t_0 is regular, and then q is a local diffeomorphism by the inverse function theorem. □

Lemma 9.4.3: Let $M := G/T \times T^{\mathrm{reg}}$ and $N := G^{\mathrm{reg}}$. Then $q^{-1}(N) = M$, every fibre $q^{-1}(y)$, $y \in N$, has exactly $|W|$ points, and every $y \in N$ has an open neighborhood V such that $q^{-1}(V)$ is a disjoint union of $|W|$ open sets, each mapped diffeomorphically onto V by q.

Proof. If $q(gT,t) \in N$, then t is conjugate to an element of T^{reg}, hence regular by Lemma 9.3.2(i); so $q^{-1}(N) = M$. Fix $(g_1T, t_1) \in q^{-1}(y)$. If $q(gT,t) = y$, then $n := g_1^{-1}g$ satisfies $ntn^{-1} = t_1$, so $n \in N(T)$ by Lemma 9.3.2(iii), and $(gT,t) = (g_1nT, n^{-1}t_1n)$. Conversely these points lie in the fibre, and they are distinct for distinct $nT \in W$. So $|q^{-1}(y)| = |W|$.

Let $q^{-1}(y) = \{x_1, \ldots, x_k\}$. By Lemma 9.4.2 choose disjoint open $U_j \ni x_j$ mapped diffeomorphically onto open sets $q(U_j)$. We claim that there is an open $V \ni y$ with $V \subset \bigcap_j q(U_j)$ and $q^{-1}(V) \subset \bigcup_j U_j$. Otherwise there are $z_m \notin \bigcup_j U_j$ with $q(z_m) \to y$; by compactness of $G/T \times T$ a subsequence converges to some z with $q(z) = y$, i.e., $z = x_j$ for some j, contradicting $z_m \notin U_j$ with U_j open. Then the sets $U_j \cap q^{-1}(V)$ have the required properties. □

Theorem 9.4.1: Weyl integration formula

Let G be compact connected with maximal torus T, and let dg, dt be the normalized Haar measures. For every $f \in C(G)$, and more generally for every Borel function $f \geq 0$ and every $f \in L^1(G)$,

$$\int_G f(g)\,dg = \frac{1}{|W|}\int_G\int_T f(gtg^{-1})\,|\Delta(t)|^2\,dt\,dg.$$

In particular, for a class function f (continuous, or Borel and ≥ 0, or integrable),

$$\int_G f(g)\,dg = \frac{1}{|W|}\int_T f(t)\,|\Delta(t)|^2\,dt.$$

Proof. Let $F \in C(G)$ with compact support in $N = G^{\mathrm{reg}}$. If the support lies in a set V as in Lemma 9.4.3, then by the change of variables formula on each of the $|W|$ sheets and Lemma 9.4.2,

$$\int_M (F\circ q)\,J\,d(\omega_{G/T}\otimes\omega_T) = |W|\int_N F\,d\omega_G.$$

By a partition of unity this holds for all $F \in C_c(N)$. For $F \in C(G)$ choose $\varphi_k \in C_c(N)$ with $0 \leq \varphi_k \uparrow 1_N$ pointwise (possible as N is open in the compact metric space G), apply the formula to $F\varphi_k$ and use dominated convergence; since $G \setminus N$ and $(G/T)\times(T\setminus T^{\mathrm{reg}})$ are null sets (Lemma 9.3.2(ii)), we obtain

$$\int_{G/T}\int_T F(gtg^{-1})J(t)\,d\omega_T(t)\,d\omega_{G/T}(gT) = |W|\int_G F\,d\omega_G.$$

The inner integral is a continuous function of gT (since $(gs)t(gs)^{-1} = gtg^{-1}$ for $s \in T$). By Lemma 9.4.1 and $J = |\Delta|^2$, there is a constant $\kappa > 0$ with

$$\int_G F(g)\,dg = \kappa\int_G\int_T F(gtg^{-1})|\Delta(t)|^2\,dt\,dg \qquad (F \in C(G)).$$

Taking $F = 1$ and using $\int_T |\Delta|^2\,dt = |W|$ (Proposition 9.2.3) gives $\kappa = 1/|W|$.

The right-hand side, as a functional of F, is integration against the image of the finite Borel measure $\frac{1}{|W|}|\Delta(t)|^2\,dt\,dg$ on $G\times T$ under $(g,t)\mapsto gtg^{-1}$. This is a finite Borel measure on the compact metrizable space G, hence regular, and it agrees with Haar measure on $C(G)$; by the uniqueness in the Riesz–Markov theorem the two measures coincide. This gives the formula for Borel $f \geq 0$ and integrable f. For class functions, $f(gtg^{-1}) = f(t)$. □

Example 9.4.4: For SU(2): $W \cong \mathbb{Z}/2$, $\Delta^+ = \{\alpha\}$ with $t_\theta^\alpha = e^{2i\theta}$, $|\Delta(t_\theta)|^2 = |1-e^{2i\theta}|^2 = 4\sin^2\theta$, and Theorem 9.4.1 reproduces Lemma 3.3.2: for a class function f,

$$\frac{1}{2}\cdot\frac{1}{2\pi}\int_0^{2\pi} f(t_\theta)\,4\sin^2\theta\,d\theta = \frac{2}{\pi}\int_0^{\pi} f(t_\theta)\sin^2\theta\,d\theta,$$

where we substituted $\theta\mapsto 2\pi-\theta$ on $[\pi,2\pi]$ and used $f(t_{-\theta}) = f(t_\theta)$.

Example 9.4.5 (Random unitary matrices): For $G = \mathrm{U}(n)$, $W = S_n$ (Example 7.2.7) and $|\Delta|^2 = \prod_{j<k}|e^{i\theta_j}-e^{i\theta_k}|^2$. So the eigenvalue angles of a Haar-random unitary matrix have joint density $\frac{1}{n!(2\pi)^n}\prod_{j<k}|e^{i\theta_j}-e^{i\theta_k}|^2$ on $[0,2\pi)^n$ (circular unitary ensemble).

10 The Weyl character formula

Definition 10.0.1: The *highest weight* of a representation π of G is the weight λ of π with maximal value $\lambda(H_0)$ (unique, since $\mu\mapsto\mu(H_0)$ is injective on $P(T)$).

For a representation π we regard $\chi_\pi\circ\exp$ as a function on $\mathfrak{t}$; by Proposition 8.1.2(ii), $\chi_\pi\circ\exp = \sum_\mu m_\mu e^\mu$, and it is W-invariant by Corollary 7.3.2(iv). Recall that $A_\rho(H)\neq 0$ iff e^H is regular (Proposition 9.2.3).

Theorem 10.0.1: Weyl character formula and theorem of the highest weight

Let G be a compact connected Lie group with maximal torus T and positive roots Δ^+.

(i) The highest weight λ of an irreducible representation π is dominant and has multiplicity 1, and for all $H \in \mathfrak{t}$

$$\chi_\pi(e^H)\, A_\rho(H) = A_{\lambda+\rho}(H), \quad \text{i.e.} \quad \chi_\pi(e^H) = \frac{\sum_{w\in W} \varepsilon(w) e^{i(w(\lambda+\rho))(H)}}{\sum_{w\in W} \varepsilon(w) e^{i(w\rho)(H)}}$$

whenever $e^H \in T^{\mathrm{reg}}$.

(ii) Two irreducible representations are equivalent if and only if they have the same highest weight.

(iii) For every $\lambda \in P_+$ there is an irreducible representation with highest weight λ.

Hence $\lambda \mapsto \pi_\lambda$ is a bijection from P_+ onto $\widehat{G}$.

Proof. Let $L := \rho + P(T)$, a W-invariant set (Lemma 9.1.2).

(i) The function $f := (\chi_\pi \circ \exp) \cdot A_\rho = \sum_{\mu,w} m_\mu \varepsilon(w) e^{\mu+w\rho}$ has exponents in L and integer coefficients, and is anti-invariant. By Lemma 9.2.1, $f = \sum_{\nu\in C\cap L} c_\nu A_\nu$ with $c_\nu \in \mathbb{Z}$. Now $|f(H)|^2 = |\chi_\pi(t)|^2 |\Delta(t)|^2$ for $t = e^H$, so by Theorems 3.2.1 and 9.4.1 and Lemma 9.2.2,

$$1 = \int_G |\chi_\pi|^2\, dg = \frac{1}{|W|} \int_T |f|^2\, dt = \sum_\nu c_\nu^2.$$

Hence $f = \pm A_\nu$ for a single $\nu \in C \cap L$. Compare the exponents with maximal value at H_0. In $\pm A_\nu$ it is ν, uniquely, with coefficient ± 1 (Theorem 9.1.1(iii)). In f, $(\mu + w\rho)(H_0) \le \lambda(H_0) + \rho(H_0)$, with equality only if $\mu(H_0) = \lambda(H_0)$ and $(w\rho)(H_0) = \rho(H_0)$, i.e., $\mu = \lambda$ and $w = 1$ (Theorem 9.1.1(iii), $\rho \in C$). So the top exponent of f is $\lambda+\rho$ with coefficient $m_\lambda \ge 1$. Therefore $\nu = \lambda + \rho$, the sign is $+$, and $m_\lambda = 1$. As $\lambda + \rho \in C$, $(\lambda+\rho)(\alpha_i^\vee) \ge 1$ is an integer (Lemma 8.3.5) and $\rho(\alpha_i^\vee) = 1$, so $\lambda(\alpha_i^\vee) \ge 0$ for all i: λ is dominant.

(ii) By (i), the highest weight determines χ_π on T^{reg}, which is dense in T (its complement is a finite union of closed subgroups of lower dimension, Lemma 9.3.2), hence on T by continuity, hence on G (Corollary 7.3.2(iv)), hence π (Corollary 3.2.3(ii)).

(iii) Suppose $\lambda \in P_+$ is not the highest weight of any irreducible representation. For $L' \in \Lambda$ and $\nu \in L$, $A_\nu(H + L') = e^{i\nu(L')} A_\nu(H)$, since $w\nu - \nu \in P(T)$. Hence $A_{\lambda+\rho}/A_\rho$ is Λ-periodic on $\{H \mid e^H \in T^{\mathrm{reg}}\}$ (the factors $e^{i(\lambda+\rho)(L')}$ and $e^{i\rho(L')}$ differ by $e^{i\lambda(L')} = 1$), and W-invariant (quotient of two anti-invariant functions). Define $f : G \to \mathbb{C}$ by $f(gtg^{-1}) := A_{\lambda+\rho}(H)/A_\rho(H)$ for $t = e^H \in T^{\mathrm{reg}}$, and $f := 0$ on $G \setminus G^{\mathrm{reg}}$. This is well defined by Corollary 7.3.2(iii), it is a class function, and it is continuous on the open set G^{reg} since locally $f = \phi \circ \mathrm{pr}_2 \circ (q|_{U_j})^{-1}$ with ϕ continuous (Lemma 9.4.3). By Theorem 9.4.1 and Lemma 9.2.2 (note $\lambda + \rho \in C \cap L$),

$$\|f\|_2^2 = \frac{1}{|W|} \int_T |A_{\lambda+\rho}|^2\, dt = 1.$$

If π is irreducible with highest weight μ, then $\mu \ne \lambda$ by assumption, and by (i), Theorem 9.4.1 (for the integrable class function $f\overline{\chi_\pi}$) and Lemma 9.2.2,

$$\langle f, \chi_\pi \rangle_{L^2} = \frac{1}{|W|} \int_T A_{\lambda+\rho} \overline{A_{\mu+\rho}}\, dt = 0.$$

This contradicts Theorem 4.2.1(iv). □

Remark 10.0.2: Part (iii) is Weyl's analytic proof via Peter–Weyl, as in Simon's book. Algebraic (Verma modules) and geometric (Borel–Weil) constructions of π_λ also exist; we do not need them.

Proposition 10.0.3: Let π be irreducible with highest weight λ. Every weight μ of π satisfies $\mu \preceq \lambda$.

Proof. For $\beta \in \Delta^+$ fix $0 \neq X_\beta \in \mathfrak{g}_\beta$ and $Y_\beta := \bar{X}_\beta \in \mathfrak{g}_{-\beta}$. By Proposition 8.2.2(iii), $d\pi(\mathfrak{g}_\gamma)V_\mu \subset V_{\mu+\gamma}$. Let $0 \neq v \in V_\lambda$. Then $d\pi(X_\beta)v = 0$ for $\beta > 0$, since $(\lambda+\beta)(H_0) > \lambda(H_0)$. Let U be the span of all vectors $d\pi(Y_{\beta_1}) \cdots d\pi(Y_{\beta_k})v$, $k \geq 0$, $\beta_j > 0$ ("words of length k"). U is invariant under $\mathfrak{t}_\mathbb{C}$ (it is spanned by weight vectors) and under all $d\pi(Y_\beta)$. We show by induction on k that $d\pi(X_\beta)u \in U$ for all $\beta > 0$ and all words u of length k. For $k = 0$ this was shown. For $u = d\pi(Y_{\beta_1})u'$ with u' of length $k-1$,

$$d\pi(X_\beta)u = d\pi(Y_{\beta_1})d\pi(X_\beta)u' + d\pi([X_\beta, Y_{\beta_1}])u'.$$

The first term lies in U by induction. Since $\dim \mathfrak{g}_\gamma = 1$, $[X_\beta, Y_{\beta_1}] \in \mathfrak{g}_{\beta-\beta_1}$ is a multiple of $X_{\beta-\beta_1}$ if $\beta - \beta_1 > 0$ (induction applies), of $Y_{\beta_1-\beta}$ if $\beta - \beta_1 < 0$ (in U by definition), lies in $\mathfrak{t}_\mathbb{C}$ if $\beta = \beta_1$, and is 0 otherwise. So U is invariant under $\mathfrak{g}_\mathbb{C}$, hence under G (Proposition 5.3.3), and $U = V$. The weights of the spanning vectors are $\lambda - \beta_1 - \cdots - \beta_k \preceq \lambda$. □

Theorem 10.0.2: Weyl dimension formula

The irreducible representation with highest weight λ has dimension $d_\lambda = \prod_{\alpha>0} \dfrac{(\lambda+\rho,\alpha)}{(\rho,\alpha)}$.

Proof. Recall $(w\mu)(H_\nu) = (w\mu,\nu) = (\mu, w^{-1}\nu)$ (W acts orthogonally) and $\varepsilon(w^{-1}) = \varepsilon(w)$. Hence, using Proposition 9.2.3 and $e^{\alpha/2}(H) - e^{-\alpha/2}(H) = 2i\sin(\alpha(H)/2)$,

$$A_{\lambda+\rho}(sH_\rho) = \sum_w \varepsilon(w) e^{is(\lambda+\rho, w^{-1}\rho)} = A_\rho(sH_{\lambda+\rho}) = \prod_{\alpha>0} 2i\sin\tfrac{s}{2}(\alpha, \lambda+\rho)\big),$$

and likewise $A_\rho(sH_\rho) = \prod_{\alpha>0} 2i\sin(\frac{s}{2}(\alpha,\rho))$, which is nonzero for small $s \neq 0$ as $(\alpha,\rho) > 0$. By Theorem 10.0.1(i) and continuity of χ_λ,

$$d_\lambda = \chi_\lambda(e) = \lim_{s\to 0} \frac{A_{\lambda+\rho}(sH_\rho)}{A_\rho(sH_\rho)} = \prod_{\alpha>0} \frac{(\alpha,\lambda+\rho)}{(\alpha,\rho)}.$$ □

Example 10.0.4 (SU(2) revisited): For SU(2), $\Delta^+ = \{\alpha\}$ with $\alpha = 2\varepsilon$, $\rho = \varepsilon$, $P(T) = \mathbb{Z}\varepsilon$, $P_+ = \mathbb{N}_0\varepsilon$, $W = \{1, s_\alpha\}$ with $s_\alpha\varepsilon = -\varepsilon$. With $x = e^{i\theta}$, the character formula reads $\chi_{n\varepsilon}(t_\theta) = \frac{x^{n+1} - x^{-n-1}}{x - x^{-1}}$ and $d_{n\varepsilon} = n+1$, in accordance with Theorem 3.3.1.

11 Representations of the classical groups

11.1 Root data of the classical groups

Lemma 11.1.1: $\mathrm{Sp}(n)$ is connected, and every element of $\mathrm{Sp}(n)$ is conjugate in $\mathrm{Sp}(n)$ to an element of the torus T of Example 7.2.4(iv).

Proof. Let $\beta(x,y) := x^T J y$, an antisymmetric bilinear form on $\mathbb{C}^{2n}$, and $j(x) := J\bar{x}$, an antilinear map. Since J is real with $J^T = -J = J^{-1}$, we have $\langle x, j(y)\rangle = (J\bar{y})^* x = y^T J^T x = -\beta(y,x) = \beta(x,y)$, $\langle j(x), j(y)\rangle = \langle y, x\rangle$ and $j^2 = -1$. In particular $\langle j(x), x\rangle = \overline{\beta(x,x)} = 0$.

Let $g \in \mathrm{Sp}(n)$. From $g^T J g = J$ and $(g^T)^{-1} = \bar{g}$ (unitarity) we get $Jg = \bar{g}J$, hence $gJ = J\bar{g}$ and $g\,j(x) = j(gx)$. Let v be a unit eigenvector, $gv = \lambda v$; then $g\,j(v) = j(\lambda v) = \bar{\lambda}\,j(v)$, and $v \perp j(v)$. The subspace $E := \mathrm{span}\{v, j(v)\}$ is invariant under g and j, and so

is $E^\perp$ (for g by unitarity; for j because for $x \in E^\perp$, $\langle j(x), j(v)\rangle = \langle v, x\rangle = 0$ and $\langle j(x), v\rangle = -\langle j(x), j(j(v))\rangle = -\langle j(v), x\rangle = 0$). By induction we obtain an orthonormal basis $v_1, \dots, v_n, j(v_1), \dots, j(v_n)$ of eigenvectors of g, with $gv_a = \lambda_a v_a$. Let k be the unitary matrix with columns $v_1, \dots, v_n, -j(v_1), \dots, -j(v_n)$. Its Gram matrix for β is J: $\beta(v_a, v_b) = \langle v_a, j(v_b)\rangle = 0$, $\beta(j(v_a), j(v_b)) = \langle j(v_a), j^2(v_b)\rangle = -\langle j(v_a), v_b\rangle = 0$, and $\beta(v_a, -j(v_b)) = -\langle v_a, j^2(v_b)\rangle = \langle v_a, v_b\rangle = \delta_{ab}$. Hence $k^T J k = J$, $k \in \mathrm{Sp}(n)$, and $k^{-1}gk = \mathrm{diag}(\lambda_1, \dots, \lambda_n, \bar\lambda_1, \dots, \bar\lambda_n) \in T$. As T is connected, $\mathrm{Sp}(n) = \bigcup_k kTk^{-1}$ is connected. □

We equip the Lie algebras with the invariant inner products $\langle X, Y\rangle = -\operatorname{tr}(XY)$ for $\mathrm{U}(n)$ and $-\frac{1}{2}\operatorname{tr}(XY)$ for $\mathrm{SO}(m)$ and $\mathrm{Sp}(n)$. With the maximal tori of Example 7.2.4, $\mathfrak{t} \cong \mathbb{R}^n$ via the angles $(\theta_1, \dots, \theta_n)$, and the functionals $e_j(\theta) := \theta_j$ form an orthonormal basis of $\mathfrak{t}^*$.

Theorem 11.1.1: Root data

For the classical groups and their maximal tori from Example 7.2.4, $P(T) = \mathbb{Z}^n$ (with respect to $e_1, \dots, e_n$), and the roots, positive roots (for a suitable H_0), simple roots, Weyl groups and Weyl vectors are as follows:

G	Δ^+	simple roots	W, $\lvert W\rvert$	ρ
$\mathrm{U}(n)$	$e_j - e_k$, $j < k$	$e_j - e_{j+1}$	S_n, $n!$	$\sum_j \frac{n+1-2j}{2} e_j$
$\mathrm{SO}(2n+1)$	$e_j \pm e_k$ $(j < k)$, e_j	$e_j - e_{j+1}$, e_n	$S_n \ltimes (\mathbb{Z}/2)^n$, $2^n n!$	$\sum_j (n - j + \frac{1}{2}) e_j$
$\mathrm{Sp}(n)$	$e_j \pm e_k$ $(j < k)$, $2e_j$	$e_j - e_{j+1}$, $2e_n$	$S_n \ltimes (\mathbb{Z}/2)^n$, $2^n n!$	$\sum_j (n - j + 1) e_j$
$\mathrm{SO}(2n)$	$e_j \pm e_k$ $(j < k)$	$e_j - e_{j+1}$, $e_{n-1} + e_n$	$S_n \ltimes (\mathbb{Z}/2)^{n-1}$, $2^{n-1} n!$	$\sum_j (n - j) e_j$

Here $\Delta = \Delta^+ \cup -\Delta^+$, S_n permutes the e_j, $(\mathbb{Z}/2)^n$ acts by sign changes $e_j \mapsto -e_j$, and for $\mathrm{SO}(2n)$ only even numbers of sign changes occur ($n \geq 2$ for $\mathrm{SO}(2n)$). Consequently, P_+ consists of the $\lambda \in \mathbb{Z}^n$ with

$$\begin{array}{ll} \mathrm{U}(n): & \lambda_1 \geq \cdots \geq \lambda_n; \\ \mathrm{SO}(2n+1),\ \mathrm{Sp}(n): & \lambda_1 \geq \cdots \geq \lambda_n \geq 0; \\ \mathrm{SO}(2n): & \lambda_1 \geq \cdots \geq \lambda_{n-1} \geq |\lambda_n|. \end{array}$$

Proof. Weight lattice. In all cases $e^H = 1$ for $H \leftrightarrow \theta$ iff all $\theta_j \in 2\pi\mathbb{Z}$, so $\Lambda = 2\pi\mathbb{Z}^n$ and $P(T) = \mathbb{Z}^n$.

Roots. $\mathrm{U}(n)$: Example 8.2.3. $\mathrm{SO}(m)$: on $\mathbb{C}^m$, the block $R(\theta_j)$ has eigenvectors $e_{2j-1} \mp i e_{2j}$ with eigenvalues $e^{\pm i\theta_j}$, so the weights of $\mathbb{C}^m$ are $\pm e_j$, together with 0 if m is odd, each with multiplicity one. The map $\Lambda^2\mathbb{C}^m \to \mathfrak{so}(m)_{\mathbb{C}} = \{X \in M_m(\mathbb{C}) \mid X^T = -X\}$, $u \wedge v \mapsto uv^T - vu^T$, is a linear isomorphism, and it is $\mathrm{SO}(m)$-equivariant since $g(uv^T - vu^T)g^{-1} = (gu)(gv)^T - (gv)(gu)^T$ for $g^{-1} = g^T$. The weights of $\Lambda^2\mathbb{C}^m$ are the sums of two weights of $\mathbb{C}^m$ belonging to distinct basis vectors. The nonzero ones are $\pm e_j \pm e_k$ $(j < k)$, and for odd m also $\pm e_j$. $\mathrm{Sp}(n)$: the weights of $\mathbb{C}^{2n}$ are $\pm e_j$. Let $\mathfrak{sp}(2n, \mathbb{C}) := \{X \in M_{2n}(\mathbb{C}) \mid X^T J + JX = 0\}$. By the block computation in Example 5.2.3, its complex dimension is $n^2 + n(n+1)$, the real dimension of $\mathfrak{sp}(n) \subset \mathfrak{sp}(2n, \mathbb{C})$; as $\mathfrak{sp}(n) \cap i\mathfrak{sp}(n) = 0$ (skew-Hermitian versus Hermitian), $\mathfrak{sp}(2n, \mathbb{C}) = \mathfrak{sp}(n)_{\mathbb{C}}$. The map $X \mapsto JX$ is a linear isomorphism of $\mathfrak{sp}(2n, \mathbb{C})$ onto the symmetric matrices, and $J(tXt^{-1}) = (t^{-1})^T (JX) t^{-1}$ for $t \in \mathrm{Sp}(n)$ (using $Jt = (t^T)^{-1} J$). So $\mathfrak{sp}(n)_{\mathbb{C}}$ is the space of symmetric bilinear forms on $\mathbb{C}^{2n}$, i.e., Sym^2 of the dual of $\mathbb{C}^{2n}$, whose weights are $\pm e_j \pm e_k$ ($j < k$, from distinct basis vectors), $\pm 2e_j$ and 0.

Positive and simple roots. Take $H_0 \leftrightarrow (n, n-1, \dots, 1)+$ a small generic perturbation; then

Δ^+ is as stated. Let S be the proposed set of simple roots. It is linearly independent, has $\dim \operatorname{span} \Delta$ elements, and every positive root is a nonnegative integer combination of S:

$$e_j - e_k = \sum_{l=j}^{k-1} (e_l - e_{l+1}), \quad e_j = (e_j - e_n) + e_n, \quad 2e_j = 2(e_j - e_n) + 2e_n,$$

$$e_j + e_k = (e_j - e_n) + (e_k - e_n) + 2e_n$$

in types B_n ($\mathrm{SO}(2n+1)$, write $2e_n = e_n + e_n$) and C_n ($\mathrm{Sp}(n)$), and $e_j + e_k = (e_j - e_{n-1}) + (e_k - e_n) + (e_{n-1} + e_n)$ for $j < k \leq n-1$, $e_j + e_n = (e_j - e_{n-1}) + (e_{n-1} + e_n)$ in type D_n ($\mathrm{SO}(2n)$). Then S is the set of simple roots: each $\alpha \in S$ is simple, since writing $\alpha = \beta + \gamma$ with $\beta, \gamma > 0$ and expanding β, γ in S with nonnegative coefficients would, by linear independence, force $\beta = 0$ or $\gamma = 0$; and the simple roots form a basis of $\operatorname{span} \Delta$ (Lemma 8.5.1(iv)), so they cannot be more than $|S|$.

Weyl groups. By Theorem 9.1.1(ii), W is generated by the reflections s_α. Here $s_{e_j - e_k}$ swaps e_j and e_k; s_{e_j} and s_{2e_j} change the sign of e_j; $s_{e_j + e_k}$ maps $e_j \mapsto -e_k$, $e_k \mapsto -e_j$. In types B_n, C_n these generate all signed permutations. In type D_n, every generator changes an even number of signs, and $s_{e_j - e_k} s_{e_j + e_k}$ changes the signs of e_j and e_k; together with the transpositions these generate all signed permutations with an even number of sign changes.

Weyl vectors and dominant weights. ρ is obtained by summing: e.g., for B_n, $\sum_{j<k} \big((e_j - e_k) + (e_j + e_k)\big) + \sum_j e_j = \sum_j (2(n-j)+1) e_j$. Dominance means $(\lambda, \alpha) \geq 0$ for the simple roots: $\lambda_j \geq \lambda_{j+1}$, and additionally $\lambda_n \geq 0$ (types B, C), resp. $\lambda_{n-1} + \lambda_n \geq 0$ (type D), which together with $\lambda_{n-1} \geq \lambda_n$ means $\lambda_{n-1} \geq |\lambda_n|$. □

11.2 The unitary groups

For $\mathrm{U}(n)$ let $\delta := (n-1, n-2, \dots, 0) = \rho + \frac{n-1}{2}(1, \dots, 1)$. Since $(1, \dots, 1)$ is W-invariant, $A_{\mu+\delta} = e^{\frac{n-1}{2}(1,\dots,1)} A_{\mu+\rho}$ for every μ, so $A_{\lambda+\rho}/A_\rho = A_{\lambda+\delta}/A_\delta$. With $x_j := e^{i\theta_j}$ and $\varepsilon(w) = \operatorname{sgn}(w)$ for $w \in S_n$,

$$A_{\lambda+\delta} = \sum_{\sigma \in S_n} \operatorname{sgn}(\sigma) \prod_j x_{\sigma(j)}^{\lambda_j + n - j} = \det\big(x_k^{\lambda_j + n - j}\big)_{j,k}, \qquad A_\delta = \det\big(x_k^{n-j}\big)_{j,k} = \prod_{j<k} (x_j - x_k)$$

(Vandermonde).

Theorem 11.2.1: Irreducible representations of U(n)

The irreducible representations π_λ of $\mathrm{U}(n)$ are labelled by $\lambda \in \mathbb{Z}^n$ with $\lambda_1 \geq \cdots \geq \lambda_n$. The character at a matrix with eigenvalues $x_1, \dots, x_n$ is the *Schur function*

$$s_\lambda(x_1, \dots, x_n) = \frac{\det\big(x_k^{\lambda_j + n - j}\big)_{j,k}}{\prod_{j<k}(x_j - x_k)}$$

(for distinct x_j; by continuity otherwise), and $d_\lambda = \prod_{j<k} \frac{\lambda_j - \lambda_k + k - j}{k - j}$.

Proof. Theorems 10.0.1, 10.0.2, 11.1.1 and the formulas above; regularity of $\operatorname{diag}(x_j)$ means $x_j \neq x_k$ for $j \neq k$, and $(\lambda + \rho, e_j - e_k) = \lambda_j - \lambda_k + k - j$, $(\rho, e_j - e_k) = k - j$. □

Example 11.2.1: (i) $\Lambda^k \mathbb{C}^n \cong \pi_{(1^k, 0^{n-k})}$, with character the elementary symmetric polynomial $e_k(x)$.

(ii) $\mathrm{Sym}^k \mathbb{C}^n \cong \pi_{(k,0,\dots,0)}$, with character the complete symmetric polynomial $h_k(x)$.

(iii) $\det^m = \pi_{(m,\dots,m)}$, and $\pi_\lambda \otimes \det^m \cong \pi_{\lambda+(m,\dots,m)}$.

(iv) $\pi_\lambda^* \cong \pi_{\lambda^*}$ with $\lambda^* = (-\lambda_n, \dots, -\lambda_1)$.

Proof. (i) The weights of $\Lambda^k\mathbb{C}^n$ are $e_{j_1}+\cdots+e_{j_k}$, $j_1 < \cdots < j_k$, each with multiplicity one, forming one S_n-orbit, the orbit of $(1^k, 0^{n-k})$. The irreducible summand containing the weight vector for $(1^k, 0^{n-k})$ has all weights of that orbit (Proposition 8.1.2(iii)), so it has dimension $\geq \binom{n}{k}$ and is everything. Its highest weight is the dominant element $(1^k, 0^{n-k})$ of the orbit (Theorem 9.1.1(iii)). (ii) $e_1^k \in \mathrm{Sym}^k\mathbb{C}^n$ has weight $(k, 0, \dots, 0)$, which is the highest weight of Sym^k (the weights are $\mu \in \mathbb{N}_0^n$ with $|\mu| = k$, and $(k, 0, \dots, 0)$ maximizes $\mu(H_0)$). So Sym^k contains $\pi_{(k,0,\dots,0)}$ (the summand containing the one-dimensional top weight space). By Theorem 11.2.1, its dimension is $\prod_{l=2}^n \frac{k+l-1}{l-1} = \binom{n+k-1}{k} = \dim \mathrm{Sym}^k\mathbb{C}^n$; so they are equal. (iii) det is one-dimensional with weight $(1, \dots, 1)$, and tensoring with it shifts all weights. (iv) The weights of π^* are the negatives of those of π (Lemma 3.2.2(iv)). The lowest weight of π_λ is $w_0\lambda = (\lambda_n, \dots, \lambda_1)$, where w_0 reverses the order: $w_0\lambda$ is a weight, and for any weight μ, $w_0\mu$ is a weight, so $\lambda - w_0\mu \in Q_+$ (Proposition 10.0.3); as w_0 maps Δ^+ to $-\Delta^+$, $w_0\lambda - \mu \in -Q_+$, i.e., $\mu(H_0) \geq (w_0\lambda)(H_0)$. Hence the highest weight of π^* is $-w_0\lambda = \lambda^*$. □

Remark 11.2.2 (Young diagrams): Highest weights with $\lambda_n \geq 0$ are partitions with at most n parts; one depicts λ by a *Young diagram* with λ_j boxes in row j. Its *columns* have lengths $\lambda'_1 \geq \lambda'_2 \geq \cdots$, where $\lambda'_i := |\{j \mid \lambda_j \geq i\}|$.

Lemma 11.2.3: Let λ be a partition with at most n parts and column lengths $\lambda'_1, \dots, \lambda'_m$. Then π_λ is a subrepresentation of $\Lambda^{\lambda'_1}\mathbb{C}^n \otimes \cdots \otimes \Lambda^{\lambda'_m}\mathbb{C}^n \subset (\mathbb{C}^n)^{\otimes|\lambda|}$.

Proof. The vector $v := \bigotimes_i (e_1 \wedge \cdots \wedge e_{\lambda'_i})$ has weight $\sum_i (1^{\lambda'_i}, 0, \dots) = \lambda$. In each factor, $e_1 \wedge \cdots \wedge e_{\lambda'_i}$ spans the weight space with maximal value at H_0; hence λ is the weight of the tensor product with maximal value at H_0, and its weight space is $\mathbb{C}v$. The irreducible summand containing v then has highest weight λ. Finally, $\Lambda^c\mathbb{C}^n$ embeds into $(\mathbb{C}^n)^{\otimes c}$ as antisymmetric tensors. □

Theorem 11.2.2: Schur–Weyl duality

Let $\mathrm{U}(n)$ act on $V := (\mathbb{C}^n)^{\otimes k}$ by $g \mapsto g^{\otimes k}$ and S_k by permuting the factors. These actions commute, and

$$V \cong \bigoplus_\lambda \pi_\lambda \otimes M_\lambda \qquad \text{as a representation of } \mathrm{U}(n) \times S_k,$$

where λ runs over the partitions of k with at most n parts, $M_\lambda := \mathrm{Hom}_{\mathrm{U}(n)}(\pi_\lambda, V)$ with S_k acting by composition, and the M_λ are irreducible and pairwise inequivalent representations of S_k.

Proof. Step 1: The commutant of S_k in $\mathrm{End}(V)$ *is* $\mathcal{A} := \mathrm{span}\{g^{\otimes k} \mid g \in \mathrm{U}(n)\}$. Clearly $\mathcal{A}$ commutes with S_k. Conversely, $\mathrm{End}(V) = \mathrm{End}(\mathbb{C}^n)^{\otimes k}$, and the commutant of S_k is the space of symmetric tensors. It is spanned by the $X^{\otimes k}$, $X \in \mathrm{End}(\mathbb{C}^n)$: the coefficient of $s_1 \cdots s_k$ in the polynomial $(s_1X_1 + \cdots + s_kX_k)^{\otimes k}$ is $\sum_{\sigma\in S_k} X_{\sigma(1)} \otimes \cdots \otimes X_{\sigma(k)}$, and coefficients of a polynomial with values in a subspace lie in that subspace. Let ℓ be a linear functional vanishing on $\mathcal{A}$. Then $Z \mapsto \ell((e^Z)^{\otimes k})$ is holomorphic on $M_n(\mathbb{C}) = \mathfrak{u}(n) \oplus i\mathfrak{u}(n) \cong \mathbb{C}^{n^2}$ and vanishes on $\mathfrak{u}(n) \cong \mathbb{R}^{n^2}$, hence vanishes identically. Since $\exp : M_n(\mathbb{C}) \to \mathrm{GL}(n, \mathbb{C})$ is surjective (by the Jordan normal form it suffices to treat a block $\mu(1 + N)$ with N nilpotent; choose c with $e^c = \mu$ and put $L := \sum_{k\geq 1}(-1)^{k+1}N^k/k$, a finite sum; then $e^{c+L} = \mu e^L = \mu(1 + N)$ by the formal identity $\exp(\log(1 + x)) = 1 + x$) and $\mathrm{GL}(n, \mathbb{C})$ is dense in $M_n(\mathbb{C})$, $\ell(X^{\otimes k}) = 0$ for all X. Hence $\mathcal{A}$ contains all symmetric tensors.

Step 2: decomposition. The map $\bigoplus_\lambda \pi_\lambda \otimes M_\lambda \to V$, $u \otimes A \mapsto Au$, is an isomorphism (isotypic decomposition, Proposition 2.2.3) intertwining $\mathrm{U}(n) \times S_k$. Every $g^{\otimes k}$ acts on $\pi_\lambda \otimes M_\lambda$ as $\pi_\lambda(g) \otimes 1$, hence so does every element of $\mathcal{A}$ (as $T_\lambda \otimes 1$ for some T_λ).

Irreducibility. If $M_\lambda = M' \oplus M''$ with S_k-invariant $M' \neq 0, M_\lambda$, let $p : M_\lambda \to M'$ be an S_k-equivariant projection (complete reducibility for the finite group S_k). The operator P equal to $1 \otimes p$ on $\pi_\lambda \otimes M_\lambda$ and 0 on the other summands commutes with S_k, so $P \in \mathcal{A}$ and $1 \otimes p = T_\lambda \otimes 1$, which forces p to be scalar, a contradiction.

Inequivalence. If $\phi : M_\lambda \to M_\mu$ is an S_k-isomorphism with $\lambda \neq \mu$, choose any linear $\psi \neq 0$ from π_λ to π_μ and let Q be $\psi \otimes \phi$ on $\pi_\lambda \otimes M_\lambda$ (with values in $\pi_\mu \otimes M_\mu$) and 0 elsewhere. Q commutes with S_k, so $Q \in \mathcal{A}$; but elements of $\mathcal{A}$ preserve each summand, while $Q \neq 0$ maps the λ-summand into the μ-summand. Contradiction.

Which λ occur. The weights of V are $e_{j_1} + \cdots + e_{j_k}$, i.e., $\mu \in \mathbb{N}_0^n$ with $\sum \mu_j = k$, so highest weights are partitions of k with at most n parts. Conversely each such λ occurs by Lemma 11.2.3. □

Remark 11.2.4: In the representation theory of symmetric groups, M_λ is identified with the Specht module $\mathcal{S}^\lambda$; we do not use this. For $k = 2$: $\mathbb{C}^n \otimes \mathbb{C}^n = \mathrm{Sym}^2\mathbb{C}^n \oplus \Lambda^2\mathbb{C}^n$, with S_2 acting trivially, resp. by the sign.

11.3 The special unitary groups

Proposition 11.3.1: The restriction of π_λ from $\mathrm{U}(n)$ to $\mathrm{SU}(n)$ is irreducible; $\pi_\lambda|_{\mathrm{SU}(n)} \cong \pi_\mu|_{\mathrm{SU}(n)}$ iff $\lambda - \mu \in \mathbb{Z}(1, \ldots, 1)$; and every irreducible representation of $\mathrm{SU}(n)$ arises in this way. Hence $\widehat{\mathrm{SU}(n)}$ is labelled by $\lambda_1 \geq \cdots \geq \lambda_{n-1} \geq \lambda_n = 0$, equivalently by the *Dynkin labels* $a_j := \lambda_j - \lambda_{j+1} \in \mathbb{N}_0$, $j = 1, \ldots, n-1$.

Proof. Every $g \in \mathrm{U}(n)$ is $g = c \cdot g'$ with $c \in \mathbb{T}$ (an n-th root of $\det g$) and $g' \in \mathrm{SU}(n)$. Scalars act by scalars on π_λ (Schur), so $\mathrm{SU}(n)$-invariant subspaces are $\mathrm{U}(n)$-invariant, and the restriction is irreducible. The maximal torus of $\mathrm{SU}(n)$ is $T' := T \cap \mathrm{SU}(n)$, with $\mathfrak{t}' = \{\theta \mid \sum \theta_j = 0\}$; restriction $\mathfrak{t}^* \to \mathfrak{t}'^*$ has kernel $\mathbb{R}(1, \ldots, 1)$, maps roots to roots and $P(T) = \mathbb{Z}^n$ onto $P(T')$ (a character of $T' \cong \mathbb{T}^{n-1}$, $(z_1, \ldots, z_{n-1}) \mapsto \mathrm{diag}(z_1, \ldots, z_{n-1}, (z_1 \cdots z_{n-1})^{-1})$, is $\prod_{j<n} z_j^{m_j}$, the restriction of $\prod_{j<n} x_j^{m_j}$). The highest weight of $\pi_\lambda|_{\mathrm{SU}(n)}$ is the restriction of λ; restrictions of λ, μ agree iff $\lambda - \mu \in \mathbb{R}(1, \ldots, 1) \cap \mathbb{Z}^n = \mathbb{Z}(1, \ldots, 1)$. Every dominant weight of $\mathrm{SU}(n)$ is the restriction of a dominant $\lambda \in \mathbb{Z}^n$ (choose a preimage and use that dominance only involves the differences $\lambda_j - \lambda_{j+1}$). Now apply Theorem 10.0.1 to $\mathrm{SU}(n)$. □

Example 11.3.2 (SU(3)): For Dynkin labels (a, b), i.e., $\lambda = (a+b, b, 0)$, Theorem 11.2.1 gives

$$d_{(a,b)} = \frac{(a+1)(b+1)(a+b+2)}{2}.$$

In physics notation: $\mathbf{1} = (0,0)$, $\mathbf{3} = (1,0) = \mathbb{C}^3$, $\bar{\mathbf{3}} = (0,1) = \Lambda^2\mathbb{C}^3 \cong (\mathbb{C}^3)^*$, $\mathbf{6} = (2,0) = \mathrm{Sym}^2\mathbb{C}^3$, $\mathbf{8} = (1,1)$, $\mathbf{10} = (3,0) = \mathrm{Sym}^3\mathbb{C}^3$. Then

$$\mathbf{3} \otimes \mathbf{3} = \mathbf{6} \oplus \bar{\mathbf{3}}, \qquad \mathbf{3} \otimes \bar{\mathbf{3}} = \mathbf{8} \oplus \mathbf{1}, \qquad \mathbf{3} \otimes \mathbf{3} \otimes \mathbf{3} = \mathbf{10} \oplus \mathbf{8} \oplus \mathbf{8} \oplus \mathbf{1},$$

and $\mathbf{8}$ is the complexified adjoint representation.

Proof. The identifications of $\mathbf{3}, \mathbf{6}, \mathbf{10}$ and $\Lambda^2\mathbb{C}^3 = \pi_{(1,1,0)}$, of Dynkin labels $(0,1)$, follow from Example 11.2.1 and Proposition 11.3.1. Also $(\mathbb{C}^3)^* \cong \pi_{(0,0,-1)}$ by Example 11.2.1(iv), and $(0,0,-1) + (1,1,1) = (1,1,0)$, so $(\mathbb{C}^3)^* \cong \Lambda^2\mathbb{C}^3$ as $\mathrm{SU}(3)$-representations.

First, $\mathbb{C}^3 \otimes \mathbb{C}^3 = \mathrm{Sym}^2\mathbb{C}^3 \oplus \Lambda^2\mathbb{C}^3 = \mathbf{6} \oplus \bar{\mathbf{3}}$. Next, $\mathbf{3} \otimes \bar{\mathbf{3}} \cong \mathbb{C}^3 \otimes (\mathbb{C}^3)^* \cong \mathrm{End}(\mathbb{C}^3) = \mathfrak{sl}(3, \mathbb{C}) \oplus \mathbb{C} \cdot 1$, with $\mathrm{SU}(3)$ acting by conjugation. The weights of $\mathfrak{sl}(3, \mathbb{C})$ are 0 and the roots $e_j - e_k$, and $e_1 - e_3$

(weight vector E_{13}) has maximal value at H_0; so $\mathfrak{sl}(3,\mathbb{C})$ contains $\pi_{(1,0,-1)} = \pi_{(2,1,0)}|_{\mathrm{SU}(3)}$, of Dynkin labels $(1,1)$ and dimension $8 = \dim \mathfrak{sl}(3,\mathbb{C})$. Hence $\mathfrak{sl}(3,\mathbb{C}) \cong \mathbf{8}$ and $\mathbf{3}\otimes\bar{\mathbf{3}} = \mathbf{8}\oplus\mathbf{1}$.

Finally, $\mathbf{3}^{\otimes 3} = (\mathbf{6}\otimes\mathbf{3})\oplus(\bar{\mathbf{3}}\otimes\mathbf{3}) = (\mathbf{6}\otimes\mathbf{3})\oplus\mathbf{8}\oplus\mathbf{1}$. The multiplication map $\mathrm{Sym}^2\mathbb{C}^3\otimes\mathbb{C}^3 \to \mathrm{Sym}^3\mathbb{C}^3$ is equivariant and surjective; by complete reducibility $\mathbf{6}\otimes\mathbf{3} \cong \mathbf{10}\oplus K$ with K its kernel, $\dim K = 8$. We compare weight multiplicities (for U(3)). In $\mathrm{Sym}^2\mathbb{C}^3\otimes\mathbb{C}^3$, the weight $(3,0,0)$ and its permutations have multiplicity 1, the permutations of $(2,1,0)$ have multiplicity 2 (e.g. $e_1^2\otimes e_2$ and $e_1e_2\otimes e_1$), and $(1,1,1)$ has multiplicity 3; in $\mathrm{Sym}^3\mathbb{C}^3$ all these have multiplicity 1. So the weights of K are the six permutations of $(2,1,0)$ (multiplicity 1) and $(1,1,1)$ (multiplicity 2), and the highest weight of K is $(2,1,0)$. Hence $K \supset \pi_{(2,1,0)} = \mathbf{8}$, and $K \cong \mathbf{8}$ by comparing dimensions. □

11.4 The orthogonal groups and spherical harmonics

Theorem 11.4.1: Irreducible representations of SO(m)

The irreducible representations of $\mathrm{SO}(2n+1)$ are labelled by $\lambda \in \mathbb{Z}^n$ with $\lambda_1 \geq \cdots \geq \lambda_n \geq 0$, those of $\mathrm{SO}(2n)$ $(n \geq 2)$ by $\lambda \in \mathbb{Z}^n$ with $\lambda_1 \geq \cdots \geq \lambda_{n-1} \geq |\lambda_n|$; characters and dimensions are given by Theorems 10.0.1 and 10.0.2 with the data of Theorem 11.1.1.

Proof. $\mathrm{SO}(m)$ is connected (Lemma 5.4.1); apply Theorems 10.0.1 and 11.1.1. □

Remark 11.4.1: For SO(3) $(n=1)$: $\Delta^+ = \{e_1\}$, $\rho = \frac{1}{2}e_1$, and $\lambda = \ell e_1$ gives $d = (\ell+\frac{1}{2})/\frac{1}{2} = 2\ell+1$, as in Proposition 5.4.2. Dominant "weights" with $\lambda_j \in \frac{1}{2}+\mathbb{Z}$ satisfy the inequalities but are not in $P(T) = \mathbb{Z}^n$; for $n=1$ they are the highest weights of the representations of SU(2) not factoring through SO(3).

Theorem 11.4.2: Spherical harmonics

Let $\mathcal{P}_\ell$ be the space of complex homogeneous polynomials of degree ℓ on $\mathbb{R}^3$ and $\mathcal{H}_\ell := \{P \in \mathcal{P}_\ell \mid \Delta P = 0\}$, with SO(3) acting by $(gP)(x) := P(g^{-1}x)$. Then $\mathcal{H}_\ell \cong D^{(\ell)}$ is irreducible of dimension $2\ell+1$, and

$$L^2(S^2) = \widehat{\bigoplus_{\ell\in\mathbb{N}_0}} \mathcal{H}_\ell|_{S^2}$$

(orthogonal Hilbert sum), each irreducible representation of SO(3) occurring exactly once.

Proof. Dimension. Define the inner product on $\mathcal{P} := \bigoplus_\ell \mathcal{P}_\ell$ by $\langle x^a, x^b\rangle := a!\,\delta_{ab}$ for multi-indices a,b. Then $\langle x_j P, Q\rangle = \langle P, \partial_j Q\rangle$ (check on monomials), so multiplication by $|x|^2$ is adjoint to Δ. Hence $\mathcal{P}_\ell = \mathcal{H}_\ell \oplus |x|^2\mathcal{P}_{\ell-2}$ (orthogonally), and as multiplication by $|x|^2$ is injective, $\dim\mathcal{H}_\ell = \binom{\ell+2}{2} - \binom{\ell}{2} = 2\ell+1$. Since Δ commutes with rotations, $\mathcal{H}_\ell$ is invariant.

Irreducibility. Let $T = \mathrm{SO}(2)$ be the rotations about the x_3-axis by angle θ. On $z := x_1 + ix_2$, $\bar z$, x_3, the element g^{-1} acts by $z \mapsto e^{-i\theta}z$, $\bar z \mapsto e^{i\theta}\bar z$, $x_3 \mapsto x_3$; so the weights of $\mathcal{P}_\ell$ lie in $\{-\ell,\dots,\ell\}$ (in units of e_1), and $\bar z^\ell = (x_1 - ix_2)^\ell$, which is harmonic, has weight ℓ. Hence $\mathcal{H}_\ell$ contains $D^{(\ell)}$ (the summand containing $\bar z^\ell$ has highest weight ℓ), and as $\dim\mathcal{H}_\ell = 2\ell+1$, $\mathcal{H}_\ell \cong D^{(\ell)}$.

Decomposition. SO(3) acts transitively on S^2, and the stabilizer of e_3 is T; so $g \mapsto ge_3$ identifies G/T with S^2, and the image of Haar measure is an SO(3)-invariant probability measure on S^2, which by the uniqueness argument of Lemma 9.4.1(iii) is the normalized surface measure. Thus $f \mapsto f(\cdot\, e_3)$ is a unitary SO(3)-isomorphism from $L^2(S^2)$ onto $L^2(G/T)$ in the sense of Proposition 4.3.2. By that proposition, $D^{(\ell)}$ occurs with multiplicity $\dim(D^{(\ell)})^T$, the multiplicity of the weight 0, which is 1. Restriction to S^2 is injective on $\mathcal{H}_\ell$ (by homogeneity), so $\mathcal{H}_\ell|_{S^2}$ is the (unique) $D^{(\ell)}$-isotypic component of $L^2(S^2)$. Isotypic components for inequivalent irreducibles are orthogonal (the orthogonal projection onto one restricted to the other is an intertwiner, hence 0 by Schur), and their sum is dense by Theorem 4.3.1. □

11.5 The compact symplectic groups

$\mathrm{Sp}(n)$ is connected (Lemma 11.1.1), so by Theorems 10.0.1 and 11.1.1 its irreducible representations are labelled by $\lambda \in \mathbb{Z}^n$ with $\lambda_1 \geq \cdots \geq \lambda_n \geq 0$. The defining representation $\mathbb{C}^{2n}$ (weights $\pm e_j$) has highest weight e_1, and the complexified adjoint representation, which is $\mathrm{Sym}^2((\mathbb{C}^{2n})^*)$ by the proof of Theorem 11.1.1, has highest weight $2e_1$. For $n = 1$, $\mathrm{Sp}(1) = \mathrm{SU}(2)$: for $g \in M_2(\mathbb{C})$ one has $g^T J g = (\det g)J$, so $\mathrm{Sp}(1) = \{g \in \mathrm{U}(2) \mid \det g = 1\}$.

12 Applications to physics

In this section, statements labelled *physical input* are taken from physics and are not proved here; all mathematical consequences drawn from them are proved. Dirac notation uses $\langle u|v\rangle := \langle v, u\rangle$, consistently with our first-argument-linear inner product.

12.1 Symmetries in quantum mechanics

Physical input. Pure states of a quantum system are rays in a Hilbert space $\mathcal{H}$, and a symmetry group G acts on $\mathcal{H}$ by unitary operators, possibly only up to phases (Wigner's theorem). For rotations, the operators can be chosen to form a unitary representation of $\mathrm{SU}(2)$ (rather than of $\mathrm{SO}(3)$). If the representation is finite-dimensional, Corollary 2.2.1 and Theorem 3.3.1 decompose it into the $D^{(j)}$, $j \in \frac{1}{2}\mathbb{N}_0$; the representations with half-integral j do not factor through $\mathrm{SO}(3)$ (Proposition 5.4.2) and describe *spin*.

12.2 Angular momentum

Let (π, V) be a finite-dimensional unitary representation of $\mathrm{SU}(2)$, $X_a = -\frac{i}{2}\sigma_a$ as in Proposition 5.4.2, and define the self-adjoint *angular momentum operators* $J_a := i\,d\pi(X_a)$. From $[X_1, X_2] = X_3$ (and cyclically) we get $[J_1, J_2] = iJ_3$ (and cyclically). Since $X_3 = -\frac{i}{2}H$ and $X_1 + iX_2 = -iE$ and $X_1 - iX_2 = -iF$, we have $d\pi(H) = 2J_3$, $d\pi(E) = J_+ := J_1 + iJ_2$, $d\pi(F) = J_- := J_1 - iJ_2$. Let $J^2 := J_1^2 + J_2^2 + J_3^2$.

Proposition 12.2.1: J^2 commutes with $\pi(g)$ for all g and acts on $D^{(j)}$ by the scalar $j(j+1)$. $D^{(j)}$ has an orthonormal basis $|j, m\rangle$, $m = -j, -j+1, \ldots, j$, with $J_3|j,m\rangle = m|j,m\rangle$ and $J_\pm|j,m\rangle = \sqrt{j(j+1) - m(m\pm 1)}\,|j, m\pm 1\rangle$.

Proof. $[J^2, J_a] = 0$ follows from the commutation relations (e.g. $[J_1^2, J_3] = J_1[J_1, J_3] + [J_1, J_3]J_1 = -i(J_1J_2 + J_2J_1)$ and $[J_2^2, J_3] = i(J_2J_1 + J_1J_2)$). So J^2 commutes with $d\pi(\mathfrak{g})$, hence with $\pi(G)$ (Proposition 5.3.3), and is a scalar on the irreducible $D^{(j)}$ by Schur. With $J_-J_+ = J_1^2 + J_2^2 + i[J_1, J_2] = J^2 - J_3^2 - J_3$, on the highest weight vector ($J_+v = 0$, $J_3v = jv$, as $Hv = 2jv$ by Theorem 5.4.1) we get $J^2v = (j^2 + j)v$. The weights of $D^{(j)} = \pi_{2j}$ under $H = 2J_3$ are $2j, 2j-2, \ldots, -2j$, each with multiplicity one; normalized eigenvectors $|j,m\rangle$ of J_3 are orthonormal (J_3 self-adjoint). Since $J_+^* = J_-$ and $J_+|j,m\rangle$ is an eigenvector of J_3 for $m+1$, $J_+|j,m\rangle = c_m|j,m+1\rangle$ with $|c_m|^2 = \langle J_-J_+|j,m\rangle, |j,m\rangle\rangle = j(j+1) - m^2 - m$; the phases can be normalized to make all $c_m \geq 0$, and then $J_-|j,m+1\rangle = J_+^*|j,m+1\rangle = c_m|j,m\rangle$. □

Corollary 3.3.3 decomposes $D^{(j_1)} \otimes D^{(j_2)}$ into $D^{(J)}$, $|j_1 - j_2| \leq J \leq j_1 + j_2$; the coefficients $\langle j_1 m_1\, j_2 m_2 | JM\rangle$ of $|J, M\rangle$ in the basis $|j_1, m_1\rangle \otimes |j_2, m_2\rangle$ (nonzero only for $m_1 + m_2 = M$, by comparing J_3-eigenvalues) are the *Clebsch–Gordan coefficients*. For example, $D^{(1/2)} \otimes D^{(1/2)} = D^{(0)} \oplus D^{(1)}$, with the singlet spanning $\Lambda^2\mathbb{C}^2$ and the triplet spanning $\mathrm{Sym}^2\mathbb{C}^2$.

Theorem 12.2.1: Wigner–Eckart

Let π be a unitary representation of SU(2) on a finite-dimensional space $\mathcal{H}$, and let T_q, $q = -k, \dots, k$, be operators on $\mathcal{H}$ such that $\pi(g) T_q \pi(g)^{-1} = \sum_{q'} D^{(k)}_{q'q}(g) T_{q'}$, where $D^{(k)}_{q'q}(g) := \langle k, q' | D^{(k)}(g) | k, q \rangle$. Let $W_1, W_2 \subset \mathcal{H}$ be irreducible invariant subspaces with $W_1 \cong D^{(j)}$, $W_2 \cong D^{(j')}$, and orthonormal bases $|j, m\rangle_1$, $|j', m'\rangle_2$ corresponding to the standard bases. Then there is a constant c (the reduced matrix element) with

$$ {}_1\langle j, m | T_q | j', m' \rangle_2 = c \, \langle j' m' \, kq | jm \rangle \qquad \text{for all } m, m', q. $$

In particular these matrix elements vanish unless $m = m' + q$ and $|j' - k| \le j \le j' + k$.

Proof. Let P be the orthogonal projection onto W_1, an intertwiner. Define $\Phi : D^{(k)} \otimes D^{(j')} \to D^{(j)}$ by $\Phi(|k, q\rangle \otimes |j', m'\rangle) := P T_q |j', m'\rangle_2$ (identifying W_1 with $D^{(j)}$). The transformation law of T_q shows that Φ is an intertwiner: $\Phi(D^{(k)}(g)|k, q\rangle \otimes \pi(g) v) = \sum_{q'} D^{(k)}_{q'q}(g) P T_{q'} \pi(g) v = P \pi(g) T_q v = \pi(g) P T_q v$. By Corollary 3.3.3 and Proposition 2.2.3, $D^{(j)}$ occurs at most once in $D^{(k)} \otimes D^{(j')}$, so $\mathrm{Hom}_G(D^{(k)} \otimes D^{(j')}, D^{(j)})$ has dimension ≤ 1, and if nonzero it is spanned by $\Phi_0 :=$ the orthogonal projection onto the $D^{(j)}$-component followed by the isometry onto $D^{(j)}$ given by the Clebsch–Gordan basis. Hence $\Phi = c\Phi_0$, and $\langle j, m | \Phi_0 (|k, q\rangle \otimes |j', m'\rangle) = \langle j' m' \, kq | jm \rangle$ (taking the factors in the order $D^{(j')} \otimes D^{(k)} \cong D^{(k)} \otimes D^{(j')}$). □

For electric dipole transitions ($k = 1$) this gives the *selection rules* $\Delta j \in \{0, \pm 1\}$ and $\Delta m \in \{0, \pm 1\}$.

12.3 The hydrogen atom and SO(4)

Physical input (Pauli 1926). For the hydrogen atom (atomic units), the angular momentum L and the Runge–Lenz vector A commute with the Hamiltonian H. On an eigenspace $\mathcal{E}$ of H with eigenvalue $E < 0$, assumed finite-dimensional, the operators L_a and $K_a := A_a / \sqrt{-2E}$ are self-adjoint and satisfy

$$ [L_a, L_b] = i\epsilon_{abc} L_c, \quad [L_a, K_b] = i\epsilon_{abc} K_c, \quad [K_a, K_b] = i\epsilon_{abc} L_c, $$

$$ L \cdot K = 0, \qquad E = -\frac{1}{2(L^2 + K^2 + 1)}. $$

Proposition 12.3.1: Under these assumptions, $E = -\frac{1}{2n^2}$ for some $n \in \mathbb{N}$, and $\mathcal{E} \cong (D^{(j)} \otimes D^{(j)})^{\oplus m}$ with $n = 2j + 1$ as a representation of $\mathfrak{su}(2) \oplus \mathfrak{su}(2)$. In particular $\dim \mathcal{E}$ is a multiple of n^2, and under the rotations L, $\mathcal{E} \cong (D^{(0)} \oplus D^{(1)} \oplus \cdots \oplus D^{(n-1)})^{\oplus m}$.

Proof. $J^{\pm} := \frac{1}{2}(L \pm K)$ satisfy $[J^{\pm}_a, J^{\pm}_b] = i\epsilon_{abc} J^{\pm}_c$ and $[J^{+}_a, J^{-}_b] = 0$ (direct computation from the relations). So $\mathfrak{su}(2) \oplus \mathfrak{su}(2)$ acts on $\mathcal{E}$ by $X_a \mapsto -iJ^{+}_a$ on the first and $-iJ^{-}_a$ on the second summand, by skew-Hermitian operators. By Theorem 5.4.1(ii) applied to the first summand, $\mathcal{E} = \bigoplus_j \mathcal{E}_j$ with $\mathcal{E}_j$ the $D^{(j)}$-isotypic component for J^{+}; as J^{-} commutes with J^{+}, each $\mathcal{E}_j$ is J^{-}-invariant, and $\mathcal{E}_j \cong D^{(j)} \otimes M_j$ with J^{-} acting on $M_j := \mathrm{Hom}_{J^{+}}(D^{(j)}, \mathcal{E})$; decomposing M_j under J^{-} gives $\mathcal{E} \cong \bigoplus D^{(j)} \otimes D^{(j')}$. Now $L \cdot K = 0$ gives $(J^{+})^2 - (J^{-})^2 = L \cdot K = 0$ (the cross terms cancel since $L \cdot K = K \cdot L$, which follows from $L \cdot K = 0$ and $K \cdot L = L \cdot K + \sum_a [K_a, L_a] = L \cdot K$), so $j = j'$ by Proposition 12.2.1. Further $L^2 + K^2 = 2((J^{+})^2 + (J^{-})^2) = 4j(j+1)$, so $E = -\frac{1}{2(2j+1)^2}$, which determines j. Finally, $L = J^{+} + J^{-}$ acts on $D^{(j)} \otimes D^{(j)}$ as the tensor product representation, which decomposes by Corollary 3.3.3. □

(Fock showed that in fact $m = 1$, i.e., the degeneracy is exactly n^2.)

12.4 Isospin and the eightfold way

Physical input. Proton and neutron form a doublet $D^{(1/2)}$ of an approximate internal *isospin* symmetry SU(2), and the pions a triplet $D^{(1)}$. Gell-Mann and Ne'eman (1961) extended this to an approximate *flavour* symmetry SU(3); in the quark model, the quarks u, d, s span $\mathbf{3}$, the antiquarks $\bar{\mathbf{3}}$, mesons are $q\bar{q}$ states and baryons qqq states. Isospin I_3 and hypercharge Y are coordinates on $\mathfrak{t}^*$ for the maximal torus of SU(3).

By Example 11.3.2, mesons occur in $\mathbf{3} \otimes \bar{\mathbf{3}} = \mathbf{8} \oplus \mathbf{1}$ (e.g. the pseudoscalar octet π, K, η, and the singlet η'), and baryons in $\mathbf{3} \otimes \mathbf{3} \otimes \mathbf{3} = \mathbf{10} \oplus \mathbf{8} \oplus \mathbf{8} \oplus \mathbf{1}$ (the octet containing p, n, Λ, Σ, Ξ; the decuplet containing Δ, Σ^*, Ξ^*, Ω^-). The Ω^-, corresponding to the weight $3e_3$ of $\mathrm{Sym}^3\mathbb{C}^3$ (a corner of the weight diagram of $\mathbf{10}$), was predicted and then discovered in 1964.

A second, exact SU(3) symmetry, *colour*, is a gauge symmetry of quantum chromodynamics, and observed hadrons are colour singlets. Representation theory explains why quarks bind in pairs $q\bar{q}$ and triples qqq: by Example 11.3.2, $\mathbf{3} \otimes \bar{\mathbf{3}}$ and $\mathbf{3}^{\otimes 3}$ contain the trivial representation exactly once (for $\mathbf{3}^{\otimes 3}$, it is $\Lambda^3\mathbb{C}^3$), whereas $\mathbf{3} \otimes \mathbf{3} = \mathbf{6} \oplus \bar{\mathbf{3}}$ does not contain it.

Further reading: B. Simon, *Representations of finite and compact groups*, AMS, 1996 (main reference); J. F. Adams, *Lectures on Lie groups*, University of Chicago Press, 1969; T. Bröcker and T. tom Dieck, *Representations of compact Lie groups*, Springer, 1985; D. Bump, *Lie groups*, 2nd ed., Springer, 2013; W. Fulton and J. Harris, *Representation theory: A first course*, Springer, 1991; B. C. Hall, *Lie groups, Lie algebras, and representations*, 2nd ed., Springer, 2015; J. E. Humphreys, *Introduction to Lie algebras and representation theory*, Springer, 1972; A. W. Knapp, *Lie groups beyond an introduction*, 2nd ed., Birkhäuser, 2002; M. R. Sepanski, *Compact Lie groups*, Springer, 2007; S. Sternberg, *Group theory and physics*, Cambridge University Press, 1994.