

Predicate Logic
&
Math Background

Predicate Logic

Predicate logic over integer expressions:

a language of logical assertions, for example

$$\forall x. x + 0 = x$$

Why discuss predicate logic?

- It is an example of a simple language
- It has simple denotational semantics
- We will use it later in program specifications

Abstract Syntax

Describes the structure of a phrase
ignoring the details of its representation.

An abstract grammar for predicate logic over integer expressions:

$$\begin{aligned} \textit{intexp} ::= & 0 \mid 1 \mid \dots \\ & \mid \textit{var} \\ & \mid -\textit{intexp} \mid \textit{intexp} + \textit{intexp} \mid \textit{intexp} - \textit{intexp} \mid \dots \\ \textit{assert} ::= & \mathbf{true} \mid \mathbf{false} \\ & \mid \textit{intexp} = \textit{intexp} \mid \textit{intexp} < \textit{intexp} \mid \textit{intexp} \leq \textit{intexp} \mid \dots \\ & \mid \neg \textit{assert} \mid \textit{assert} \wedge \textit{assert} \mid \textit{assert} \vee \textit{assert} \\ & \mid \textit{assert} \Rightarrow \textit{assert} \mid \textit{assert} \Leftrightarrow \textit{assert} \\ & \mid \forall \textit{var}. \textit{assert} \mid \exists \textit{var}. \textit{assert} \end{aligned}$$

Resolving Notational Ambiguity

- Using parentheses: $(\forall x. (((x) + (0)) + 0) = (x))$

- Using precedence and parentheses: $\forall x. (x + 0) + 0 = x$

arithmetic operators ($*$ / rem ...) with the usual precedence
relational operators ($=$ \neq $<$ \leq ...)

\neg

\wedge

\vee

\Rightarrow

\Leftrightarrow

- The body of a quantified term extends to a delimiter.

Carriers and Constructors

- Carriers: sets of abstract phrases (e.g. *intexp*, *assert*)
- Constructors: specify abstract grammar productions

$$\begin{array}{lll} \textit{intexp} ::= 0 & \longrightarrow & c_0 \in \{\langle \rangle\} \rightarrow \textit{intexp} \\ \textit{intexp} ::= \textit{intexp} + \textit{intexp} & \longrightarrow & c_+ \in \textit{intexp} \times \textit{intexp} \rightarrow \textit{intexp} \end{array}$$

Note: Independent of the concrete pattern of the production:

$$\textit{intexp} ::= \mathbf{plus} \textit{intexp} \textit{intexp} \longrightarrow c_+ \in \textit{intexp} \times \textit{intexp} \rightarrow \textit{intexp}$$

- Constructors must be injective and have disjoint ranges
- Carriers must be either predefined or their elements must be constructible in finitely many constructor applications

Inductive Structure of Carrier Sets

With these properties of constructors and carriers,
carriers can be defined inductively:

$$\begin{aligned} \text{intexp}^{(0)} &= \{\} \\ \text{intexp}^{(j+1)} &= \{c_0\langle\rangle, \dots\} \cup \{c_+(x_0, x_1) \mid x_0, x_1 \in \text{intexp}^{(j)}\} \cup \dots \\ \text{assert}^{(0)} &= \{\} \\ \text{assert}^{(j+1)} &= \{c_{\text{true}}\langle\rangle, c_{\text{false}}\langle\rangle\} \\ &\quad \cup \{c_=(x_0, x_1) \mid x_0, x_1 \in \text{intexp}^{(j)}\} \cup \dots \\ &\quad \cup \{c_{\neg}(x_0) \mid x_0 \in \text{assert}^{(j)}\} \cup \dots \\ \\ \text{intexp} &= \bigcup_{j=0}^{\infty} \text{intexp}^{(j)} \\ \text{assert} &= \bigcup_{j=0}^{\infty} \text{assert}^{(j)} \end{aligned}$$

Denotational Semantics of Predicate Logic

The meaning of a term $e \in \text{intexp}$ is $\llbracket e \rrbracket_{\text{intexp}}$

i.e. the function $\llbracket - \rrbracket_{\text{intexp}}$ maps *intexp* objects to their meanings.

What is the set of meanings?

The meaning $\llbracket 5 + 37 \rrbracket_{\text{intexp}}$ of the term $\underbrace{5 + 37}$ could be the integer 42.
(that is, $c_+(c_5\langle \rangle, c_{37}\langle \rangle)$)

However the term $x + 5$ contains the **free variable** x ,
so the meaning of an *intexp* in general cannot be an integer...

Mathematical Background

- Sets
- Relations
- Functions
- Sequences
- Products and Sums

Sets

$x \in S$	membership	$\{\}$	the empty set
$x \in! S$	$S = \{x\}$	\mathbf{N}	natural numbers
$S \subseteq T$	inclusion	\mathbf{Z}	integers
$S \subseteq^{\text{fin}} T$	finite subset	\mathbf{B}	$= \{\text{true}, \text{false}\}$
$\{E \mid P\}$	set comprehension		
$S \cap T$	intersection	$= \{x \mid x \in S \text{ and } x \in T\}$	x is a bound variable
$S \cup T$	union	$= \{x \mid x \in S \text{ or } x \in T\}$	
$S - T$	difference	$= \{x \mid x \in S \text{ and not } x \in T\}$	
$\mathcal{P} S$	powerset	$= \{T \mid T \subseteq S\}$	
$m \text{ to } n$	integer range	$= \{x \mid m \leq x \text{ and } x \leq n\}$	

Generalized Set Operations

$$\cup S \stackrel{\text{def}}{=} \{x \mid \exists T \in S. x \in T\} \quad \bigg| \quad \cap S \stackrel{\text{def}}{=} \{x \mid \forall T \in S. x \in T\}$$

$$\bigcup_{i \in I} S \stackrel{\text{def}}{=} \cup \{S \mid i \in I\} \quad \bigg| \quad \bigcap_{i \in I} S \stackrel{\text{def}}{=} \cap \{S \mid i \in I\} \dots$$

$$\bigcup_{i=m}^n S \stackrel{\text{def}}{=} \bigcup_{i \in m \text{ to } n} S \quad \bigg| \quad \bigcap_{i=m}^n S \stackrel{\text{def}}{=} \bigcap_{i \in m \text{ to } n} S$$

$$\cup \{\} = \{\}$$

$$\cap \{\} \quad \text{meaningless}$$

Examples:

$$A \cup B = \cup \{A, B\}$$

$$\cup \{i \text{ to } (i + 1) \mid i \in \{j^2 \mid j \in 1 \text{ to } 3\}\} = \{1, 2, 4, 5, 9, 10\}$$

Relations

A **relation** ρ is a set of **primitive pairs** $[x, y]$.

$$\rho \text{ relates } x \text{ and } y \iff x \rho y \iff [x, y] \in \rho$$

$$\rho \text{ is an identity relation} \iff (\forall x, y. x \rho y \Rightarrow x = y)$$

$$\text{the identity on } S \quad I_S \stackrel{\text{def}}{=} \{[x, x] \mid x \in S\}$$

$$\text{the domain of } \rho \quad \text{dom } \rho \stackrel{\text{def}}{=} \{x \mid \exists y. x \rho y\}$$

$$\text{the range of } \rho \quad \text{ran } \rho \stackrel{\text{def}}{=} \{x \mid \exists y. y \rho x\}$$

$$\text{composition of } \rho \text{ with } \rho' \quad \rho' \cdot \rho \stackrel{\text{def}}{=} \{[x, z] \mid \exists y. x \rho y \text{ and } y \rho' z\}$$

$$\text{reflection of } \rho \quad \rho^\dagger \stackrel{\text{def}}{=} \{[y, x] \mid [x, y] \in \rho\}$$

Relations: Properties and Examples

$$(\rho_3 \cdot \rho_2) \cdot \rho_1 = \rho_3 \cdot (\rho_2 \cdot \rho_1)$$

$$\rho \cdot I_S \subseteq \rho \supseteq I_T \cdot \rho$$

$$\text{dom } I_S = S = \text{ran } I_S$$

$$I_T \cdot I_S = I_{T \cap S}$$

$$I_S^\dagger = I_S$$

$$(\rho^\dagger)^\dagger = \rho$$

$$(\rho_2 \cdot \rho_1)^\dagger = \rho_1^\dagger \cdot \rho_2^\dagger$$

$$\rho \cdot \{\} = \{\} = \{\} \cdot \rho$$

$$I_{\{\}} = \{\} = \{\}^\dagger$$

$$\text{dom } \rho = \{\} \Rightarrow \rho = \{\}$$

$$I_{\mathbf{N}} = \{[0, 0], [1, 1], [2, 2], \dots\}$$

$$< = \{[0, 1], [0, 2], [1, 2], \dots\}$$

$$\leq = \{[0, 0], [0, 1], [1, 1], [0, 2], \dots\}$$

$$\geq = \{[0, 0], [1, 0], [1, 1], [2, 0], \dots\}$$

$$< \subseteq \leq$$

$$< \cup I_{\mathbf{N}} = \leq$$

$$\leq \cap \geq = I_{\mathbf{N}}$$

$$< \cap \geq = \{\}$$

$$< \cdot \leq = <$$

$$\leq \cdot \leq = \leq$$

$$\geq = \leq^\dagger$$

Functions

A relation f is a **function** if

$$\forall x, x', x''. ([x, x'] \in f \text{ and } [x, x''] \in f) \Rightarrow x' = x''$$

If f is a function,

$$f\ x = y \iff f_x = y \iff f \text{ maps } x \text{ to } y \iff [x, y] \in f$$

I_S and $\{\}$ are functions.

If f and g are functions, then $g \cdot f$ is a function: $(g \cdot f)\ x = g(f\ x)$

f^\dagger is not necessarily a function:

consider $f = \{[\text{true}, \{\}], [\text{false}, \{\}]\}$

f is an **injection** if both f and f^\dagger are functions.

Notation for Functions

Typed abstraction: $\lambda x \in S. E \stackrel{\text{def}}{=} \{[x, E] \mid x \in S\}$

Defined only when E is defined for all $x \in S$

(consider $\lambda g \in \mathbb{N}. g \ 3$)

$$I_S = \lambda x \in S. x$$

$$g \cdot f = \lambda x \in \text{dom } f. g(f \ x), \text{ if } \text{ran } f \subseteq \text{dom } g.$$

Placeholder: E with a dash $(-)$ standing for the bound variable

$$g \ (-) \ h = \lambda x \in S. (g \ (x)) \ h \qquad - + 42 = \lambda x \in \mathbb{N}. x + 42$$

Variation of a function f : $[f \mid x : y] \ z = \begin{cases} y, & \text{if } z = x \\ f \ z, & \text{otherwise} \end{cases}$

$$\text{dom } [f \mid x : y] = (\text{dom } f) \cup \{x\}$$

$$\text{ran } [f \mid x : y] = ((\text{ran } f) - \{z \mid [x, z] \in f\}) \cup \{y\}$$

Sequences

$$[f \mid x_1 : y_1 \mid \dots \mid x_n : y_n] \stackrel{\text{def}}{=} [\dots [f \mid x_1 : y_1] \dots \mid x_n : y_n]$$

$$[x_1 : y_1 \mid \dots \mid x_n : y_n] \stackrel{\text{def}}{=} [\{\} \mid x_1 : y_1 \mid \dots \mid x_n : y_n]$$

$$\langle x_0, \dots x_{n-1} \rangle \stackrel{\text{def}}{=} [0 : x_0 \mid \dots n-1 : x_{n-1}]$$

$[] = \{\}$ — the empty function

$\langle \rangle = [] = \{\}$ — the empty sequence

$\langle x_0, \dots x_{n-1} \rangle$ — an *n*-tuple

$\langle x, y \rangle$ — a (non-primitive) pair

$\text{dom } \langle x_0, \dots x_{n-1} \rangle = 0 \text{ to } (n-1)$

$\langle x_0, \dots x_{n-1} \rangle_i = x_i$ when $i \in 0 \text{ to } (n-1)$

Products

Let θ be an indexed family of sets (a function with sets in its range).

The **Cartesian product** of θ is

$$\prod \theta \stackrel{\text{def}}{=} \{f \mid \text{dom } f = \text{dom } \theta \text{ and } \forall i \in \text{dom } \theta. f \ i \in \theta \ i\}$$

$$\prod \langle \mathbf{B}, \mathbf{B} \rangle$$

$$= \prod (\lambda x \in 0 \text{ to } 1. \mathbf{B})$$

$$= \{[0 : \text{true}, 1 : \text{true}], [0 : \text{true}, 1 : \text{false}], \\ [0 : \text{false}, 1 : \text{true}], [0 : \text{false}, 1 : \text{false}]\}$$

$$= \{\langle \text{true}, \text{true} \rangle, \langle \text{true}, \text{false} \rangle, \langle \text{false}, \text{true} \rangle, \langle \text{false}, \text{false} \rangle\}$$

More Products

$$\prod_{x \in T} S \stackrel{\text{def}}{=} \prod_{\lambda x \in T} S$$

$$S_1 \times \dots \times S_n \stackrel{\text{def}}{=} \prod_{i=1}^n \textcolor{red}{S_i}$$

$$\prod_{i=m}^n S \stackrel{\text{def}}{=} \prod_{i \in (m \text{ to } n)} S$$

$$S^T \stackrel{\text{def}}{=} \prod_{x \in T} S$$

$$S^n \stackrel{\text{def}}{=} S^{0 \text{ to } (n-1)} = \underbrace{S \times \dots \times S}_{n \text{ times}}$$

$$\Pi\langle \mathbf{B}, \mathbf{B} \rangle = \mathbf{B} \times \mathbf{B} = \mathbf{B}^2$$

$$S^0 = S^{\{\}} = \{\langle \rangle\} = \{\{\}\}$$

Sets of Sequences

Let $\mathbf{U} = \{\langle \rangle\}$

$$S^+ \stackrel{\text{def}}{=} \bigcup_{i=1}^{\infty} S^i$$

$$\mathbf{U}^+ = \{\langle \langle \rangle \rangle, \langle \langle \rangle, \langle \rangle \rangle, \langle \langle \rangle, \langle \rangle, \langle \rangle \rangle, \dots \text{(finite)}\}$$

$$S^* \stackrel{\text{def}}{=} S^0 \cup S^+$$

$$\mathbf{U}^* = \{\langle \rangle, \langle \langle \rangle \rangle, \langle \langle \rangle, \langle \rangle \rangle, \langle \langle \rangle, \langle \rangle, \langle \rangle \rangle, \dots \text{(finite)}\}$$

$$S^{\infty} \stackrel{\text{def}}{=} S^* \cup S^{\mathbf{N}}$$

$$\mathbf{U}^{\infty} = \{\langle \rangle, \langle \langle \rangle \rangle, \langle \langle \rangle, \langle \rangle \rangle, \langle \langle \rangle, \langle \rangle, \langle \rangle \rangle, \dots \text{(infinite)}\}$$

Sums

The **disjoint union** (**sum**) of θ is

$$\Sigma \theta \stackrel{\text{def}}{=} \{ \langle i, x \rangle \mid i \in \text{dom } \theta \text{ and } x \in \theta i \}$$

$$\sum_{x \in T} S \stackrel{\text{def}}{=} \sum \lambda x \in T. S \qquad S_1 + \dots + S_n \stackrel{\text{def}}{=} \sum_{i=1}^n \textcolor{red}{S_i}$$

$$\sum_{i=m}^n S \stackrel{\text{def}}{=} \sum_{i \in (m \text{ to } n)} S \qquad \textcolor{blue}{T \times S} = \sum_{x \in T} S$$

$$n \times S = (0 \text{ to } (n - 1)) \times S = \underbrace{S + \dots + S}_{n \text{ times}}$$

$$\begin{aligned} \mathbf{B} + \mathbf{B} &= \Sigma \langle \mathbf{B}, \mathbf{B} \rangle = \{ \langle 0, \text{true} \rangle, \langle 0, \text{false} \rangle, \langle 1, \text{true} \rangle, \langle 1, \text{false} \rangle \} \\ &= 2 \times \mathbf{B} \end{aligned}$$

Sums

Let θ be an indexed family of sets (a function with sets in its range).

The **disjoint union** (**sum**) of θ is

$$\Sigma \theta \stackrel{\text{def}}{=} \{ \langle i, x \rangle \mid i \in \text{dom } \theta \text{ and } x \in \theta i \}$$

$$\sum_{x \in T} S \stackrel{\text{def}}{=} \sum \lambda x \in T. S \qquad S_1 + \dots + S_n \stackrel{\text{def}}{=} \sum_{i=1}^n \textcolor{red}{S_i}$$

$$\sum_{i=m}^n S \stackrel{\text{def}}{=} \sum_{i \in (m \text{ to } n)} S \qquad \textcolor{blue}{T \times S} = \sum_{x \in T} S$$

$$n \times S = (0 \text{ to } (n - 1)) \times S = \underbrace{S + \dots + S}_{n \text{ times}}$$

$$\begin{aligned} \mathbf{B} + \mathbf{B} &= \Sigma \langle \mathbf{B}, \mathbf{B} \rangle = \{ \langle 0, \text{true} \rangle, \langle 0, \text{false} \rangle, \langle 1, \text{true} \rangle, \langle 1, \text{false} \rangle \} \\ &= 2 \times \mathbf{B} \end{aligned}$$

Functions of Multiple Arguments

- Use **tuples** instead of multiple arguments:

$$f (a_0, \dots a_{n-1}) \longrightarrow f \langle a_0, \dots a_{n-1} \rangle$$

Syntactic sugar:

$$\lambda \langle x_0 \in S_0, \dots, x_{n-1} \in S_{n-1} \rangle. E$$

$$\stackrel{\text{def}}{=} \lambda x \in S_0 \times \dots \times S_{n-1}. (\lambda x_0 \in S_0. \dots \lambda x_{n-1} \in S_{n-1}. E) \\ (x\ 0) \dots (x(n-1))$$

- Use **Currying**:

$$f (a_0, \dots a_{n-1}) \longrightarrow f\ a_0\ \dots\ a_{n-1} \\ = (\dots (f\ a_0)\ \dots)\ a_{n-1}$$

where f is a Curried function $\lambda x_0 \in S_0. \dots \lambda x_{n-1} \in S_{n-1}. E$.

Relations Between Sets

ρ is a relation from S to T

$$\iff \rho \in S \xrightarrow{\text{REL}} T$$

$$\iff \text{dom } \rho \subseteq S \text{ and } \text{ran } \rho \subseteq T.$$

Relation on $S \stackrel{\text{def}}{=} \text{relation from } S \text{ to } S.$

$$I_S \in S \xrightarrow{\text{REL}} S$$

$$\rho \in S \xrightarrow{\text{REL}} T \Rightarrow \rho^\dagger \in T \xrightarrow{\text{REL}} S$$

$$\text{For all } S \text{ and } T, \{\} \in S \xrightarrow{\text{REL}} T$$

$$\{\} \in! S \xrightarrow{\text{REL}} \{\}$$

$$\{\} \in! \{\} \xrightarrow{\text{REL}} T$$

Total Relations


$\rho \in S \xrightarrow{\text{REL}} T$ is a **total relation from S to T**

$$\iff \rho \in S \xrightarrow{\text{TREL}} T$$

$$\iff \forall x \in S. \exists y \in T. x \rho y$$

$$\iff \text{dom } \rho = S$$

$$\iff I_S \subseteq \rho^\dagger \cdot \rho$$



$$\rho \in (\text{dom } \rho) \xrightarrow{\text{TREL}} T \iff T \supseteq \text{ran } \rho$$

Functions Between Sets

f is a **partial function from S to T**

$$\iff f \in S \xrightarrow{\text{PFUN}} T$$

$$\iff f \in S \xrightarrow{\text{REL}} T \text{ and } f \text{ is a function.}$$

“Partial”: $f \in S \xrightarrow{\text{REL}} T \Rightarrow \text{dom } f \subseteq S$

$f \in S \xrightarrow{\text{PFUN}} T$ is a **(total) function from S to T**

$$\iff f \in S \rightarrow T$$

$$\iff \text{dom } f = S.$$

- $S \rightarrow T = T^S = \prod_{x \in S} T$
- $S \rightarrow T \rightarrow U = S \rightarrow (T \rightarrow U)$


Surjections, Injections, Bijections

f is a **surjection** from S to $T \iff \text{ran } f = T$

f is a **injection** from S to $T \iff f^\dagger \in T \xrightarrow{\text{PFUN}} S$

f is a **bijection** from S to $T \iff f^\dagger \in T \rightarrow S$

$\iff f$ is an **isomorphism** from S to T



Back to Predicate Logic

$intexp ::= 0 \mid 1 \mid \dots$

$\mid var$

$\mid -intexp \mid intexp + intexp \mid intexp - intexp \mid \dots$

$assert ::= \mathbf{true} \mid \mathbf{false}$

$\mid intexp = intexp \mid intexp < intexp \mid intexp \leq intexp \mid \dots$

$\mid \neg assert \mid assert \wedge assert \mid assert \vee assert$

$\mid assert \Rightarrow assert \mid assert \Leftrightarrow assert$

$\mid \forall var. assert \mid \exists var. assert$

Denotational Semantics of Predicate Logic

The meaning of term $e \in \text{intexp}$ is $\llbracket e \rrbracket_{\text{intexp}}$

i.e. the function $\llbracket - \rrbracket_{\text{intexp}}$ maps objects from intexp to their meanings.

What is the set of meanings?

The meaning $\llbracket 5 + 37 \rrbracket_{\text{intexp}}$ of the term $5 + 37$ could be the integer 42.

But the term $x + 5$ contains the free variable x ...

Environments

...hence we need an **environment** (variable assignment, state)

$$\sigma \in \Sigma \stackrel{\text{def}}{=} \text{var} \rightarrow \mathbf{Z}$$

to give meaning to free variables.

The meaning of a term is a function from the states to \mathbf{Z} or \mathbf{B} .

$$\begin{array}{ll} \llbracket - \rrbracket_{intexp} & \in \quad intexp \rightarrow \Sigma \rightarrow \mathbf{Z} \\ \llbracket - \rrbracket_{assert} & \in \quad assert \rightarrow \Sigma \rightarrow \mathbf{B} \end{array}$$

$$\begin{array}{ll} \text{if } \sigma = [x : 3, y : 4], & \text{then } \llbracket x+5 \rrbracket_{intexp} \sigma = 8 \\ & \llbracket \exists z. x < z \wedge z < y \rrbracket \sigma = \text{false} \end{array}$$

Direct Semantics Equations for Predicate Logic

$v \in \text{var}$

$e \in \text{intexp}$

$p \in \text{assert}$

$$\begin{aligned} \llbracket 0 \rrbracket_{\text{intexp}} \sigma &= 0 \\ (\text{really } \llbracket c_0 \langle \rangle \rrbracket_{\text{intexp}} \sigma &= 0) \end{aligned}$$

$$\llbracket v \rrbracket_{\text{intexp}} \sigma = \sigma v$$

$$\llbracket e_0 + e_1 \rrbracket_{\text{intexp}} \sigma = \llbracket e_0 \rrbracket_{\text{intexp}} \sigma + \llbracket e_1 \rrbracket_{\text{intexp}} \sigma$$

$$\llbracket \text{true} \rrbracket_{\text{assert}} \sigma = \text{true}$$

$$\llbracket e_0 = e_1 \rrbracket_{\text{assert}} \sigma = \llbracket e_0 \rrbracket_{\text{intexp}} \sigma = \llbracket e_1 \rrbracket_{\text{intexp}} \sigma$$

$$\llbracket \neg p \rrbracket_{\text{assert}} \sigma = \neg(\llbracket p \rrbracket_{\text{assert}} \sigma)$$

$$\llbracket p_0 \wedge p_1 \rrbracket_{\text{assert}} \sigma = \llbracket p_0 \rrbracket_{\text{assert}} \sigma \wedge \llbracket p_1 \rrbracket_{\text{assert}} \sigma$$

$$\llbracket \forall v. p \rrbracket_{\text{assert}} \sigma = \forall n \in \mathbf{Z}. \llbracket p \rrbracket_{\text{assert}} [\sigma | v : n]$$

Example: The Meaning of a Term

$$\llbracket \forall x. x+0=x \rrbracket_{assert} \sigma$$

$$= \forall n \in \mathbf{Z}. \llbracket x+0=x \rrbracket_{assert} [\sigma | x : n]$$

$$= \forall n \in \mathbf{Z}. \llbracket x+0 \rrbracket_{intexp} [\sigma | x : n] = \llbracket x \rrbracket_{intexp} [\sigma | x : n]$$

$$= \forall n \in \mathbf{Z}. \llbracket x \rrbracket_{intexp} [\sigma | x : n] + \llbracket 0 \rrbracket_{intexp} [\sigma | x : n] = \llbracket x \rrbracket_{intexp} [\sigma | x : n]$$

$$= \forall n \in \mathbf{Z}. [\sigma | x : n](x) + 0 = [\sigma | x : n](x)$$

$$= \forall n \in \mathbf{Z}. n + 0 = n$$

$$= \mathbf{true}$$

Properties of the Semantic Equations

- They are **syntax-directed** (**homomorphic**):
 - exactly one equation for each abstract grammar production (constructor)
 - result expressed using functions (meanings) of subterms only (arguments of constructor)

\Rightarrow they have exactly one solution $\langle \llbracket - \rrbracket_{intexp}, \llbracket - \rrbracket_{assert} \rangle$
(proof by induction on the structure of terms).
- They define **compositional** semantic functions
(depending only on the **meaning** of the subterms)

\Rightarrow “equivalent” subterms can be substituted

Validity of Assertions

p holds/is true in $\sigma \iff \sigma$ satisfies $p \iff \llbracket p \rrbracket_{assert} \sigma = \mathbf{true}$

p is valid $\iff \forall \sigma \in \Sigma. p$ holds in σ

p is unsatisfiable $\iff \forall \sigma \in \Sigma. \llbracket p \rrbracket_{assert} \sigma = \mathbf{false}$

$\iff \neg p$ is valid

p is stronger than p' $\iff \forall \sigma \in \Sigma. (p' \text{ holds if } p \text{ holds})$

$\iff (p \Rightarrow p')$ is valid

p and p' are equivalent $\iff p$ is stronger than p'
and p' is stronger than p

Inference Rules

<i>Class</i>	<i>Examples</i>
$\vdash p$ (Axiom) $\frac{}{\vdash p}$ (Axiom Schema) $\frac{\vdash p_0 \quad \dots \quad \vdash p_{n-1}}{\vdash p}$ (Rule)	$\vdash x + 0 = x$ (xPlusZero) $\frac{}{\vdash e_1 = e_0 \Rightarrow e_0 = e_1}$ (SymmObjEq) $\frac{\vdash p \quad \vdash p \Rightarrow p'}{\vdash p'}$ (ModusPonens) $\frac{\vdash p}{\vdash \forall v. p}$ (Generalization)

Formal Proofs

A set of inference rules defines a **logical theory** \vdash .

A **formal proof** (in a logical theory):

a sequence of **instances** of the inference rules, where the premisses of each rule occur as conclusions earlier in the sequence.

1. $\vdash x + 0 = x$ (xPlusZero)
2. $\vdash x + 0 = x \Rightarrow x = x + 0$ (SymmObjEq)
[$e_0 : x \mid e_1 : x + 0$]
3. $\vdash x = x + 0$ (ModusPonens, 1, 2)
[$p : x + 0 = x \mid p' : x = x + 0$]
4. $\vdash \forall x. x = x + 0$ (Generalization, 3)
[$v : x \mid p : x = x + 0$]

Tree Representation of Formal Proofs

$$\frac{\frac{\frac{\vdash x + 0 = x}{\vdash x = x + 0} \text{ (MP)}}{\vdash \forall x. x = x + 0} \text{ (Gen)}}{\vdash x + 0 = x \Rightarrow x = x + 0} \text{ (SymmObjEq)}$$

Soundness of a Logical Theory

An inference rule is **sound** if in every instance of the rule the conclusion is valid if all the premisses are.

A logical theory \vdash is sound if all inference rules in it are sound.

If \vdash is sound and there is a formal proof of $\vdash p$, then p is valid.

Object vs Meta implication:

$\vdash p \Rightarrow \forall v. p$ is not a sound rule, although $\frac{\vdash p}{\vdash \forall v. p}$ is.

Completeness of a Logical Theory

A logical theory \vdash is **complete** if

for every valid p there is a formal proof of $\vdash p$.



A logical theory \vdash is **axiomatizable** if

there exists a finite set of inference rules



from which can be constructed formal proofs of all assertions in \vdash .

No first-order theory of arithmetic is complete and axiomatizable.

Variable Binding



Variable Binding



Bound and Free Variables

In $\forall v. p$, v is the **binding occurrence** (**binder**) and p is its **scope**.

If a non-binding occurrence of v is within the scope of a binder for v , then it is a **bound** occurrence; otherwise it's a **free** one.

$$FV_{interp}(0) = \{\}$$

$$FV_{assert}(\mathbf{true}) = \{\}$$

$$FV(v) = \{v\}$$

$$FV(e_0 = e_1) = FV(e_0) \cup FV(e_1)$$

$$FV(-e) = FV(e)$$

$$FV(\neg p) = FV(p)$$

$$FV(e_0 + e_1) = FV(e_0) \cup FV(e_1) \quad FV(p_0 \wedge p_1) = FV(p_0) \cup FV(p_1)$$

$$FV(\forall v. p) = FV(p) - \{v\}$$

Example:

$$FV(\exists y. x < y \wedge \exists x. x > y) = \{x\}$$

Only Assignment of Free Variables Matters

Coincidence Theorem:

If $\sigma v = \sigma' v$ for all $v \in FV_\theta(p)$, then $\llbracket p \rrbracket_\theta \sigma = \llbracket p \rrbracket_\theta \sigma'$
(where p is a phrase of type θ).

Proof: By structural induction.

Inductive hypothesis:

The statement of the theorem holds
for all phrases of depth less than that of the phrase p' .

Base cases:

$$p' = 0 \Rightarrow \llbracket 0 \rrbracket_{intexp} \sigma = 0 = \llbracket 0 \rrbracket_{intexp} \sigma'$$

$$p' = v \Rightarrow \llbracket v \rrbracket_{intexp} \sigma = \sigma v = \sigma' v = \llbracket v \rrbracket_{intexp} \sigma', \text{ since } FV(v) = \{v\}.$$

Proof of Concidence Theorem, cont'd

Coincidence Theorem:

If $\sigma v = \sigma' v$ for all $v \in FV_\theta(p)$, then $\llbracket p \rrbracket_\theta \sigma = \llbracket p \rrbracket_\theta \sigma'$.

Inductive cases:

$p' = e_0 + e_1$: by IH $\llbracket e_i \rrbracket_{intexp} \sigma = \llbracket e_i \rrbracket_{intexp} \sigma', i \in \{1, 2\}$.

$$\begin{aligned}\llbracket p' \rrbracket_{intexp} \sigma &= \llbracket e_0 \rrbracket_{intexp} \sigma + \llbracket e_1 \rrbracket_{intexp} \sigma \\ &= \llbracket e_0 \rrbracket_{intexp} \sigma' + \llbracket e_1 \rrbracket_{intexp} \sigma' = \llbracket p' \rrbracket_{intexp} \sigma'\end{aligned}$$

$p' = \forall u. q$: $\sigma v = \sigma' v, \quad \forall v \in FV(p') = FV(q) - \{u\}$
then $[\sigma|u : n]v = [\sigma'|u : n]v, \forall v \in FV(q), n \in \mathbf{Z}$

Then by IH $\llbracket q \rrbracket_{assert} [\sigma|u : n] = \llbracket q \rrbracket_{assert} [\sigma'|u : n]$ for all $n \in \mathbf{Z}$,

hence $\forall n \in \mathbf{Z}. \llbracket q \rrbracket_{assert} [\sigma|u : n] = \forall n \in \mathbf{Z}. \llbracket q \rrbracket_{assert} [\sigma'|u : n]$

$$\llbracket \forall u. q \rrbracket_{assert} \sigma = \llbracket \forall u. q \rrbracket_{assert} \sigma'.$$

Substitution

$$\left. \begin{array}{l} -/\delta \in \textit{intexp} \rightarrow \textit{intexp} \\ -/\delta \in \textit{assert} \rightarrow \textit{assert} \end{array} \right\} \text{when } \delta \in \textit{var} \rightarrow \textit{intexp}$$

$$0/\delta = 0$$

$$v/\delta = \delta v$$

$$(-e)/\delta = -(e/\delta)$$

$$(p_0 \wedge p_1)/\delta = (p_0/\delta) \wedge (p_1/\delta)$$

$$(e_0 + e_1)/\delta = (e_0/\delta) + (e_1/\delta)$$

$$(\forall v. p)/\delta = \forall v'. (p/[\delta|v : v']),$$

...

$$\text{where } v' \notin \bigcup_{u \in FV(p) - \{v\}} FV(\delta u)$$


Examples:

$$(x < 0 \wedge \exists x. x \leq y)/[x : y+1] = y+1 < 0 \wedge \exists x. x \leq y$$


$$(x < 0 \wedge \exists x. x \leq y)/[y : x+1] = x < 0 \wedge \exists z. z \leq x+1$$

Preserving Binding Structure

$$(x < 0 \wedge \exists x. x \leq y) / [\boxed{x} : y+1] = y+1 < 0 \wedge \exists x. x \leq y$$




\longrightarrow




Avoiding Variable Capture

$$(x < 0 \wedge \exists x. x \leq y) / [\boxed{y} : x+1] = x < 0 \wedge \exists z. z \leq x+1$$



\longrightarrow



Substitution Theorems

Substitution Theorem:

If $\sigma = \llbracket - \rrbracket_{intexp} \sigma' \cdot \delta$ on $FV(p)$, then $(\llbracket - \rrbracket \sigma)p = (\llbracket - \rrbracket \sigma' \cdot (-/\delta))p$.

Finite Substitution Theorem:

$$\llbracket p/v_0 \rightarrow e_0, \dots v_{n-1} \rightarrow e_{n-1} \rrbracket \sigma = \llbracket p \rrbracket [\sigma|v_0 : \llbracket e_0 \rrbracket \sigma, \dots].$$

where

$$p/v_0 \rightarrow e_0, \dots v_{n-1} \rightarrow e_{n-1} \stackrel{\text{def}}{=} p/[cvar|v_0 : e_0 | \dots | v_{n-1} : e_{n-1}].$$

Renaming:

If $u \notin FV(q) - \{v\}$, then $\llbracket \forall u. (q/v \rightarrow u) \rrbracket_{boolexp} = \llbracket \forall v. q \rrbracket_{boolexp}$.