
Supplementary Figure S13 | A state-resolved secondary IL12RB2 regulatory component

A


Credible-set regulatory architecture

Variant	Signal	PBC PIP	ENCODE cCRE	Activated T-cell IL12RB2 E2G
rs6679356	L1	0.94	—	—
rs6702599	L1	0.06	—	—
rs3828069	L2	0.58	—	—
rs3790569	L2	0.26	dELS	—
rs11209064	L2	0.16	dELS	8
rs112588573	L3	0.82	—	—
rs72932868	L3	0.14	—	—


B



C



D



E

Cross-autoimmune context at 1p31.3

Trait	Regional lead P	max r² to PBC	PP.H3	PP.H4	Signal interpretation
PSC	P = 2.53e-04	0.005	0.205	0.017	No detectable regional association
RA	P = 1.25e-05	Not available	0.077	0.016	No detectable regional association
IBD	P = 4.59e-111	0.039	1.000	0.000	Distinct IL23R-side component
Psoriasis	P = 4.09e-49	0.037	1.000	0.000	Distinct IL23R-side component

Default coloc prior $p_{12} = 1 \times 10^{-5}$. IBD and psoriasis support H3, not a shared PBC signal.

Activation-state eQTL and LD compatibility are supportive; no formal genotype-by-stimulation interaction was established.