

Sparsity-Promoting Dynamic Mode Decomposition Applied to Sea Surface Temperature Fields

Zhicheng Zhang*

(Joint work with Yoshihiko Susuki*, & Atsushi Okazaki[†])

**Department of Electrical Engineering, Kyoto University*

[†]*Institute of Advanced Academic Research, Chiba University*

2025 SICE Festival with Annual Conference (SICE FS'25)

September 11, 2025, Chiang Mai, Thailand

Purpose and Outline of Presentation

Goal: Koopman/Dynamic mode decomposition (K/DMD) applied to climate systems.

- **Ques 1:** Why should we capture the **dominant** Koopman modes embedded in sea surface temperature (SST) data field?
(i.e., Why use KMD? Why explore SST?)
- **Ques 2:** Why we focus on Sparsity-Promoting technique?
 - I. Background on long-term SST and its real-world (physical information)
(e.g., monthly, seasonal, annual cycle of SST data)
 2. Review of **Sparsity-Promoting** Dynamic Mode Decomposition (SPDMD)
 3. Discussion on extracted dominant Koopman modes
(including **accuracy & model reduction**, **reconstruction**, etc...)
 4. Summary



State-of-the-Art Reviews on Climate Systems

nature communications



Article

<https://doi.org/10.1038/s41467-024-48033-6>

Revealing trends and persistent cycles of non-autonomous systems with autonomous operator-theoretic techniques

Received: 13 August 2023

Gary Froyland¹✉, Dimitrios Giannakis^{2,3}, Edoardo Luna⁴ & Joanna Slawinska²

Accepted: 16 April 2024

Exponentially decaying modes

2021 NAVARRA ET AL.

Estimation of Koopman Transfer Operators for the Equatorial Pacific SST

ANTONIO NAVARRA,^{a,b} JOE TRIBBIA,^c AND STEFAN KLUS^d

ARTICLE

<https://doi.org/10.1038/s41467-021-26357-x>

OPEN

Spectral analysis of climate dynamics with operator-theoretic approaches

Gary Froyland¹, Dimitrios Giannakis^{2,3✉}, Benjamin R. Lintner⁴, Maxwell Pike⁴ & Joanna Slawinska^{5,6,7}

Giannakis^{1*}

Geophysical Research Letters®

RESEARCH LETTER

10.1029/2023GL102743

Key Points:

- An operator-theoretic approach,

Identification of the Madden–Julian Oscillation With Data-Driven Koopman Spectral Analysis

Benjamin R. Lintner^{1,2} , Dimitrios Giannakis³ , Max Pike¹, and Joanna Slawinska³

SCIENTIFIC REPORTS

natureresearch

Check for updates

2024

NAVARRA ET AL.

Variability of SST through Koopman Modes

ANTONIO NAVARRA,^{a,b} JOE TRIBBIA,^c STEFAN KLUS,^d AND PAULA LORENZO-SÁNCHEZ^{a,e}

SCIENTIFIC REPORTS

natureresearch



Statistical ENSO operator- methods for nonlinear

Recently, (Koopman) operator theoretic methods are being increasingly popular and practical in climate science and weather forecasting!


So... Why?




Why Koopman Operator for Time-Series Modeling Works?



Climate systems go beyond toy model (Chaotic system)!




Observable Space $f, g \in \mathcal{F}$



Linear evolution of the observables

$$U\phi_i = \lambda_i\phi_i$$


$$U^t f = f \circ T^t$$




- The (flow) dynamics acts on **finite-dim. state-space**, while the system is **nonlinear**.
- Koopman operator *lifts* a nonlinear system into a **linear representation** acting on the **space of observables**, but it is **infinite-dimensional**.

Background: Why is (Niño-3) SST Data worth Exploring?



Atmosphere-ocean feedbacks during El Niño-Southern Oscillation
El Niño



Atmosphere-ocean feedbacks during El Niño-Southern Oscillation
Neutral



Atmosphere-ocean feedbacks during El Niño-Southern Oscillation
La Niña



Why SST data?

- ❑ Niño-3 tracks SST anomalies in the eastern tropical Pacific (5°N – 5°S , 150°W – 90°W), is a vital dataset for understanding **global climate patterns**, such as El Niño-Southern Oscillation (ENSO).



Purpose:

- ❑ Use Koopman Mode Decomposition (KMD) capture **dominant climate patterns** by using a few modes!

Ref.) Pictures comes from NOAA Climate, <https://www.climate.gov/news-features/blogs/enso/>


Animation takes from NASA Scientific Visualization Studio, https://svs.gsfc.nasa.gov/4695#section_credits

Niño-3 Index Sea Surface Temperature (SST) Data



Nino-3 index (Original) Data:



- Area Region:
Latitude: **5N-5S**, Longitude: **90W-150W**
- Total Grids (Lat. x Longt.):
10 x 60 = 600 grids
- Month/Year Range:
Jan. 1891 to Dec. 2019 (129 yrs)
- Snapshots: t = 1548/mths
- Climate modes: ENSO events, etc...



Niño 3 index Observed SST Data

Time-Series Data Collection: Data-Driven Climate Modeling

Niño 3 index (Monthly) SST Data Field



Data Processing:

- Extract the snapshot matrices from time-series
- Stack the data matrix into a high-dim. vector y
- Span the vectors as a new big matrix \mathbf{Y} .


$$\mathbf{Y} := \begin{Bmatrix} | & | & | \\ y_0, y_1, \dots, y_{N-1} \\ | & | & | \end{Bmatrix}$$

Matrix \leftarrow Vectors


$$y_t \in \mathbb{R}^{p \times 1}, t = 0, 1, N - 1$$

Method: Dynamic Mode Decomposition (DMD)


Data $p \times N$




Modes $p \times r$



\approx



Amplitudes
 $r \times r$



Dynamics
 $r \times N$

Refs.) Rowley et al, Spectral analysis of nonlinear flows, *J. Fluid Mech.*'2009
Schmid, Dynamic mode decomposition of numerical and experimental data, *J. Fluid Mech.*'2010

Singular Value Decomposition (SVD)

$$y_t \approx \sum_{j=1}^N \phi_j \lambda_j^t b_j \quad \xrightarrow{\text{SVD}} \quad y_t \approx \sum_{j=1}^r \Phi_j \Sigma_j^t b_j$$

$$\mathbf{Y} = \mathbf{U}\Sigma\mathbf{V}^*$$

Linear Time Invariant (LTI):

$$\mathbf{Y}^+ = \mathbf{AY} \quad \xleftarrow{} \quad y_{t+1} = \mathbf{Ay}_t$$

$$\mathbf{Y} \approx \Phi \text{Diag}(\mathbf{b}) \Sigma$$

$$J(\mathbf{b}) = \min_{\mathbf{b}} \|\mathbf{Y} - \Phi \text{Diag}(\mathbf{b}) \Sigma\|_F^2$$

➤ DMD via Least Square Error Optimization

$$\underbrace{\begin{bmatrix} | & \cdots & | \\ y_0 & \cdots & y_{N-1} \\ | & \cdots & | \end{bmatrix}}_{\mathbf{Y} \in \mathbb{R}^{p \times N}} \approx \underbrace{\begin{bmatrix} | & \cdots & | \\ \Phi_1 & \cdots & \Phi_r \\ | & \cdots & | \end{bmatrix}}_{\Phi} \underbrace{\begin{bmatrix} b_1 & & \\ & \ddots & \\ b_r & & \end{bmatrix}}_{\text{Diag}(\mathbf{b})} \underbrace{\begin{bmatrix} 1 & \lambda_1 & \cdots & \lambda_1^{N-1} \\ \vdots & \vdots & \ddots & \vdots \\ 1 & \lambda_r & \cdots & \lambda_r^{N-1} \end{bmatrix}}_{\Sigma}$$




Connection between DMD & KMD:


- ❖ DMD is a **numerical** counterparts of Koopman mode decomposition (KMD)
- ❖ Koopman operator: **linear evolutions of observables** (directly measured from data)
- ❖ DMD modes \rightarrow Koopman modes

Method: Sparsity-Promoting Dynamic Mode Decomposition (SPDMD)

Data $p \times N$




Modes $p \times r$



\approx

Amplitudes
 $r \times r$

Diag(\mathbf{b})




Sparse Amps.

$$S \times S \\ S \ll r$$

Ref.) Kutz et al, Dynamic mode decomposition: Data-driven modeling of complex systems, SIAM'2016
Jovanovic, Schmid & Nichols, Sparsity-promoting dynamic mode decomposition, Physics. Fluids'2014

$$J(\mathbf{b}) = \min_{\mathbf{b}} \|\mathbf{Y} - \Phi \text{Diag}(\mathbf{b}) \mathbf{\Xi}\|_F^2$$



Dynamics
 $r \times N$





- Promote sparsity on “Amplitudes”!
- Take L1 norm regularization!
(Convex program, ADMM Algorithm)

\mathbf{Y}


\approx

Φ



\approx

$\mathbf{\Xi}$



$$J_\gamma(\mathbf{b}) = \min_{\mathbf{b}} \|\mathbf{Y} - \Phi \text{Diag}(\mathbf{b}) \mathbf{\Xi}\|_F^2 + \gamma \sum_{i=1}^r |b_i|$$

- Dominant modes determined by order of the *magnitude of amplitudes* $|b_i|, i = 1, \dots, r$
- Associated with leading pairs (λ_i, ϕ_i)

Numerical Benchmark : Seasonal Cycle of SST Data Experiment

Seasonal cycle SST data (3-mth period)

$y_t \in \mathbb{R}^{1800}, p = 1800, N = 515$



Tall and **Skinny** matrix: $\mathbf{Y} \in \mathbb{R}^{1800 \times 515}$ estimates \mathbf{A}

One column denotes 3-mth (medium-resolution)


DMD (●), SPDMD (★)

DMD and SPDMD estimates the Koopman SST eigenvalues distributed around unit cycle!

Seasonal SST data enjoys **steady state!**



Extraction: Dominant Koopman (Spatial & Temporal) Modes



- SPDMD extracts leading **spatial** & **temporal** climate modes.
 - It helps us understand the variability of climate dynamics, e.g., **ENSO** (El Niño, La Niña events)

Trade-off: Accuracy vs. Model Reduction

















Table 2: Model reduction with the number of modes, the cost, and the performance loss in seasonal cycle of SST.

Weight γ	Modes # i	Cost $\mathcal{J}(\mathbf{b}^*)$	Loss $\Pi\%$
0.0001	511	2.2308×10^4	0.6010
0.0105	501	1.1602×10^5	0.8277
0.5503	490	1.8414×10^5	1.0224
8.5573	488	2.2331×10^5	1.1172
100.24	478	2.3250×10^5	1.1983
691.05	303	3.4864×10^5	1.8255
1714.3	107	4.7381×10^5	2.3815
4948.2	19	7.7276×10^5	3.0350
5757.2	9	8.6025×10^5	3.3761
16000	4	1.6447×10^6	3.5880

Ques:

- Does the SPDMD method perform effectively?
- How does the number of modes (few/many) impact the accuracy of the climate model?

SPDMD Reproducibility: Reconstruction, Validation of Seasonal SST



Abs(-) / Magnitude of Reconstructed Data

28

26

24

22

20

28

26

24

22

20

28

26

24

22

20

28

26

24

22

20

28

26


24

22


20

The Take-Home Messages:


- ✓ Koopman analysis for SST data by using SPDMD method to identify the dominant (spatial & temporal) modes.
- ✓ The captured Koopman modes (growth/decay) reveal climate patterns, suggesting the climate changes/variability...
- ✓ By reconstruction, KMD with SPDMD is effective method to make trade-off between accuracy and model reduction!




○ #259 Modes



Original SST (#515 Modes)



▶ #57 Modes



arXiv:2507.05711
SPDMD applied to SST

Thank you!

ขอบคุณครับ/ค่ะ

Suggestions and Comments are Welcome!

zhang.zhicheng.2c@kyoto-u.ac.jp



京都大学
KYOTO UNIVERSITY

Backup Slides


Koopman Operator: A (Bounded) Linear Operator

Koopman Framework


$$x_{t+1} = T(x_t)$$

REALIZATION LIFTING
 $\mathcal{K}^t g = g \circ T^t$
 $g_{t+1} = \mathcal{K}^t g_t = g \circ T = g(x_{t+1})$

- Lifts the **nonlinear evolution of state** into a **linear evolution of functional observable**
- Linearity: $\mathcal{K}^t(\alpha_1 g_1 + \alpha_2 g_2) = \alpha_1 \mathcal{K}^t g_1 + \alpha_2 \mathcal{K}^t g_2$
 $\forall \alpha_1, \alpha_2 \in \mathbb{R}, g_1, g_2 \in \mathcal{O}$



Refs.) Koopman, PNAS'1931; Koopman & von Neumann, PNAS'1932; Mezic, Nonlinear Dynamics'05



Point: Capture the system's dominant behavior via linear evolution in a low-dimensional Koopman latent space.

$$\begin{aligned}\mathcal{K}\phi_j &= \lambda_j\phi_j \\ \mathcal{K}^t f &= f \circ T^t\end{aligned}$$

