

Graph Rectifiability Summer Research with Lisa Naples

Daily Report

By

Charles Zhang

Summer 2021

Contents

1	May 25	3
1.1	Basic Set Theory	3
1.2	Functions and Limits	4

1 May 25

1.1 Basic Set Theory

Review and summary of some definitions and theorems:

Definition 1.1.1 (Countable) *An infinite set A is countable if its elements can be listed in the form x_1, x_2, \dots with every element appearing at a specific place in the list; otherwise, the set is uncountable*

Definition 1.1.2 (Open) $A \subset \mathbb{R}^n$ is open if, $\forall x \in A, \exists B(x, r) \subset A$ where $r > 0$.

Definition 1.1.3 (Closed) $A \subset \mathbb{R}^n$ is closed if, whenever $\{x_k\} \in A, x_k \rightarrow x \in \mathbb{R}^n$, then $x \in A$.

Definition 1.1.4 (Closure) \bar{A} is the intersection of all the closed sets containing a set A .

Definition 1.1.5 (Interior) $\text{int}(A)$ is the union of all open sets contained in A .

Definition 1.1.4 and 1.1.5 shows that The closure of A is thought of as the **smallest closed set** containing A , and the interior as the **largest open set** contained in A .

Definition 1.1.6 (Boundary) $\partial A = \bar{A} \setminus \text{int}(A)$

Theorem 1.1.1 $x \in \partial A \Leftrightarrow \forall r > 0, B(x, r) \cap A \neq \emptyset, B(x, r) \cap A^C \neq \emptyset$

Definition 1.1.7 (Dense) Set B is a dense in A if $A \subset \bar{B}$, that is, if there are points of B arbitrarily close to each point of A .

Definition 1.1.8 (Compact) A is compact if any collection of open sets that covers A has a finite subcollection which also covers A .

Theorem 1.1.2 A compact subset of \mathbb{R}^n is both closed and bounded.

Theorem 1.1.3 The intersection of any collection of compact sets is compact.

Definition 1.1.9 (Connected) $A \subset \mathbb{R}^n$ is connected if there not exists open sets U and V s.t. $A \subset U \cup V$ with disjoint and nonempty $A \cap U$ and $A \cap V$.

Definition 1.1.10 (Connected Component) Connected component of x is the largest connected subset of A containing a point x .

Definition 1.1.11 (Disconnect) The set A is totally disconnected if the connected component of each point consists of just that point.

The definition of disconnect also can be as: \exists open sets U and V s.t. $x \in U, y \in V$ and $A \subset U \cup V$.

Definition 1.1.12 (Borel Set) Borel Sets is the smallest collection of subsets of \mathbb{R}^n with the following properties:

1. Every open set and every closed set is a Borel set.
2. The union of every finite or countable collection of Borel sets is a Borel set, and the intersection of every finite or countable collection of Borel sets is a Borel set.

In short, Any set that can be constructed using a sequence of countable unions or intersections starting with the open sets or closed sets will certainly be Borel.

1.2 Functions and Limits

Definition 1.2.1 (Congruence) The transformation $S : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is congruence or isometry if it preserves distances i.e. if $|S(x) - S(y)| = |x - y|$ for $x, y \in \mathbb{R}^n$

Special cases include *translations*, which are of the form $S(x) = x + a$ and have the effect of shifting points parallel to the vector a , *rotations* which have a centre a such that $|S(x) - a| = |x - a|$ for all x (for convenience, we also regard the identity transformation given by $I(x) = x$ as a rotation) and *reflections*, which maps points to their mirror images in some $(n - 1)$ -dimensional plane. A congruence that may be achieved by a combination of a rotation and a translation, that is, does not involve reflection, is called a *rigid motion* or *direct congruence*. A transformation $S : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is a *similarity* of ratio or scale $c > 0$ if $|S(x) - S(y)| = c|x - y|$ for all x, y in \mathbb{R}^n . A similarity transforms sets into geometrically similar ones with all lengths multiplied by the factor c .

Definition 1.2.2 (Linear Transformation) A transformation $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is linear if $\forall x, y \in \mathbb{R}^n, T(x + y) = T(x) + T(y)$ and $T(\lambda x) = \lambda T(x), \lambda \in \mathbb{R}$

Such a linear transformation is *non-singular* if $T(x) = 0$ if and only if $x = 0$. If $S : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is of the form $S(x) = T(x) + a$, where T is a non-singular linear transformation and a is a vector in \mathbb{R}^n , then S is called an *affine transformation* or an *affinity*. An affinity may be thought of as a shearing transformation; its contracting or expanding effect need not be the same in every direction. However, if T is orthonormal, then S is a congruence, and if T is a scalar multiple of an orthonormal transformation, then T is a similarity.

Definition 1.2.3 (Hölder Function) A function $f : X \rightarrow Y$ is called a Hölder function of exponent α if

$$|f(x) - f(y)| \leq c|x - y|^\alpha \quad (x, y \in X)$$

for some constant $c \geq 0$.

Definition 1.2.4 (Lipschitz Function) The function f is called Lipschitz if

$$|f(x) - f(y)| \leq c|x - y| \quad (x, y \in X)$$

and bi-Lipschitz if

$$c_1|x - y| \leq |f(x) - f(y)| \leq c_2|x - y| \quad (x, y \in X)$$

for $0 < c_1 \leq c_2 < \infty$, in which case both f and $f^{-1} : f(X) \rightarrow X$ are Lipschitz functions.

Definition 1.2.5 (Lower Limit)

$$\liminf_{x \rightarrow 0} f(x) \equiv \lim_{r \rightarrow 0} (\inf \{f(x) : 0 < x < r\})$$

Note: $\inf \{f(x) : 0 < x < r\}$ is either $-\infty$ for all positive r or else increases as r decreases, $\liminf_{x \rightarrow 0} f(x)$ always exists.

Definition 1.2.6 (Upper Limit)

$$\overline{\lim}_{x \rightarrow 0} f(x) \equiv \lim_{r \rightarrow 0} (\sup\{f(x) : 0 < x < r\})$$

Note: The lower and upper limits exist as real numbers or $-\infty$ or ∞ for every function f and are indicative of the variation of f for x close to 0, shown in Figure 1.




Figure 1: The upper and lower limits of a function.

We write $f(x) \sim g(x)$ to mean that $f(x)/g(x) \rightarrow 1$ as $x \rightarrow 0$.

Theorem 1.2.1 (Lipschitz and Hölder functions are continuous)

Proof: for the function $f : X \rightarrow Y$ is a Hölder function of exponent α s.t. $|f(x) - f(y)| \leq c|x - y|^\alpha$ ($x, y \in X$) for some constant $c \geq 0$. Then, $\forall \epsilon > 0$, let $\delta = \frac{\epsilon}{c}$. Hence, $\forall x, y \in X, |x - y| < \delta \Rightarrow |x - y| < \frac{\epsilon}{c} \Rightarrow |f(x) - f(y)| \leq c|x - y| \leq c \cdot \frac{\epsilon}{c} = \epsilon \Rightarrow$ Hölder functions are continuous, and let exponent $\alpha = 1$, then we have Lipschitz functions are continuous.

Definition 1.2.7 (Homeomorphism) If $f : X \rightarrow Y$ is a continuous bijection with continuous inverse $f^{-1} : Y \rightarrow X$, then f is called a homeomorphism, and X and Y are termed homeomorphic sets.

Corollary 1.2.1 Congruences, similarities and affine transformations on \mathbb{R}^n are examples of homeomorphisms.

Definition 1.2.8 (Differentiable) If $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$, we say that f is differentiable at x and has derivative given by the linear mapping $f'(x) : \mathbb{R}^n \rightarrow \mathbb{R}^n$ if

$$\lim_{h \rightarrow 0} \frac{|f(x + h) - f(x) - f'(x)h|}{|h|} = 0.$$

Definition 1.2.9 (Pointwise Convergence) For a sequence of functions: $f_k : X \rightarrow Y$ where X and Y are subsets of Euclidean spaces. f_k converge pointwise to a function $f : X \rightarrow Y$ if $f_k(x) \rightarrow f(x)$ as $k \rightarrow \infty$.

Definition 1.2.10 (Uniform Convergence) For a sequence of functions: $f_k : X \rightarrow Y$ where X and Y are subsets of Euclidean spaces. f_k converge uniformly to a function $f : X \rightarrow Y$ if $\sup_{x \in X} |f_k(x) - f(x)| \rightarrow 0$ as $k \rightarrow \infty$.

Note: Uniform convergence is a stronger property than pointwise convergence i.e. Uniform convergence implies pointwise convergence, but not the other way around

Theorem 1.2.2 If the functions f_k are continuous and converge uniformly to f , then f is continuous.

proof: TODO

Theorem 1.2.3 (Logarithms) Apparently, $a^c = b^{c \log a / \log b}$