Section 1

The proposed approach processes physiological signals collected from participants. Each recording is segmented into fixed windows and normalized before analysis. Results are reported on a held out test split to avoid optimistic estimates. The proposed approach processes physiological signals collected from participants. Each recording is segmented into fixed windows and normalized before analysis. Results are reported on a held out test split to avoid optimistic estimates.

Section 2

The proposed approach processes physiological signals collected from participants. Each recording is segmented into fixed windows and normalized before analysis. Results are reported on a held out test split to avoid optimistic estimates. The proposed approach processes physiological signals collected from participants. Each recording is segmented into fixed windows and normalized before analysis. Results are reported on a held out test split to avoid optimistic estimates.

Section 3

The proposed approach processes physiological signals collected from participants. Each recording is segmented into fixed windows and normalized before analysis. Results are reported on a held out test split to avoid optimistic estimates. The proposed approach processes physiological signals collected from participants. Each recording is segmented into fixed windows and normalized before analysis. Results are reported on a held out test split to avoid optimistic estimates.

Section 4

The proposed approach processes physiological signals collected from participants. Each recording is segmented into fixed windows and normalized before analysis. Results are reported on a held out test split to avoid optimistic estimates. The proposed approach processes physiological signals collected from participants. Each recording is segmented into fixed windows and normalized before analysis. Results are reported on a held out test split to avoid optimistic estimates.