

# Molecular Dynamics Simulation of the D102A Variant of Chymotrypsin

David Li, Patrick Fleming

December 12, 2017

## 1 INTRODUCTION

Almost one third of all proteases can be classified as serine proteases [1]. These proteins are named for the nucleophilic Ser residue at the active site. Serine proteases fall into two broad categories based on their structure – chymotrypsin-like (trypsin-like) or subtilisin-like [2]. Chymotrypsin-like proteases are the most abundant in nature and can be found in eukaryotes, prokaryotes, archaea, and viruses. They are involved in many critical physiological processes, such as hemostasis, apoptosis, digestion, and reproduction [1].

In order to hydrolyze a peptide bond, these well-studied enzymes must overcome three major mechanistic barriers: (1) amide bonds are extremely stable due to electron donation from the amide nitrogen to the carbonyl; (2) water is a poor nucleophile and proteases always activate water via a general base; (3) and amines are poor leaving groups because proteases protonate the amine prior to expulsion [1].

To overcome these three reaction barriers, serine proteases contain a group of three residues called the catalytic triad that use hydrogen bonding to increase reaction favorability. In chymotrypsin, the triad is composed of serine, histidine, and aspartate, and is part of a larger hydrogen bonding network [1].

Mutagenesis experiments of catalytic triad residues show decreased catalytic activity; substitution of Ser195 or His57 with Ala effectively disables the triad [1]. For this project, we hypothesize that the Asp-His hydrogen bond restrains

the conformational flexibility of the His ring so that it can more strongly hydrogen bond to Ser, thus activating serine for a nucleophilic attack.

To test this, we carried out two all-atom molecular dynamics simulations of two chymotrypsin series variants: the wild type sequence, and with Asp-102 substituted for alanine (mutation D102A). This enabled us to determine rotamer distributions of the His side chain and the time-dependent frequencies of His-Ser bond formations.

## 2 METHODS

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

Nam dui ligula, fringilla a, euismod sodales, sollicitudin vel, wisi. Morbi auctor lorem non

justo. Nam lacus libero, pretium at, lobortis vitae, ultricies et, tellus. Donec aliquet, tortor sed accumsan bibendum, erat ligula aliquet magna, vitae ornare odio metus a mi. Morbi ac orci et nisl hendrerit mollis. Suspendisse ut massa. Cras nec ante. Pellentesque a nulla. Cum sociis natoque penatibus et magnis dis parturient montes, nascetur ridiculus mus. Aliquam tincidunt urna. Nulla ullamcorper vestibulum turpis. Pellentesque cursus luctus mauris.

Nulla malesuada porttitor diam. Donec felis erat, congue non, volutpat at, tincidunt tristique, libero. Vivamus viverra fermentum felis. Donec nonummy pellentesque ante. Phasellus adipiscing semper elit. Proin fermentum massa ac quam. Sed diam turpis, molestie vitae, placerat a, molestie nec, leo. Maecenas lacinia. Nam ipsum ligula, eleifend at, accumsan nec, suscipit a, ipsum. Morbi blandit ligula feugiat magna. Nunc eleifend consequat lorem. Sed lacinia nulla vitae enim. Pellentesque tincidunt purus vel magna. Integer non enim. Praesent euismod nunc eu purus. Donec bibendum quam in tellus. Nullam cursus pulvinar lectus. Donec et mi. Nam vulputate metus eu enim. Vestibulum pellentesque felis eu massa.

### 3 RESULTS




Figure 1: RMSD of wild-type (WT) and D102A simulations




Figure 2:




Figure 3:




Figure 4:




Figure 5:




Figure 6:

## 4 CONCLUSION

## ACKNOWLEDGEMENTS

## 5 REFERENCES

### REFERENCES

- [1] Hedstrom L. Serine protease mechanism and specificity. Chemical reviews. 2002 Dec 11;102(12):4501-24.
- [2] Madala PK, Tyndall JD, Nall T, Fairlie DP. Update 1 of: Proteases universally recognize beta strands in their active sites. Chemical reviews. 2011 Apr 8;110(6):PR1-31.

## A HYDROGEN BOND LIFETIME CODE

Minted portion to be added later.