

Chapter 14 - Problem Set 2

Calculus 3

Andry Paez

Section 5: The Chain Rule

3-7 (odd)

Use The Chain Rule to find $\frac{dz}{dt}$ or $\frac{dw}{dt}$.

3. $z = xy^3 - x^2y, \quad x = t^2 + 1, \quad y = t^2 - 1$

5. $z = \sin x \cos y, \quad x = \sqrt{t}, \quad y = 1/t$

7. $w = xe^{y/z}, \quad x = t^2, \quad y = 1 - t, \quad z = 1 + 2t$

11-15 (odd)

Use the Chain Rule to find $\frac{\partial z}{\partial s}$ and $\frac{\partial z}{\partial t}$

11. $z = (x - y)^5, \quad x = s^2t, \quad y = st^2$

13. $z = \ln(3x + 2y), \quad x = s \sin t, \quad y = t \cos s$

15. $z = (\sin \theta)/r, \quad r = st, \quad \theta = s^2 + t^2$

25-29 (odd)

Use the Chain Rule to find the indicated partial derivatives.

25. $z = x^4 + x^2y, \quad x = s + 2t - u, \quad y = stu^2;$

$\frac{\partial z}{\partial s}, \frac{\partial z}{\partial t}, \frac{\partial z}{\partial u}$ **when** $s = 4, t = e, u = 1$

27. $w = xy + yz + zx, \quad x = r \cos \theta, \quad y = r \sin \theta, \quad z = r\theta;$

$\frac{\partial w}{\partial r}, \frac{\partial w}{\partial \theta}$ **when** $r = 2, \theta = \pi/2$

29. $N = \frac{p+q}{p+r}, \quad p = u + vw, \quad q = v + uw, \quad r = w + uv$

$\frac{\partial N}{\partial u}, \frac{\partial N}{\partial v}, \frac{\partial N}{\partial w}$ **when** $u = 2, v = 3, w = 4$

Use Equation 5 to find $\frac{dy}{dx}$

31. $y \cos x = x^2 + y^2$

Use Equations 6 to find $\frac{\partial z}{\partial x}$ and $\frac{\partial z}{\partial y}$

35. $x^2 + 2y^2 + 3z^2 = 1$

Section 6: Directional Derivatives and the Gradient Vector

5, 7

Find the directional derivative of f at the given point in the direction indicated by the angle θ .

5.

7.

(a) Find the gradient of f

(b) Evaluate the gradient at the point P

(c) Find the rate of change of f at P in the direction of the vector \vec{u}

9.

13, 15

Find the directional derivative of the function at the given point in the direction of the vector \vec{v} .

13.

15.

21, 23

Find the directional derivative of the function at the point P in the direction of the point Q .

21.

23.

27, 29

Find the maximum rate of change of f at the given point and the direction in which it occurs.

27.

29.

37.

The temperature T in a metal ball is inversely proportional to the distance from the center of the ball, which we take to be the origin. The temperature at the point $(1, 2, 2)$ is 120°

(a) Find the rate of change of T at $(1, 2, 2)$ in the direction toward the point $(2, 1, 3)$.

(b) Show that at any point in the ball the direction of greatest increase in temperature is given by a vector that points toward the origin

47-51 (odd)

Find equations of (a) the tangent plane and (b) the normal line to the given surface at the specified point.

47.

49.

51.

Section 7: Maximum and Minimum Values

5-21 (odd)

Find the local maximum and minimum values and saddle point(s) of the function. You are encouraged to use a calculator or computer to graph the function with a domain and viewpoint that reveals all the important aspects of the function.

- 5.
- 7.
- 9.
- 11.
- 13.
- 15.
- 17.
- 19.
- 21.

33-39 (odd)

Find the absolute maximum and minimum values of f on the set D .

- 33.
- 35.
- 37.
- 39.

43

Find the shortest distance from the point $(2, 0, -3)$ to the plane $x + y + z = 1$.

45

Find the points on the cone $z^2 = x^2 + y^2$ that are closest to the point $(4, 2, 0)$.

47

Find three positive numbers whose sum is 100 and whose product is a maximum.

55

A cardboard box without a lid is to have a volume of $32,000 \text{ cm}^3$. Find the dimensions that minimize the amount of cardboard used.