

Chapter 14 - Problem Set 2

Calculus 3

Andry Paez

Section 5: The Chain Rule

3-7 (odd)

Use The Chain Rule to find $\frac{dz}{dt}$ or $\frac{dw}{dt}$.

3. $z = xy^3 - x^2y, \quad x = t^2 + 1, \quad y = t^2 - 1$

Solution

$$\begin{aligned}\frac{dz}{dt} &= \frac{dz}{dx} \left(\frac{dx}{dt} \right) + \frac{dz}{dy} \left(\frac{dy}{dt} \right) \\ \frac{dz}{dt} &= [y^3 - 2xy](2t) + [3xy^2 - x^2](2t) \\ \boxed{\frac{dz}{dt} &= 2t[y^3 - 2xy + 3xy^2 - x^2]}\end{aligned}$$

5. $z = \sin x \cos y, \quad x = \sqrt{t}, \quad y = 1/t$

Solution

$$\begin{aligned}\frac{dz}{dt} &= \frac{dz}{dx} \left(\frac{dx}{dt} \right) + \frac{dz}{dy} \left(\frac{dy}{dt} \right) \\ \frac{dz}{dt} &= [\cos x \cos y] \left(\frac{1}{2} t^{-\frac{1}{2}} \right) + [-\sin x \sin y] (-t^{-2}) \\ \boxed{\frac{dz}{dt} &= \frac{1}{2\sqrt{t}} \cos x \cos y + \frac{1}{t^2} \sin x \sin y}\end{aligned}$$

7. $w = xe^{y/z}, \quad x = t^2, \quad y = 1 - t, \quad z = 1 + 2t$

Solution

$$\begin{aligned}\frac{dw}{dt} &= \frac{dw}{dx} \left(\frac{dx}{dt} \right) + \frac{dw}{dy} \left(\frac{dy}{dt} \right) + \frac{dw}{dz} \left(\frac{dz}{dt} \right) \\ \frac{dw}{dt} &= [e^{y/z}] (2t) + \left[\frac{x}{z} e^{y/z} \right] (-1) + \left[-\frac{xy}{z^2} e^{y/z} \right] (2) \\ \boxed{\frac{dw}{dt} &= e^{y/z} \left[2t - \frac{x}{z} - \frac{2xy}{z^2} \right]}\end{aligned}$$

11-15 (odd)

Use the Chain Rule to find $\frac{\partial z}{\partial s}$ and $\frac{\partial z}{\partial t}$

11. $z = (x - y)^5, \quad x = s^2t, \quad y = st^2$

Solution

$$\begin{aligned}\frac{\partial z}{\partial s} &= \frac{\partial z}{\partial x} \left(\frac{\partial x}{\partial s} \right) + \frac{\partial z}{\partial y} \left(\frac{\partial y}{\partial s} \right) \\ \frac{\partial z}{\partial s} &= [5(x - y)^4] (2st) + [-5(x - y)^4] (t^2)\end{aligned}$$

$$\boxed{\frac{\partial z}{\partial s} = 5(x - y)^4 [2st - t^2]}$$

$$\begin{aligned}\frac{\partial z}{\partial t} &= \frac{\partial z}{\partial x} \left(\frac{\partial x}{\partial t} \right) + \frac{\partial z}{\partial y} \left(\frac{\partial y}{\partial t} \right) \\ \frac{\partial z}{\partial t} &= [5(x - y)^4] (s^2) + [-5(x - y)^4] (2st)\end{aligned}$$

$$\boxed{\frac{\partial z}{\partial t} = 5(x - y)^4 [s^2 - 2st]}$$

13. $z = \ln(3x + 2y), \quad x = s \sin t, \quad y = t \cos s$

Solution

$$\begin{aligned}\frac{\partial z}{\partial s} &= \frac{\partial z}{\partial x} \left(\frac{\partial x}{\partial s} \right) + \frac{\partial z}{\partial y} \left(\frac{\partial y}{\partial s} \right) \\ \frac{\partial z}{\partial s} &= \left[\frac{3}{3x + 2y} \right] (\sin t) + \left[\frac{2}{3x + 2y} \right] (-t \sin s)\end{aligned}$$

$$\boxed{\frac{\partial z}{\partial s} = \frac{3 \sin t - 2t \sin s}{3x + 2y}}$$

$$\begin{aligned}\frac{\partial z}{\partial t} &= \frac{\partial z}{\partial x} \left(\frac{\partial x}{\partial t} \right) + \frac{\partial z}{\partial y} \left(\frac{\partial y}{\partial t} \right) \\ \frac{\partial z}{\partial t} &= \left[\frac{3}{3x + 2y} \right] (s \cos t) + \left[\frac{2}{3x + 2y} \right] (\cos s)\end{aligned}$$

$$\boxed{\frac{\partial z}{\partial t} = \frac{3s \cos t + 2 \cos s}{3x + 2y}}$$

15. $z = (\sin \theta)/r, \quad r = st, \quad \theta = s^2 + t^2$

Solution

$$\frac{\partial z}{\partial s} = \frac{\partial z}{\partial r} \left(\frac{\partial r}{\partial s} \right) + \frac{\partial z}{\partial \theta} \left(\frac{\partial \theta}{\partial s} \right)$$

$$\frac{\partial z}{\partial s} = \left[-\frac{\sin \theta}{r^2} \right] (t) + \left[\frac{\cos \theta}{r} \right] (2s)$$

$$\boxed{\frac{\partial z}{\partial s} = -\frac{t \sin \theta}{r^2} + \frac{2s \cos \theta}{r}}$$

$$\frac{\partial z}{\partial t} = \frac{\partial z}{\partial r} \left(\frac{\partial r}{\partial t} \right) + \frac{\partial z}{\partial \theta} \left(\frac{\partial \theta}{\partial t} \right)$$

$$\frac{\partial z}{\partial t} = \left[-\frac{\sin \theta}{r^2} \right] (s) + \left[\frac{\cos \theta}{r} \right] (2t)$$

$$\boxed{\frac{\partial z}{\partial t} = -\frac{s \sin \theta}{r^2} + \frac{2t \cos \theta}{r}}$$

25-29 (odd)

Use the Chain Rule to find the indicated partial derivatives.

25. $z = x^4 + x^2y, \quad x = s + 2t - u, \quad y = stu^2;$

$\frac{\partial z}{\partial s}, \frac{\partial z}{\partial t}, \frac{\partial z}{\partial u}$ when $s = 4, t = 2, u = 1$

Solution

When $s = 4, t = 2$, and $u = 1 \Rightarrow x = 7, y = 8$

$$\frac{\partial z}{\partial s} = [4x^3 + 2xy] (1) + [x^2] (tu^2) = 4x^3 + 2xy + x^2tu^2$$

$$\frac{\partial z}{\partial s} = 4(7)^3 + 2(7)(8) + (7)^2(2)(1)^2$$

$$\boxed{\frac{\partial z}{\partial s} = 1582}$$

$$\frac{\partial z}{\partial t} = [4x^3 + 2xy] (2) + [x^2] (su^2) = 8x^3 + 4xy + x^2su^2$$

$$\frac{\partial z}{\partial t} = 8(7)^3 + 4(7)(8) + (7)^2(4)(1)^2$$

$$\boxed{\frac{\partial z}{\partial t} = 3164}$$

$$\frac{\partial z}{\partial u} = [4x^3 + 2xy] (-1) + [x^2] (2stu) = -4x^3 - 2xy + 2x^2stu$$

$$\frac{\partial z}{\partial u} = -4(7)^3 - 2(7)(8) + 2(7)^2(4)(2)(1)$$

$$\boxed{\frac{\partial z}{\partial u} = -700}$$

27. $w = xy + yz + zx, \quad x = r \cos \theta, \quad y = r \sin \theta, \quad z = r\theta;$

$$\frac{\partial w}{\partial r}, \frac{\partial w}{\partial \theta} \quad \text{when } r = 2, \theta = \pi/2$$

Solution

When $r = 2$ and $\theta = \pi/2 \Rightarrow x = 0, y = 2, z = \pi$

$$\frac{\partial w}{\partial r} = [y + z](\cos \theta) + [x + z](\sin \theta) + [x + y](\theta)$$

$$\frac{\partial w}{\partial r} = [2 + \pi](0) + [0 + \pi](1) + [0 + 2](\pi/2) = 0 + \pi + \pi$$

$$\boxed{\frac{\partial w}{\partial r} = 2\pi}$$

$$\frac{\partial w}{\partial \theta} = [y + z](-r \sin \theta) + [x + z](r \cos \theta) + [x + y](r)$$

$$\frac{\partial w}{\partial \theta} = [2 + \pi](-2 \cdot 1) + [0 + \pi](0) + [0 + 2](2) = -4 - 2\pi + 0 + 4$$

$$\boxed{\frac{\partial w}{\partial \theta} = -2\pi}$$

$$29. N = \frac{p+q}{p+r}, \quad p = u + vw, \quad q = v + uw, \quad r = w + uv$$

$$\frac{\partial N}{\partial u}, \frac{\partial N}{\partial v}, \frac{\partial N}{\partial w} \quad \text{when } u = 2, v = 3, w = 4$$

Solution

$$\text{When } u = 2, v = 3, \text{ and } w = 4 \Rightarrow p = 14, q = 11, r = 10$$

$$\frac{\partial N}{\partial u} = \left[\frac{(p+r) - (p+q)}{(p+r)^2} \right] (1) + \left[\frac{1}{p+r} \right] (w) + \left[-\frac{p+q}{(p+r)^2} \right] (v)$$

$$\frac{\partial N}{\partial u} = \frac{r-q}{(p+r)^2} + \frac{w}{p+r} - \frac{v(p+q)}{(p+r)^2}$$

$$\frac{\partial N}{\partial u} = \frac{10-11}{(14+10)^2} + \frac{4}{14+10} - \frac{3(14+11)}{(14+10)^2}$$

$$\frac{\partial N}{\partial u} = \frac{-1}{576} + \frac{4}{24} - \frac{75}{576} = \frac{-1+96-75}{576} = \frac{20}{576}$$

$$\boxed{\frac{\partial N}{\partial u} = \frac{5}{144}}$$

$$\frac{\partial N}{\partial v} = \left[\frac{r-q}{(p+r)^2} \right] (w) + \left[\frac{1}{p+r} \right] (1) + \left[-\frac{p+q}{(p+r)^2} \right] (u)$$

$$\frac{\partial N}{\partial v} = \frac{w(r-q)}{(p+r)^2} + \frac{1}{p+r} - \frac{u(p+q)}{(p+r)^2}$$

$$\frac{\partial N}{\partial v} = \frac{4(10-11)}{(14+10)^2} + \frac{1}{14+10} - \frac{2(14+11)}{(14+10)^2}$$

$$\frac{\partial N}{\partial v} = \frac{-4}{576} + \frac{1}{24} - \frac{50}{576} = \frac{-4+24-50}{576} = \frac{-30}{576}$$

$$\boxed{\frac{\partial N}{\partial v} = -\frac{5}{96}}$$

$$\frac{\partial N}{\partial w} = \left[\frac{r-q}{(p+r)^2} \right] (v) + \left[\frac{1}{p+r} \right] (u) + \left[-\frac{p+q}{(p+r)^2} \right] (1)$$

$$\frac{\partial N}{\partial w} = \frac{v(r-q)}{(p+r)^2} + \frac{u}{p+r} - \frac{p+q}{(p+r)^2}$$

$$\frac{\partial N}{\partial w} = \frac{3(10-11)}{(14+10)^2} + \frac{2}{(14+10)} - \frac{14+11}{(14+10)^2}$$

$$\frac{\partial N}{\partial w} = \frac{-3}{576} + \frac{2}{24} - \frac{25}{576} = \frac{-3+48-25}{576} = \frac{20}{576}$$

$$\boxed{\frac{\partial N}{\partial w} = \frac{5}{144}}$$

Use Equation 5 to find $\frac{dy}{dx}$

31. $y \cos x = x^2 + y^2$

Solution

$$\frac{dy}{dx} = -\frac{\frac{\partial F}{\partial x}}{\frac{\partial F}{\partial y}} = -\frac{F_x}{F_y}$$

Let $F = y \cos x - x^2 - y^2$

$$F_x = -y \sin x - 2x$$

$$F_y = \cos x - 2y$$

$$\frac{dy}{dx} = -\frac{F_x}{F_y} = -\frac{(-y \sin x - 2x)}{\cos x - 2y} = \frac{2x + y \sin x}{\cos x - 2y}$$

Use Equations 6 to find $\frac{\partial z}{\partial x}$ and $\frac{\partial z}{\partial y}$

35. $x^2 + 2y^2 + 3z^2 = 1$

Solution

$$\frac{\partial z}{\partial x} = -\frac{F_x}{F_z} \quad \frac{\partial z}{\partial y} = -\frac{F_y}{F_z}$$

Let $F = x^2 + 2y^2 + 3z^2 - 1$

$$F_x = 2x \quad F_y = 4y \quad F_z = 6z$$

$$\frac{\partial z}{\partial x} = -\frac{F_x}{F_z} = -\frac{2x}{6z} = -\frac{x}{3z}$$

$$\frac{\partial z}{\partial y} = -\frac{F_y}{F_z} = -\frac{4y}{6z} = -\frac{2y}{3z}$$

Section 6: Directional Derivatives and the Gradient Vector

5, 7

Find the directional derivative of f at the given point in the direction indicated by the angle θ .

5. $f(x, y) = xy^3 - x^2$, $(1, 2)$, $\theta = \pi/3$

7. $f(x, y) = \arctan(xy)$, $(2, -3)$, $\theta = 3\pi/4$

(a) Find the gradient of f

(b) Evaluate the gradient at the point P

(c) Find the rate of change of f at P in the direction of the vector \vec{u}

9. $f(x, y) = x/y$, $P(2, 1)$, $\vec{u} = \frac{3}{5}\hat{i} + \frac{4}{5}\hat{j}$

13, 15

Find the directional derivative of the function at the given point in the direction of the vector \vec{v} .

13. $f(x, y) = e^x \sin y$, $(0, \pi/3)$, $\vec{v} = \langle -6, 8 \rangle$

15. $g(s, t) = s\sqrt{t}$, $(2, 4)$, $\vec{v} = 2\hat{i} - \hat{j}$

21, 23

Find the directional derivative of the function at the point P in the direction of the point Q .

21. $f(x, y) = x^2y^2 - y^3$, $P(1, 2)$, $Q(-3, 5)$

23. $f(x, y) = \sqrt{xy}$, $P(2, 8)$, $Q(5, 4)$

27, 29

Find the maximum rate of change of f at the given point and the direction in which it occurs.

27. $f(x, y) = 5xy^2$, $(3, -2)$

29. $f(x, y) = \sin(xy)$, $(1, 0)$

37

The temperature T in a metal ball is inversely proportional to the distance from the center of the ball, which we take to be the origin. The temperature at the point $(1, 2, 2)$ is 120°

(a) Find the rate of change of T at $(1, 2, 2)$ in the direction toward the point $(2, 1, 3)$.

(b) Show that at any point in the ball the direction of greatest increase in temperature is given by a vector that points toward the origin

47-51 (odd)

Find equations of (a) the tangent plane and (b) the normal line to the given surface at the specified point.

47. $2(x-2)^2 + (y-1)^2 + (z-3)^2 = 10$, $(3, 3, 5)$

49. $xy^2z^3 = 8$, $(2, 2, 1)$

51. $x + y + z = e^{xyz}$, $(0, 0, 1)$

Section 7: Maximum and Minimum Values

5-21 (odd)

Find the local maximum and minimum values and saddle point(s) of the function. You are encouraged to use a calculator or computer to graph the function with a domain and viewpoint that reveals all the important aspects of the function.

5. $f(x, y) = x^2 + xy + y^2 + y$

7. $f(x, y) = 2x^2 - 8xy + y^4 - 4y^3$

9. $f(x, y) = (x - y)(1 - xy)$

11. $f(x, y) = y\sqrt{x} - y^2 - 2x + 7y$

13. $f(x, y) = x^3 - 3x + 3xy^2$

15. $f(x, y) = x^4 - 2x^2 + y^3 - 3y$

17. $f(x, y) = xy - x^2y - xy^2$

19. $f(x, y) = e^x \cos y$

21. $f(x, y) = y^2 - 2y \cos x, \quad -1 \leq x \leq 7$

33-39 (odd)

Find the absolute maximum and minimum values of f on the set D .

33. $f(x, y) = x^2 + y^2 - 2x,$

D is the closed triangular region with vertices $(2, 0)$, $(0, 2)$, and $(0, -2)$

35. $f(x, y) = x^2 + y^2 + x^2y + 4,$

$D = \{(x, y) \mid |x| \leq 1, |y| \leq 1\}$

37. $f(x, y) = x^2 + 2y^2 - 2x - 4y + 1,$

$D = \{(x, y) \mid 0 \leq x \leq 2, 0 \leq y \leq 3\}$

39. $f(x, y) = 2x^3 + y^4,$

$D = \{(x, y) \mid x^2 + y^2 \leq 1\}$

43

Find the shortest distance from the point $(2, 0, -3)$ to the plane $x + y + z = 1$.

45

Find the points on the cone $z^2 = x^2 + y^2$ that are closest to the point $(4, 2, 0)$.

47

Find three positive numbers whose sum is 100 and whose product is a maximum.

55

A cardboard box without a lid is to have a volume of $32,000 \text{ cm}^3$. Find the dimensions that minimize the amount of cardboard used.
