


## Backpropagation: a simple example

$$f(x, y, z) = (x + y)z$$

## Backpropagation: a simple example


$$f(x, y, z) = (x + y)z$$



## Backpropagation: a simple example

$$f(x, y, z) = (x + y)z$$

e.g.  $x = -2$ ,  $y = 5$ ,  $z = -4$



## Backpropagation: a simple example


$$f(x, y, z) = (x + y)z$$

e.g.  $x = -2$ ,  $y = 5$ ,  $z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$

Want:  $\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z}$



## Backpropagation: a simple example


$$f(x, y, z) = (x + y)z$$

e.g.  $x = -2, y = 5, z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$

Want:  $\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z}$



$$\frac{\partial f}{\partial f}$$

## Backpropagation: a simple example


$$f(x, y, z) = (x + y)z$$

e.g.  $x = -2, y = 5, z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$

Want:  $\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z}$



$$\frac{\partial f}{\partial f}$$

A pink arrow points from this box to the output  $f$  of the multiplication node in the computational graph above.

## Backpropagation: a simple example


$$f(x, y, z) = (x + y)z$$

e.g.  $x = -2, y = 5, z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$

Want:  $\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z}$



$$\frac{\partial f}{\partial z}$$

## Backpropagation: a simple example


$$f(x, y, z) = (x + y)z$$

e.g.  $x = -2, y = 5, z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$

Want:  $\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z}$



$$\frac{\partial f}{\partial z}$$



## Backpropagation: a simple example


$$f(x, y, z) = (x + y)z$$

e.g.  $x = -2, y = 5, z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$

Want:  $\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z}$



$$\frac{\partial f}{\partial q}$$

## Backpropagation: a simple example


$$f(x, y, z) = (x + y)z$$

e.g.  $x = -2, y = 5, z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$

Want:  $\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z}$



$$\frac{\partial f}{\partial q}$$

## Backpropagation: a simple example


$$f(x, y, z) = (x + y)z$$

e.g.  $x = -2, y = 5, z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$

Want:  $\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z}$



$$\frac{\partial f}{\partial y}$$

Chain rule:

$$\frac{\partial f}{\partial y} = \frac{\partial f}{\partial q} \frac{\partial q}{\partial y}$$

Upstream  
gradient

Local  
gradient

## Backpropagation: a simple example


$$f(x, y, z) = (x + y)z$$

e.g.  $x = -2, y = 5, z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$

Want:  $\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z}$



$$\frac{\partial f}{\partial y}$$

Chain rule:

$$\frac{\partial f}{\partial y} = \frac{\partial f}{\partial q} \frac{\partial q}{\partial y}$$

Upstream  
gradient

Local  
gradient

## Backpropagation: a simple example


$$f(x, y, z) = (x + y)z$$

e.g.  $x = -2, y = 5, z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$

Want:  $\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z}$



$$\frac{\partial f}{\partial x}$$

Chain rule:

$$\frac{\partial f}{\partial x} = \frac{\partial f}{\partial q} \frac{\partial q}{\partial x}$$

Upstream  
gradient

Local  
gradient

## Backpropagation: a simple example


$$f(x, y, z) = (x + y)z$$

e.g.  $x = -2$ ,  $y = 5$ ,  $z = -4$

$$q = x + y \quad \frac{\partial q}{\partial x} = 1, \frac{\partial q}{\partial y} = 1$$

$$f = qz \quad \frac{\partial f}{\partial q} = z, \frac{\partial f}{\partial z} = q$$

Want:  $\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z}$




$$\frac{\partial f}{\partial x}$$


Chain rule:

$$\frac{\partial f}{\partial x} = \frac{\partial f}{\partial q} \frac{\partial q}{\partial x}$$


Upstream  
gradient


Local  
gradient



















Another example:  $f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$




Another example:

$$f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$$



Another example:


$$f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$$



$f(x) = e^x$	$\rightarrow$	$\frac{df}{dx} = e^x$		$f(x) = \frac{1}{x}$	$\rightarrow$	$\frac{df}{dx} = -1/x^2$
$f_a(x) = ax$	$\rightarrow$	$\frac{df}{dx} = a$		$f_c(x) = c + x$	$\rightarrow$	$\frac{df}{dx} = 1$

Another example:

$$f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$$



$$f(x) = e^x$$

→

$$\frac{df}{dx} = e^x$$

$$f_a(x) = ax$$

→

$$\frac{df}{dx} = a$$

$$f(x) = \frac{1}{x}$$

→

$$\frac{df}{dx} = -1/x^2$$

$$f_c(x) = c + x$$


→

$$\frac{df}{dx} = 1$$



Another example:

$$f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$$



$$f(x) = e^x$$

→

$$\frac{df}{dx} = e^x$$

$$f_a(x) = ax$$

→

$$\frac{df}{dx} = a$$

$$f(x) = \frac{1}{x}$$

→

$$\frac{df}{dx} = -1/x^2$$

$$f_c(x) = a + x$$


Upstream gradient

→

$$\frac{df}{dx} = 1$$

Another example:


$$f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$$



$f(x) = e^x$	$\rightarrow$	$\frac{df}{dx} = e^x$		$f(x) = \frac{1}{x}$	$\rightarrow$	$\frac{df}{dx} = -1/x^2$
$f_a(x) = ax$	$\rightarrow$	$\frac{df}{dx} = a$		$f_c(x) = c + x$	$\rightarrow$	$\frac{df}{dx} = 1$

Another example:


$$f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$$



$f(x) = e^x$	$\rightarrow$	$\frac{df}{dx} = e^x$		$f(x) = \frac{1}{x}$	$\rightarrow$	$\frac{df}{dx} = -1/x^2$
$f_a(x) = ax$	$\rightarrow$	$\frac{df}{dx} = a$		<div style="border: 1px solid red; padding: 5px;"><math>f_c(x) = c + x</math></div>	$\rightarrow$	<div style="border: 1px solid red; padding: 5px;"><math>\frac{df}{dx} = 1</math></div>

Another example:

$$f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$$



$$f(x) = e^x$$

$\rightarrow$

$$\frac{df}{dx} = e^x$$

$$f_a(x) = ax$$

$\rightarrow$

$$\frac{df}{dx} = a$$

$$f(x) = \frac{1}{x}$$

$\rightarrow$

$$\frac{df}{dx} = -1/x^2$$


$$f_c(x) = c + x$$

$\rightarrow$

$$\frac{df}{dx} = 1$$

Another example:

$$f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$$



$$f(x) = e^x$$

→

$$\frac{df}{dx} = e^x$$

$$f_a(x) = ax$$

→

$$\frac{df}{dx} = a$$

$$f(x) = \frac{1}{x}$$

→

$$\frac{df}{dx} = -1/x^2$$


$$f_c(x) = c + x$$

→

$$\frac{df}{dx} = 1$$

Another example:

$$f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$$



$$f(x) = e^x \rightarrow \frac{df}{dx} = e^x$$


$$f_a(x) = ax \rightarrow \frac{df}{dx} = a$$

$$f(x) = \frac{1}{x} \rightarrow \frac{df}{dx} = -1/x^2$$

$$f_c(x) = c + x \rightarrow \frac{df}{dx} = 1$$

Another example:

$$f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$$



$$f(x) = e^x \quad \rightarrow \quad \frac{df}{dx} = e^x$$


$$f_a(x) = ax \quad \rightarrow \quad \frac{df}{dx} = a$$

$$f(x) = \frac{1}{x} \quad \rightarrow \quad \frac{df}{dx} = -1/x^2$$

$$f_c(x) = c + x \quad \rightarrow \quad \frac{df}{dx} = 1$$

Another example:

$$f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$$



$$f(x) = e^x$$

→

$$\frac{df}{dx} = e^x$$

$$f_a(x) = ax$$

→

$$\frac{df}{dx} = a$$

$$f(x) = \frac{1}{x}$$

→

$$\frac{df}{dx} = -1/x^2$$

$$f_c(x) = c + x$$


→

$$\frac{df}{dx} = 1$$



Another example:

$$f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$$



[upstream gradient] x [local gradient]

$$[0.2] \times [1] = 0.2$$

$$[0.2] \times [1] = 0.2 \text{ (both inputs!)}$$

$$f(x) = e^x$$

→

$$\frac{df}{dx} = e^x$$

$$f_a(x) = ax$$

→

$$\frac{df}{dx} = a$$

$$f(x) = \frac{1}{x}$$

→

$$\frac{df}{dx} = -1/x^2$$


$$f_c(x) = c + x$$

→

$$\frac{df}{dx} = 1$$

Another example:

$$f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$$



$$f(x) = e^x$$

→

$$\frac{df}{dx} = e^x$$

$$f_a(x) = ax$$

→

$$\frac{df}{dx} = a$$

$$f(x) = \frac{1}{x}$$

→

$$\frac{df}{dx} = -1/x^2$$


$$f_c(x) = c + x$$

→

$$\frac{df}{dx} = 1$$

Another example:

$$f(w, x) = \frac{1}{1 + e^{-(w_0x_0 + w_1x_1 + w_2)}}$$



[upstream gradient] x [local gradient]

$$x_0: [0.2] \times [2] = 0.4$$

$$w_0: [0.2] \times [-1] = -0.2$$

$$f(x) = e^x$$

→

$$\frac{df}{dx} = e^x$$

$$f_a(x) = ax$$

→

$$\frac{df}{dx} = a$$

$$f(x) = \frac{1}{x}$$

→

$$\frac{df}{dx} = -1/x^2$$

$$f_c(x) = c + x$$

→

$$\frac{df}{dx} = 1$$


$$f(w, x) = \frac{1}{1 + e^{-(w_0 x_0 + w_1 x_1 + w_2)}}$$

Computational graph representation may not be unique. Choose one where local gradients at each node can be easily expressed!

$$\sigma(x) = \frac{1}{1 + e^{-x}}$$

sigmoid function

$$\frac{d\sigma(x)}{dx} = \frac{e^{-x}}{(1 + e^{-x})^2} = \left( \frac{1 + e^{-x} - 1}{1 + e^{-x}} \right) \left( \frac{1}{1 + e^{-x}} \right) = (1 - \sigma(x)) \sigma(x)$$




$$f(w, x) = \frac{1}{1 + e^{-(w_0 x_0 + w_1 x_1 + w_2 x_2)}}$$

Computational graph representation may not be unique. Choose one where local gradients at each node can be easily expressed!

$$\sigma(x) = \frac{1}{1 + e^{-x}}$$

sigmoid function


$$\frac{d\sigma(x)}{dx} = \frac{e^{-x}}{(1 + e^{-x})^2} = \left( \frac{1 + e^{-x} - 1}{1 + e^{-x}} \right) \left( \frac{1}{1 + e^{-x}} \right) = (1 - \sigma(x)) \sigma(x)$$



[upstream gradient] x [local gradient]  
 $[1.00] \times [(1 - 0.73) (0.73)] = 0.2$

# Patterns in backward flow


**add** gate: gradient distributor



# Patterns in backward flow

**add** gate: gradient distributor


Q: What is a **max** gate?



# Patterns in backward flow

**add** gate: gradient distributor

**max** gate: gradient router






# Patterns in backward flow

**add** gate: gradient distributor

**max** gate: gradient router

Q: What is a **mul** gate?




# Patterns in backward flow


**add** gate: gradient distributor

**max** gate: gradient router

**mul** gate: gradient switcher




# Gradients add at branches



# Gradients for vectorized code


(x,y,z are  
now vectors)

This is now the  
**Jacobian matrix**  
(derivative of each  
element of z w.r.t. each  
element of x)



gradients

# Vectorized operations




# Vectorized operations

$$\frac{\partial L}{\partial x} = \boxed{\frac{\partial f}{\partial x}} \frac{\partial L}{\partial f}$$

Jacobian matrix

4096-d  
input vector



4096-d  
output vector


Q: what is the  
size of the  
Jacobian matrix?

# Vectorized operations

$$\frac{\partial L}{\partial x} = \boxed{\frac{\partial f}{\partial x}} \frac{\partial L}{\partial f}$$

Jacobian matrix

4096-d  
input vector



4096-d  
output vector

Q: what is the  
size of the  
Jacobian matrix?  
[4096 x 4096!]

A vectorized example:  $f(x, W) = ||W \cdot x||^2 = \sum_{i=1}^n (W \cdot x)_i^2$



A vectorized example:  $f(x, W) = ||W \cdot x||^2 = \sum_{i=1}^n (W \cdot x)_i^2$





Diagram illustrating the vectorization of the variables  $x$  and  $W$  in the function  $f(x, W)$ . Two gray arrows point downwards from  $x$  and  $W$  in the equation above to their respective domain specifications below.

$$\begin{matrix} \downarrow & \downarrow \\ \in \mathbb{R}^n & \in \mathbb{R}^{n \times n} \end{matrix}$$


A vectorized example:  $f(x, W) = ||W \cdot x||^2 = \sum_{i=1}^n (W \cdot x)_i^2$



A vectorized example:  $f(x, W) = ||W \cdot x||^2 = \sum_{i=1}^n (W \cdot x)_i^2$

$$\begin{bmatrix} 0.1 & 0.5 \\ -0.3 & 0.8 \end{bmatrix} \mathbf{W}$$


$$\begin{bmatrix} 0.2 \\ 0.4 \end{bmatrix} \mathbf{x}$$



$$q = W \cdot x = \begin{pmatrix} W_{1,1}x_1 + \cdots + W_{1,n}x_n \\ \vdots \\ W_{n,1}x_1 + \cdots + W_{n,n}x_n \end{pmatrix}$$


$$f(q) = ||q||^2 = q_1^2 + \cdots + q_n^2$$

A vectorized example:  $f(x, W) = ||W \cdot x||^2 = \sum_{i=1}^n (W \cdot x)_i^2$




$$q = W \cdot x = \begin{pmatrix} W_{1,1}x_1 + \cdots + W_{1,n}x_n \\ \vdots \\ W_{n,1}x_1 + \cdots + W_{n,n}x_n \end{pmatrix}$$
$$f(q) = ||q||^2 = q_1^2 + \cdots + q_n^2$$

A vectorized example:  $f(x, W) = ||W \cdot x||^2 = \sum_{i=1}^n (W \cdot x)_i^2$



$$q = W \cdot x = \begin{pmatrix} W_{1,1}x_1 + \cdots + W_{1,n}x_n \\ \vdots \\ W_{n,1}x_1 + \cdots + W_{n,n}x_n \end{pmatrix}$$
$$f(q) = ||q||^2 = q_1^2 + \cdots + q_n^2$$

A vectorized example:  $f(x, W) = ||W \cdot x||^2 = \sum_{i=1}^n (W \cdot x)_i^2$




$$q = W \cdot x = \begin{pmatrix} W_{1,1}x_1 + \cdots + W_{1,n}x_n \\ \vdots \\ W_{n,1}x_1 + \cdots + W_{n,n}x_n \end{pmatrix}$$

$$f(q) = ||q||^2 = q_1^2 + \cdots + q_n^2$$

$$\frac{\partial f}{\partial q_i} = 2q_i$$

$$\nabla_q f = 2q$$

A vectorized example:  $f(x, W) = ||W \cdot x||^2 = \sum_{i=1}^n (W \cdot x)_i^2$




$$q = W \cdot x = \begin{pmatrix} W_{1,1}x_1 + \cdots + W_{1,n}x_n \\ \vdots \\ W_{n,1}x_1 + \cdots + W_{n,n}x_n \end{pmatrix}$$

$$f(q) = ||q||^2 = q_1^2 + \cdots + q_n^2$$

$$\frac{\partial f}{\partial q_i} = 2q_i$$

$$\nabla_q f = 2q$$

A vectorized example:  $f(x, W) = ||W \cdot x||^2 = \sum_{i=1}^n (W \cdot x)_i^2$



$$q = W \cdot x = \begin{pmatrix} W_{1,1}x_1 + \cdots + W_{1,n}x_n \\ \vdots \\ W_{n,1}x_1 + \cdots + W_{n,n}x_n \end{pmatrix}$$


$$f(q) = ||q||^2 = q_1^2 + \cdots + q_n^2$$



A vectorized example:  $f(x, W) = ||W \cdot x||^2 = \sum_{i=1}^n (W \cdot x)_i^2$

$$\begin{bmatrix} 0.1 & 0.5 \\ -0.3 & 0.8 \end{bmatrix} W$$

$$\begin{bmatrix} 0.2 \\ 0.4 \end{bmatrix} x$$




$$\frac{\partial q_k}{\partial W_{i,j}} = \mathbf{1}_{k=i} x_j$$

$$\begin{aligned} \frac{\partial f}{\partial W_{i,j}} &= \sum_k \frac{\partial f}{\partial q_k} \frac{\partial q_k}{\partial W_{i,j}} \\ &= \sum_k (2q_k) (\mathbf{1}_{k=i} x_j) \\ &= 2q_i x_j \end{aligned}$$

$$q = W \cdot x = \begin{pmatrix} W_{1,1}x_1 + \cdots + W_{1,n}x_n \\ \vdots \\ W_{n,1}x_1 + \cdots + W_{n,n}x_n \end{pmatrix}$$

$$f(q) = ||q||^2 = q_1^2 + \cdots + q_n^2$$

A vectorized example:  $f(x, W) = ||W \cdot x||^2 = \sum_{i=1}^n (W \cdot x)_i^2$




$$\frac{\partial q_k}{\partial W_{i,j}} = \mathbf{1}_{k=i} x_j$$

$$\begin{aligned} \frac{\partial f}{\partial W_{i,j}} &= \sum_k \frac{\partial f}{\partial q_k} \frac{\partial q_k}{\partial W_{i,j}} \\ &= \sum_k (2q_k) (\mathbf{1}_{k=i} x_j) \\ &= 2q_i x_j \end{aligned}$$

$$q = W \cdot x = \begin{pmatrix} W_{1,1}x_1 + \cdots + W_{1,n}x_n \\ \vdots \\ W_{n,1}x_1 + \cdots + W_{n,n}x_n \end{pmatrix}$$

$$f(q) = ||q||^2 = q_1^2 + \cdots + q_n^2$$

A vectorized example:  $f(x, W) = ||W \cdot x||^2 = \sum_{i=1}^n (W \cdot x)_i^2$




$$\nabla_W f = 2q \cdot x^T$$

$$q = W \cdot x = \begin{pmatrix} W_{1,1}x_1 + \cdots + W_{1,n}x_n \\ \vdots \\ W_{n,1}x_1 + \cdots + W_{n,n}x_n \end{pmatrix}$$

$$f(q) = ||q||^2 = q_1^2 + \cdots + q_n^2$$

$$\begin{aligned} \frac{\partial q_k}{\partial W_{i,j}} &= \mathbf{1}_{k=i}x_j \\ \frac{\partial f}{\partial W_{i,j}} &= \sum_k \frac{\partial f}{\partial q_k} \frac{\partial q_k}{\partial W_{i,j}} \\ &= \sum_k (2q_k)(\mathbf{1}_{k=i}x_j) \\ &= 2q_i x_j \end{aligned}$$

A vectorized example:  $f(x, W) = ||W \cdot x||^2 = \sum_{i=1}^n (W \cdot x)_i^2$




$$q = W \cdot x = \begin{pmatrix} W_{1,1}x_1 + \cdots + W_{1,n}x_n \\ \vdots \\ W_{n,1}x_1 + \cdots + W_{n,n}x_n \end{pmatrix}$$

$$f(q) = ||q||^2 = q_1^2 + \cdots + q_n^2$$

$$\begin{aligned} \frac{\partial q_k}{\partial W_{i,j}} &= \mathbf{1}_{k=i} x_j \\ \frac{\partial f}{\partial W_{i,j}} &= \sum_k \frac{\partial f}{\partial q_k} \frac{\partial q_k}{\partial W_{i,j}} \\ &= \sum_k (2q_k) (\mathbf{1}_{k=i} x_j) \\ &= 2q_i x_j \end{aligned}$$

A vectorized example:  $f(x, W) = ||W \cdot x||^2 = \sum_{i=1}^n (W \cdot x)_i^2$




$$\frac{\partial q_k}{\partial x_i} = W_{k,i}$$

$$q = W \cdot x = \begin{pmatrix} W_{1,1}x_1 + \cdots + W_{1,n}x_n \\ \vdots \\ W_{n,1}x_1 + \cdots + W_{n,n}x_n \end{pmatrix}$$

$$f(q) = ||q||^2 = q_1^2 + \cdots + q_n^2$$

A vectorized example:  $f(x, W) = ||W \cdot x||^2 = \sum_{i=1}^n (W \cdot x)_i^2$




$$q = W \cdot x = \begin{pmatrix} W_{1,1}x_1 + \cdots + W_{1,n}x_n \\ \vdots \\ W_{n,1}x_1 + \cdots + W_{n,n}x_n \end{pmatrix}$$

$$f(q) = ||q||^2 = q_1^2 + \cdots + q_n^2$$

$$\begin{aligned} \frac{\partial q_k}{\partial x_i} &= W_{k,i} \\ \frac{\partial f}{\partial x_i} &= \sum_k \frac{\partial f}{\partial q_k} \frac{\partial q_k}{\partial x_i} \\ &= \sum_k 2q_k W_{k,i} \end{aligned}$$

A vectorized example:  $f(x, W) = ||W \cdot x||^2 = \sum_{i=1}^n (W \cdot x)_i^2$



$$\nabla_x f = 2W^T \cdot q$$

$$q = W \cdot x = \begin{pmatrix} W_{1,1}x_1 + \cdots + W_{1,n}x_n \\ \vdots \\ W_{n,1}x_1 + \cdots + W_{n,n}x_n \end{pmatrix}$$

$$f(q) = ||q||^2 = q_1^2 + \cdots + q_n^2$$

$$\begin{aligned} \frac{\partial q_k}{\partial x_i} &= W_{k,i} \\ \frac{\partial f}{\partial x_i} &= \sum_k \frac{\partial f}{\partial q_k} \frac{\partial q_k}{\partial x_i} \\ &= \sum_k 2q_k W_{k,i} \end{aligned}$$