

|                                | Yes       | No                                  |
|--------------------------------|-----------|-------------------------------------|
| Diagnosis Discrepancy          |           | <input checked="" type="checkbox"/> |
| Primary Tumor Site Discrepancy |           | <input checked="" type="checkbox"/> |
| IIHAA Discrepancy              |           | <input checked="" type="checkbox"/> |
| Prior Malignancy History       |           | <input checked="" type="checkbox"/> |
| Dual/Synchronous Primary Tumor |           | <input checked="" type="checkbox"/> |
| Case is (circle):              | QUALIFIED | DISQUALIFIED                        |
| Reviewer Initials              | RB        | Date Reviewed: 10/18/11             |

UUID: 25667964-D12A-43A9-885E-DD8FDB694D74  
TCGA-ET-A3D0-01A-PR Redacted



SPECIMEN TAKEN:

FINAL DIAGNOSIS -----

TOTAL THYROIDECTOMY:

SPECIMEN TYPE: Total thyroidectomy

TUMOR SITE: Right lobe (main mass) and left lobe

HISTOLOGIC TYPE: Papillary carcinoma

1CD-0-3

TUMOR SIZE: Greatest dimension: 2.5 cm

Carcinoma, papillary, thyroid

FOCALITY: Multifocal

8260/3

Site: thyroid, nos C13.9

LYMPH NODES: None submitted

EXTENT OF INVASION

PRIMARY TUMOR:

pT2: Tumor >2cm but <4 cm, limited to thyroid

hw  
10/24/11

REGIONAL LYMPH NODES:

pNX: Cannot be assessed

DISTANT METASTASIS:

pMX: Cannot be assessed

MARGINS: Margins are uninvolved

Distance of invasive carcinoma from nearest margin: 1 mm (soft tissue margin)

VENOUS/LYMPHATIC INVASION:

Not identified

ADDITIONAL PATHOLOGIC FINDINGS:

Lymphocytic thyroiditis

Note: The largest focus of tumor (2.5 cm) involves the right lobe. A separate focus of papillary carcinoma (0.4 cm) is located in the left lobe. This case was shown at the