


ILD-0-3
 Carcinoma, lobular
 infiltration 8520/3
 Site: @ Breast, NOS
 JW 4/2/13 C50.9

UUID:F2409024-D848-4B02-A371-92BEA1105EC4
 TCGA-AR-A5QP-01A-PR

Redacted



A. Breast, left, lumpectomy: Infiltrating lobular carcinoma, Nottingham grade I (of III) [tubules 3/3, nuclei 1/3, mitoses 1/3; Nottingham score 5/9], forming a 2.2 x 2.1 x 1.6 cm mass. Angiolymphatic invasion is absent. The non-neoplastic breast parenchyma shows nonproliferative fibrocystic changes. Biopsy site changes present. All surgical resection margins are negative for tumor (minimum tumor free margin, 0.4 cm, deep margin, and 0.6 cm, anterior margin).

B. Lymph node, left, sentinel biopsy: One (of 2) axillary sentinel lymph nodes is positive with multiple (5) metastases, the largest measuring 3.5 mm in greatest dimension. Extranodal extension is focally present.

With available surgical material [AJCCpT2N1(sn)], (7th edition, 2010).

Seen in consultation with

| Criteria | Yes | No |
|--------------------------------|------------------------|-------------------------------------|
| Diagnosis Discrepancy | | <input checked="" type="checkbox"/> |
| Primary Tumor Site Discrepancy | | <input checked="" type="checkbox"/> |
| HiPAA Discrepancy | | <input checked="" type="checkbox"/> |
| Prior Malignancy History | | <input checked="" type="checkbox"/> |
| Dual/Synchronous Primary Noted | | <input checked="" type="checkbox"/> |
| Case is (circle): | QUALIFIED | DISQUALIFIED |
| Reviewer Initials | Date Reviewed: 1/28/13 | |