

Redacted



1CD-0-3

carcinoma, infiltrating duct, NOS 8500/3
 Site breast, NOS C50.9

Breast, left, modified radical mastectomy: Infiltrating ductal carcinoma, Nottingham grade II (of III), forming a mass (2.3 x 2.1 x 1.7 cm) located in the superior/central aspect of the breast [AJCC pT2]. Ductal carcinoma in situ, intermediate nuclear grade, comprises approximately 5-25% of the tumor volume. Angiolymphatic invasion is present. The non-neoplastic breast parenchyma shows nonproliferative fibrocystic changes. The tumor does not involve the nipple, overlying skin, or underlying chest wall. All surgical resection margins, after re-excision of the anterior/superior margin, are negative for tumor (minimum tumor free margin, 0.7 cm, anterior/superior margin). The deep margin is negative for tumor (free by 0.8 cm). Multiple (22) left mid/low axillary lymph nodes are negative for metastatic carcinoma.

Lymph nodes, left high axillary, excision: Multiple (3) left high axillary lymph nodes are negative for tumor.

Lymph node, left axillary sentinel, excision: A single left axillary sentinel lymph node is positive for metastatic adenocarcinoma with one metastasis measuring 2.5 x 2.5 x 1.3 cm [AJCC pN1]. Extranodal extension is not present. Blue dye is not identified.

Criteria	Yes	No
Diagnosis Discrepancy		X
Primary Tumor Site Discrepancy		X
HIPAA Discrepancy		X
Prior Malignancy History		X
Dual/Synchronous Primary Note		
Case Is (circle):	QUALIFIED	DISQUALIFIED
Reviewer Initials	9/1/10	