

UUID:AF3A1FB4-1FC6-4CB7-93EF-0E3E2FEE3E8B
TCGA-VS-A9UQ-01A-PR

Redacted



(MM/DD/YYYY)

ICD O-3
Adenocarcinoma, mucinous
endocervical type 848213
Cervix NOS C53.9

9/3/14

PATHOLOGY REPORT:

PRIMARY SITE: Cervix

- 1-) Previous surgical scar:
 - Cicatricial fibrosis of the dermis.
- 3-) Right iliac obturator lymph nodes:
 - Free of neoplasia (0/2).
- 4-) Right obturator lymphadenectomy:
 - Lymph nodes free of neoplasia (0/10).
- 5-) Left iliac obturator lymphadenectomy:
 - Lymph nodes free of neoplasia (0/10).
- 6-) Uterus + parametria + Fallopian tubes + vaginal cuff:
 - Moderately differentiated mucinous adenocarcinoma of endocervical type, measuring 4.0cm of maximum extension and 1.5cm of maximum depth of invasion.
 - The infiltrative tumor is present in the anterior half of the uterine cervix.
 - Presence of foci of adenocarcinoma in situ in the posterior wall of the cervix.
 - The tumor infiltrates almost all of the anterior cervical stromal thickness, being less than 1mm distant from the circumferential anterior surgical margin.
 - Isthmus, uterine corpus and vagina free of neoplasia.
 - No indubitable focus of angiolymphatic and perineural infiltration was observed.
 - Vaginal margin free of neoplasia.
 - Parametria free of neoplasia.
 - Leiomyomas in submucosal, subserosal and intramural topographies.
 - Fallopian tubes free of neoplasia.

Criteria	Yes	No
Diagnosis Discrepancy		✓
Primary Tumor Site Discrepancy		✓
HIPAA Discrepancy		✓
Prior Malignancy History		✓
Dual/Synchronous Primary Notes		✓
Case is (circle):	QUALIFIED	DISQUALIFIED
Reviewer Initials	MA	Date Reviewed: 12/7/13