

General description of catalogues:

Aperture: Kron convolved

Bands: all NIRCам and HST available

Selection criteria: $\text{SN}(\text{F444W}) \geq 3$ and $3 \leq z_{\text{Eazy}} \leq 9$

N_{total} = number of galaxies that satisfy selection criteria

N_{final} = number of galaxies that were successfully fit and that have $3 \leq z_{\text{Prospector}} \leq 9$.

The columns are explained below, in every case “elo” and “ehi” are the lower and upper errors, they were estimated from the 16th, 50th and 84th percentile and represent q50-q16 and q84-q50, respectively. The minus in the lower error bars is just a convention.

Name	Description
ID	NIRCам identifier from the photometric catalogue used
Nphot(HST+JWST)	total number of photometric bands used in the fit
red_chi2(HST+JWST)	reduced χ^2 using all the photometric points
Nphot(JWST)	number of NIRCам photometric bands used in the fit
red_chi2(JWST)	reduced χ^2 using only the NIRCам bands
SN(F277W), SN(F356W), SN(F444W)	the signal-to-noise in F277W, F356W, and F444W
z	redshift
log(Mstar)	logarithm of stellar mass in units of solar masses
SFR5, SFR10, SFR50, SFR100, SFR2000	star formation rates averaged over the past 5, 10, 50, 100, and 200 Myr (they are <u>NOT</u> logged, and are in units of solar masses per year)
t50, t90	time of half-mass (and 90%) assembly in Myr
log(nion)	logarithm of flux of ionising photons in units of s^{-1}
log(xion)	logarithm of the ionising photon production efficiency in units of Hz erg^{-1}
M1500obs, M1500int	observed and intrinsic UV magnitude (AB mag)
dust2	dust2 parameter from Conroy+2009
log(U)	logarithm of ionising parameter
log(Z)	logarithm of metallicity

Prospector settings and priors:

- **IMF:** Chabrier
- **redshift:** clipped normal using EAZY photometric redshifts (thanks to Kevin Hainline) as mean plus errors
- **log(stellar mass):** TopHat between 6 and 12
- **SFH:** continuity with 8 bins, student t_width = 0.3
- **log(stellar metallicity):** clipped normal between -2 and 0, with mean -1.5
- **log(gas metallicity), log(U):** free
- **dust attenuation law:** Conroy+2009
- We include IGM scaling, nebular emission, and impose an error floor of 5% in the photometry

GOODS-S

Photometry catalogue: hlsp_jades_jwst_nircam_goodss_photometry_v0.9.3_catalog

$N_{\text{total}} = 37272$

$N_{\text{final}} = 35442$

GOODS-N

Photometry catalogue: hlsp_jades_jwst_nircam_goodsn_photometry_v0.9.1_catalog

$N_{\text{total}} = 24683$

$N_{\text{final}} = 22421$