

Algebra

Textbook: Real Analysis

FRED YUAN

Fall 2024

Homework of our real variables class. Textbook by Folland. Professor: Shukun Wu

Last updated on Monday 9th September, 2024.

1 Week 1

Exercise 1.1. Prove demorgan's law: $(\cup_{a \in A} E_a)^c = \cap_{a \in A} E_a^c$

Proof. Let $x \in \cup_{a \in A} E_a$. This means x must be in some $E_{a_0}, E_{a_1}, \dots, E_{a_n}$. Then $x^c \in E_{a_0}^c, E_{a_1}^c, \dots, E_{a_n}^c$ Where $E_{a_n} \in \cap_{a \in A} E_a$

By definition of intersection, $x \in E_{a_0}^c \cap E_{a_1}^c \cap \dots E_{a_n}^c$, $x = \cap_{a \in A} E_a^c$ □

Exercise 1.2. Let X be a finite set. Prove that $|2^X| = 2^{|X|}$

Proof. Intuitively, we can mark each element as in or out Proof by induction: $X =$

1. Let $X = \{1\}$

$$P(X) = \{\emptyset, \{1\}\} = 2$$

$$\text{We WTS: } |2^{X+1}| = 2^{|X+1|}$$

$$2^{|X+1|} = P(X) \cup X_{n+1} = |P(X)| \cup X_{n+1}. \text{ You have two choices, include } X_{n+1} \text{ or not, so}$$

$$\implies |P(X)| \cdot 2 = |2^{X+1}|.$$

□

Exercise 1.3. Let $f: X \rightarrow Y$ be a mapping. Prove that $f^{-1}(\cup_{a \in A} E_a) = \cup_{a \in A} f^{-1} E_a$

Proof. 1. Let $x \in f^{-1} \cup_{a \in A} E_a \implies y \in \cup_{a \in A} E_a$. Then for some a_0 , $y \in E_{a_0}$ Where $E_{a_0} \subseteq E_a$ for all $a \in A \implies x \in f^{-1} E_{a_0} \subseteq \cup_{a \in A} f^{-1}(E_a)$

2. The proof goes the other way as well. (It's the same) Let $x \in \cup_{a \in A} f^{-1} E_a \implies x \in f^{-1}(E_{a_0}) \implies y \in E_{a_0} \subseteq \cup_{a \in A} E_a \implies x \subseteq f^{-1}(\cup_{a \in A} E_a)$ □

Exercise 1.4. Show that there does not exist a linear order of " \leq " of \mathbb{C} with the following additional assumptions:

1. For any 3 numbers a, b, c , $a \leq b$ implies $a + c \leq b + c$.
2. For any two numbers a, b , $a \geq 0$ and $b \geq 0$ implies $ab \geq 0$
3. $0 < 1$

Proof. we must define an ordering for i . Either $i \geq 0$ or $i \leq 0$. However, neither of these cases make sense.

If $i \geq 0$, square both sides to get a contradiction.

If $i \leq 0$, ⁴ both sides to get a contradiction.

For below, assume $i \leq 0$

1. If $i \geq 0$, then $a = 0$, $b = i$, $c = 0$. We have $0 \leq b$ but after squaring both sides, we get $0 \leq -1$. Contradiction.

If $i \leq 0$, $a = i$, $b = 0$, $c = 0$. Starting with $i \leq 0$, if we ⁴ both sides, we end up with $1 \geq 0$. Contradiction.

2. $a = 0a + 1i$, $b = 0a + ia$. $ab = -1$

3. If $0 < 1$, then $0a + 2i < 1a + 0i$. However, $(0a + 2i)^4 = 16 > 1^4 = 1$

□

Exercise 1.5. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be an increasing function (this is, $f(x_1) \geq f(x_2)$ if $x_1 \geq x_2$). Let $E := \{x \in \mathbb{R} : f(x) \text{ is not continuous in } x\}$. Prove that E is countable

Proof. Observation: Because f is increasing, let's say we start at arbitrary $-n$. Every $f(x_2), f(x_3), \dots \geq -n$. Let's say $f(x_2) = -n + \epsilon$. And since E is not continuous, this means $f(x)$ must be jumping.

Observation: Since this sequence is increasing, if we can show that there forms either a bijection or injection onto \mathbb{Q} , then E is countable.

If ϵ is a rational number, then we are already done.

So if $\epsilon = \pi$ or some irrational value, we need to find some $\frac{p}{q} < \epsilon$ so that we can form a bijection. Because \mathbb{Q} is dense in \mathbb{R} , we know there always exists a $0 < \frac{p}{q} < \epsilon$.

Thus, we can map E to \mathbb{Q} . Since $|E| \leq |\mathbb{Q}|$, E is countable.

□

Exercise 1.6. Suppose $E \subset \mathbb{R}$ is a countable set. Prove that there exists $r \in \mathbb{R}$ such that $E \cap (r + E) = \emptyset$. Here $r + E := \{r + x : x \in E\}$

Proof. Since E is countable, $|E| \leq |\mathbb{Q}|$, meaning E can be listed. $E := \{E_0, E_1, \dots, E_n\}$. WLOG, assume E is increasing and sorted.

Now if we let $x \in E$, the most concerning element is $x = E_0$. Let $r = E_n + 2|E_0|$.

We have two cases:
$$\begin{cases} E_0 > 0 : r + x = E_n + 2E_0 + x > E_n \\ E_0 < 0 : E_n + 2|E_0| + x > E_n \end{cases}$$

Therefore, if we let $r := E_n + 2|E_0|$, all elements $r + E$ will be $>$ E . Thus, $E \cap (r + E) = \emptyset$

□

2 Week 2

Exercise 2.1. Show that a countable union of countable sets is still countable

Proof. Suppose we have some countable sets, $S_0, S_1, S_2, \dots, S_n$ and $S := \bigcup_{i \in \mathbb{N}} S_i$. Since each of these sets are countable, we can list them all out.

$S_0 = \{S_{0_0}, S_{0_1}, \dots, S_{0_n}\}$, $S_1 = \{S_{1_0}, S_{1_1}, \dots, S_{1_n}\}, \dots$. We can re-arrange these sets in a $n \times n$ grid similar to the rationals. Then we can find a mapping from $|S|$ to $|\mathbb{Q}|$ just like cantor's snake.

i.e. There is a mapping $f: |\mathbb{N} \times \mathbb{N}| \rightarrow |\mathbb{N}|$ that lists everything in the grid. Therefore, S is countable. \square

Exercise 2.2. Show that \mathbb{Q} is a dense subset of \mathbb{R} under the euclidian metric.

Proof. To be dense in \mathbb{R} , we must show that for every element $r \in \mathbb{R}$, $\exists q \in \mathbb{Q}$ such that $d(q, r) < \epsilon$ given arbitrary ϵ .

Take $q := \frac{m}{n}$ and suppose r is irrational.

WTF: q such that $|r - \frac{m}{n}| < \epsilon$. Assume $r > q$ so $r - \frac{m}{n} < \epsilon$

By the archemedian property, we know there exists a $n > N$ s.t. $\frac{1}{n} < \epsilon$.
Also, for any $r \in \mathbb{R}$, we can find a m such that $m \leq nr < m + 1$.

$$= 0 \leq nr - m < 1 \rightarrow 0 \leq r - \frac{m}{n} < \frac{1}{n}$$

Finally, since $\epsilon > \frac{1}{n}$, we arrive at $|r - \frac{m}{n}| < \frac{1}{n} < \epsilon$. \square