



Overview
(/study/ap
aa-
hl/sid-
134-
cid-
761926/o

Teacher view



(https://intercom.help/kognity)

**Index**

- The big picture
- The 3D coordinate system
- The right pyramid
- The cone
- The sphere and hemisphere
- Combined solids
- Angles between lines
- Angle between a line and a plane
- Checklist
- Investigation



Table of
contents



Notebook

3. Geometry and trigonometry / 3.1 Three-dimensional space



Glossary



Reading
assistance

The big picture

Geometric figures and their transformations are widely used in design and architecture. Apart from engineering calculations of buildings, architects use geometry to define the spatial form of a building, to create harmonious forms and also to create environmentally conscious buildings. The Petronas Twin Towers in Kuala Lumpur, Malaysia, are among the many skyscrapers built in recent history. The design of the towers starts with a square where the vertices of the square represent unity, harmony, stability and rationality. This square is rotated through 90° and small circular infills complete the floor plate of the towers, as seen in the photograph. The next stage of the design moves from 2D to 3D using the base plate, which changes in size as the tower is built upwards. From a distance, the towers give the illusion of cylinders at the bottom and cones or pyramids on the upper floors.

How many square metres of glass do you think are used in these towers? What is the total volume of the structure, and how many air conditioning units are needed to cool the building in the hot tropical weather?



Student
view



To help answer these types of question, you will be learning about 3D shapes and forms, and their relation to 2D objects.

Overview

(/study/ap

aa-

hl/sid-

134-


cid-

761926/o



The Petronas Twin Towers, Kuala Lumpur, Malaysia

Credit: Nikada Getty Images



Concept



Student
view

3D shapes such as the right pyramid, cone and sphere are widely used in design and architecture. Identifying elements such as faces, edges and vertices helps you to calculate surface area and volume which



Overview
(/study/ap

aa-
hl/sid-
134-
cid-
761926/o


define the space occupied, and you can use modelling to create new designs.

3. Geometry and trigonometry / 3.1 Three-dimensional space

The 3D coordinate system

Plotting points in 3D

The Cartesian coordinate system consists of three perpendicular planes intersecting at a point called ‘the origin’. Similar to the 2D coordinate plane, each point is defined by its perpendicular distance from each plane, as shown in the diagram below. By convention, you use (x, y, z) , always in the same order, to represent the coordinates of a point in 3D.



Student
view



Overview
(/study/app)

aa-
hl/sid-
134-
cid-
761926/o

More information

The image depicts a 3D Cartesian coordinate system with three perpendicular planes representing the x, y, and z axes. These planes intersect at the origin, marked as O. The x-axis extends horizontally, the y-axis extends depth-wise, and the z-axis is vertical. Each axis is labeled accordingly. A point labeled (x, y, z) is marked within the space, located at a specific position on each axis, forming a rectangular cuboid shape in the 3D space with the origin as one vertex. The edges of this cuboid run parallel to each axis, illustrating the point's coordinates in the 3D space. Light blue planes emphasize the 3D nature and spatial division by the axes.

[Generated by AI]

As it is a perpendicular system, you can think of the coordinates of a point as a vertex of a cuboid, where one of the vertices is at the origin, and three of the edges are on each axis. In the diagram below, point A (1,2,1) is the vertex of a cuboid with edges of lengths 1, 2 and 1 drawn on the x, y and z axes, respectively.

Section

Student... (0/0)

Feedback

Print (/study/app/math-aa-hl/sid-

Assign


134-cid-761926/book/the-big-picture-id-25407/print/)



Student
view



Overview
 (/study/ap
 aa-
 hl/sid-
 134-
 cid-
 761926/o



More information

The image depicts a 3D coordinate system with x, y, and z axes intersecting at the origin labeled as "O." A cuboid is positioned within this system. In the upper diagram, point A (1,2,1) is part of a cuboid, with one vertex at the origin and the others extending along the axes. The cuboid is outlined with dashed lines along the x and y axes and solid lines along the z axis. The edges of the cuboid have lengths of 1, 2, and 1 along the x, y, and z axes, respectively. The lower diagram shows the same structure rotated to highlight the y-axis perspective. The point A is consistently marked at (1,2,1) with a bold dot, and the cuboid edges are either dashed or solid, confirming their measurements on respective axes.

[Generated by AI]



Student
view




Overview
 (/study/ap
 aa-
 hl/sid-
 134-
 cid-
 761926/o

✓ Important

You use (x, y, z) to represent the coordinates of a point in 3D space.

Example 1



More information

The image depicts a cuboid drawn in a 3D coordinate system. The cuboid is displayed with vertices labeled as A, B, C, D, E, F, G, and H. The coordinate axes are labeled x, y, and z. The axis lines have scale markings: the x-axis extends from 0 to 3, the y-axis from 0 to 3, and the z-axis from 0 to 3, each marked at unit intervals. The vertices are connected by dotted lines, forming a rectangular prism. The cuboid sits on the base formed by points A, B, E, and F with its top formed by points C, D, G, and H. Each axis appears to define the space in which the cuboid is centered, with the point O at the origin.



Student view

[Generated by AI]



Overview
(/study/app/math-aa-hl/sid-134-cid-761926)

aa-
hl/sid-
134-
cid-
761926/o


The figure above shows a cuboid drawn in the 3D coordinate system.

a) Write down the letters with coordinates $(0,0,0)$ and $(2,0,0)$ respectively.

b) Write down the coordinates of each of the vertices of the cuboid.

a) F $(0,0,0)$ and A $(2,0,0)$.

b) Using (x, y, z) as (length, width, height) gives B $(2,1,0)$, C $(2,1,3)$, D $(2,0,3)$, E $(0,1,0)$, G $(0,0,3)$ and H $(0,1,3)$.



Student
view



Midpoint of a line segment

Overview


- (/study/ap aa-hl/sid-134-cid-761926/o) Similar to the midpoint formula in the 2D Cartesian plane, the midpoint of a line segment in 3D is the average of the coordinates of the endpoints.

! Exam tip

The formula booklet gives you the following formula:

The coordinates of the midpoint of a line segment with endpoints (x_1, y_1, z_1) and (x_2, y_2, z_2) are given by

$$M \left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}, \frac{z_1 + z_2}{2} \right).$$



More information

The image is a diagram showing a line segment with three labeled points: A, M, and B, displaying a collinear relationship. Each point is marked by a dot on the line. The text $AM = MB$ indicates that point M is equidistant from points A and B, making M the midpoint of the line segment AB. The diagram visually represents the concept of a midpoint in geometry.

[Generated by AI]



Student view

Example 2

Overview
(/study/ap

aa-
hl/sid-
134-
cid-
761926/o



If the endpoints of a line segment are A (1, -1, 2) and B (2, -2, 3) find the coordinates of the midpoint of the segment AB.

Steps	Explanation
Use	$M\left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}, \frac{z_1 + z_2}{2}\right)$
which gives	$M\left(\frac{1+2}{2}, \frac{-1+(-2)}{2}, \frac{2+3}{2}\right)$ $M\left(\frac{3}{2}, -\frac{3}{2}, \frac{5}{2}\right)$
So the midpoint has coordinates	$M\left(\frac{3}{2}, -\frac{3}{2}, \frac{5}{2}\right)$

Example 3



The coordinates of the two vertices of a parallelogram ABCD are A (2, 0, 0) and B (0, 0, 2).


If the intersection of diagonals of the parallelogram is M (0, 2, 0), find the coordinates of the other vertices.



Student
view



Overview
 (/study/ap
 aa-
 hl/sid-
 134-
 cid-
 761926/o



More information

This is a diagram of a parallelogram. It has four vertices labeled as A, B, C, and D. The point M, labeled at the center, represents the intersection of the diagonals of the parallelogram. The diagonals are shown as dashed lines, connecting vertex A to C, and B to D. The solid lines connect the vertices to form the parallelogram shape. The diagram seems to use a coordinate system with midpoint M at (0, 2, 0), although specific coordinates for other points are not shown in the image.

[Generated by AI]

M (0, 2, 0) is the midpoint of AC. A has coordinates (2, 0, 0), and C has coordinates (x_1, y_1, z_1) , so,

$$0 = \frac{2 + x_1}{2} \Rightarrow x_1 = -2$$

$$2 = \frac{0 + y_1}{2} \Rightarrow y_1 = 4$$



Student
view

$$0 = \frac{0 + z_1}{2} \Rightarrow z_1 = 0.$$



Therefore C has coordinates $(-2, 4, 0)$.

Overview
 (/study/ap
 aa-
 hl/sid-
 134-
 cid-
 761926/o
 —

As M $(0, 2, 0)$ is also the midpoint of BD, where the endpoints are B $(0, 0, 2)$ and D (x_2, y_2, z_2) ,

$$0 = \frac{0 + x_2}{2} \Rightarrow x_2 = 0$$

$$2 = \frac{0 + y_2}{2} \Rightarrow y_2 = 4$$

$$0 = \frac{2 + z_2}{2} \Rightarrow z_2 = -2$$

So D has coordinates $(0, 4, -2)$.

Distance between two points

The distance between A (x_1, y_1, z_1) and B (x_2, y_2, z_2) could be found using Pythagoras' theorem, as shown in the diagram below. First, using right-angled triangle AKC, you can find the length of AC, and then using triangle ACB you can find the length of AB.

$$AB^2 = (x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2$$

or

$$AB = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2}$$



① Exam tip

Student
view


The formula booklet gives you the following formula:



The distance between two points (x_1, y_1, z_1) and (x_2, y_2, z_2) is given by

Overview
 (/study/ap
 aa-
 hl/sid-
 134-
 cid-
 761926/o

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2}.$$



More information

The image is a 3D geometric diagram showing a cube with a triangular plane inside, connecting points A(x_1, y_1, z_1), B(x_2, y_2, z_2), C, and K. Point A is at the front bottom left and Point B is at the top right corner of the cube. Lines connecting these points form an orange triangular plane, marked as ABC. Point K is on the base of the cube under point B, and Point C is located on the bottom right corner of the face near the observer. The lines forming the cube and triangle are dashed, while the edges of the triangle are solid, highlighting its shape inside the cube. A right angle is indicated at point C, suggesting a perpendicular relationship with the base of the cube.

[Generated by AI]



Example 4

Student view





a) Find the exact distance between points A (-1,2,3) and B (-1,-2,4).

Overview
 (/study/ap
 aa-
 hl/sid-
 134-
 cid-
 761926/o

b) Round your answer in (a) to 3 significant figures.

$$AB = \sqrt{(-1 - (-1))^2 + (2 - (-2))^2 + (3 - 4)^2}$$

a) $AB = \sqrt{17}$ (exact)

b) $AB = 4.12$ (3 significant figures)

! Exam tip

Pay attention to the accuracy that the question asks for. An exact answer should be left in surd form or as a multiple of π .

For example, $\frac{\pi}{2}$ is an exact value but when rounded to 3 significant figures it is 1.57.

Example 5



A remote-controlled drone is 500 m north and 200 m east of its operator, and it is 20 m high. It is travelling due north at a constant height at a speed of 2 km h^{-1} .

a) What will be its coordinates to nearest metre after 30 minutes?

b) At this time, how far will the drone be from the operator? Give your answer to the nearest metre.

x
 Student
 view

a) Initial coordinates of the drone (north, east, height) are (500,200,20) 30 min = 0.5 hour so the distance it will travel is given by



Overview
 (/study/ap
 aa-
 hl/sid-
 134-
 cid-
 761926/o

$$\text{distance} = \text{speed} \times \text{time} = 0.5 \times 2 = 1 \text{ km} = 1000 \text{ m.}$$

As it is travelling due north the new coordinates will be (1500,200,20).


b) The operator is at the origin, at coordinates (0,0,0) and the new coordinates of the drone are (1500,200,20).

Using the distance formula $d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2}$

$$d = \sqrt{1500^2 + 200^2 + 20^2} = 1513 \text{ m to nearest metre.}$$

4 section questions ^

Question 1



Student
view

More information

ABCDEFGH is a cuboid. Give the letter of the point with coordinates (4, 3, 0).



Overview
(/study/ap
aa-
hl/sid-
134-
cid-
761926/o

- 1 B
2 G
3 H



- 4 F

Explanation


The perpendicular line to the x coordinate is $x = 4$.

The perpendicular line to y coordinate is $y = 3$.

As point B is on the x—y plane, z coordinate is 0.

Therefore (4,3,0) is point B.

Question 2



Student
view



Overview
(/study/ap

aa-
hl/sid-
134-
cid-
761926/o

More information

ABCDEFGH is a cuboid. What are the coordinates of point G?

1 (4,3,2)



2 (2,3,4)

3 (2,4,3)

4 (3,2,4)

Explanation

The perpendicular line to the x coordinate is $x = 4$.

The perpendicular line to the y coordinate is $y = 3$.

As point G is on the $x-y$ plane, z coordinate is 2.


Therefore G has coordinates (4,3,2).

Question 3



Student
view

Home
Overview
(/study/app
aa-
hl/sid-
134-
cid-
761926/o



More information

ABCDEF GH is a cuboid. Which of the following could be used to calculate the distance between points G and D?

1 $\sqrt{4^2 + 3^2 + 2^2}$

2 $\sqrt{(4 - 2)^2 + (3 - 3)^2 + (2 - 0)^2}$

3 $\sqrt{(4 - 3)^2 + (3 - 2)^2 + (2 - 0)^2}$

4 $\sqrt{4^2 + (3 - 2)^2 + 0^2}$

Explanation

G (4,3,2) and D (0,0,0).

Using the formula

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2}$$

then,



Student
view



Overview
(/study/ap

aa-
hl/sid-
134-
cid-
761926/o

$$\begin{aligned} d &= \sqrt{(4-0)^2 + (3-0)^2 + (2-0)^2} \\ &= \sqrt{4^2 + 3^2 + 2^2}. \end{aligned}$$

Question 4



Which one of the following is the exact distance between points?

$(-1, -3, 4)$ and $(0, 1, 2)$?

1 $\sqrt{21}$



2 3

3 $\sqrt{33}$

4 4

Explanation

Using the formula

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2}$$

then,

$$\begin{aligned} d &= \sqrt{(0 - (-1))^2 + (1 - (-3))^2 + (2 - 4)^2} \\ &= \sqrt{21}. \end{aligned}$$

3. Geometry and trigonometry / 3.1 Three-dimensional space

The right pyramid



Student
view




Overview
(/study/ap-
aa-
hl/sid-
134-
cid-
761926/o

☆ Definition

A pyramid is a polyhedron formed by connecting a polygonal base to its apex. Each triangle formed by the edge of the base and the apex is called a lateral face.

If the apex of the pyramid is directly above the centre of the base, then it is called a right pyramid.

The diagram shows a right pyramid with a rectangular base ABCD. Point E is its apex and G is the centre of the base. Line segment EG is perpendicular to the base of the pyramid.



More information



Student
view




Overview
(/study/ap
aa-
hl/sid-
134-
cid-
761926/o

The diagram features a right pyramid with a rectangular base labeled ABCD. Point E is the apex of the pyramid, and point G is the center of the rectangular base. The line segment EG is perpendicular to the base, indicating that E is directly above G. The four corners of the rectangular base are labeled B, C, A, and D in a clockwise order. The pyramid's apex and base are connected by triangular faces.

The base, ABCD, is shown as a parallelogram to represent the three-dimensional shape of the base when viewed from an angle. The vertical segment EG represents the height of the pyramid, and a right angle is marked at point G to highlight the perpendicular relationship between the line segment EG and the base ABCD.

[Generated by AI]

The base of a pyramid can have three or more sides. On the next diagram, you can see the triangular based pyramid which is also known as a tetrahedron.



More information



Student
view



Overview
(/study/ap
aa-
hl/sid-
134-
cid-
761926/o

The image is a diagram representing a triangular based pyramid also known as a tetrahedron. The tetrahedron is composed of a triangle with three vertices labeled A, B, and C. Lines connect these three points, forming the triangular base of the pyramid. There are no additional labels or text within the triangle. The diagram highlights the structure and shape of the tetrahedron as described in the accompanying text, emphasizing the triangular nature of the base and the overall pyramid-like structure of the shape.

[Generated by AI]

The diagram above shows a right pyramid with a triangle as the base (tetrahedron).

In right pyramids, the apex is exactly above the centre of the base. The illustration in the **Important** box shows a regular hexagonal based pyramid.

✓ **Important**

- The centre of the base is the intersection of diagonals, G .
- Length GK is the perpendicular height of the pyramid, h .
- Point H is the midpoint of edge ED .
- Connecting K to H gives the height of triangle KDE .
- The height of face KDE is the slant height of the pyramid, l .
- Connecting centre G to point H gives the height of the triangle formed by points GED; let GH = b .
- Triangle KGH is a right-angled triangle.


You need to be able to spot right-angled triangles to be able to calculate information about the pyramid.



Student
view



Overview
(/study/ap
aa-
hl/sid-
134-
cid-
761926/o



Interactive 3. Net of a Square Based Pyramid.

Credit: GeoGebra  (<https://www.geogebra.org/m/Aur2JZd5>) Lee W Fisher

 More information for interactive 3

The diagram illustrates a right pyramid with a regular hexagonal base. In the right pyramid the apex K is positioned directly above the center G of the base. The perpendicular height h of the pyramid is represented by the line segment GK, which is perpendicular to the base. Point H is the midpoint of edge ED, and connecting K to H gives the slant height ' l ' of the pyramid, which is the height of the triangular face KDE.

Connecting the center G to point H provides the height b of the triangle formed by points GED. The right-angled triangle KGH is essential for calculating various properties of the pyramid. By identifying and analyzing right-angled triangles within the pyramid, such as KGH, one can determine important measurements like the perpendicular height, slant height, and other geometric properties. This visualization helps in understanding the structural relationships and performing calculations related to the pyramid's dimensions.




Student
view



Surface area of a right pyramid

Overview

- (/study/ap aa-hl/sid-134-cid-761926/o) The net of a pyramid includes a base of n sides and n triangular faces, one on each edge of the base.
- 134-cid-761926/o

 [More information](#)

The image shows the net of a square-based pyramid. It consists of a square in the center, representing the base, with a triangle attached to each of the square's sides. The net forms a symmetric arrangement, where the triangles extend outward from each side of the square. This structure can be folded along the edges to form a three-dimensional square pyramid, where the triangles become the pyramid's lateral faces.

[Generated by AI]

The square-based pyramid above has a base with four sides. Therefore, the net is made up of four triangles and a square base.




Student view

The square-based pyramid has five faces in total, shown below.



Overview
(/study/app/math-aa-hl/sid-134-cid-761926/o)



More information

The image is a diagram showing the net of a square-based pyramid. It appears as a diamond shape with four congruent triangles forming the sides, arranged around a central square. Each triangle is connected to one side of the square, creating a cross-like pattern. The four triangular faces are meant to fold up from the center square to form the pyramid's shape, with the square acting as the base. Small squares are marked near each connecting edge, possibly indicating tabs or connection points for assembly. This diagram illustrates how the two-dimensional net of a pyramid can be folded to form a three-dimensional shape.

Section

Student... (0/0)

Feedback

Print (/study/app/math-aa-hl/sid-134-cid-761926/book/the-3d-coordinate-system-id-25408/print/)

Assign

[Generated by AI]


An applet for this net can be found below.



Student
view



Overview
(/study/ap
aa-
hl/sid-
134-
cid-
761926/o



interactive 1.Surface Area of a Right Pyramid.

More information for interactive 1

This interactive visualization demonstrates how to find the surface area of a right pyramid. By adjusting the slider 'c', users can observe the transition between the 2D net and the 3D pyramid. As c moves from 0 to 1, the pyramid's faces gradually unfold. At c = 0, the pyramid is fully constructed. As c increases, the triangular faces lift outward, revealing the 2D net. At c = 1, the net is completely unfolded, showing the square base and four separate triangular faces.

This helps in understanding how the total surface area is composed. As the slider moves to lower values, the triangular faces gradually fold upward, forming the pyramid. At the lowest position of the slider, the pyramid is fully constructed, showing how the lateral faces meet at the apex. This interactive approach




Student
view



Overview
(/study/ap
aa-
hl/sid-
134-
cid-
761926/o

allows users to explore how the surface area is determined by adding the base area and the combined area of the triangular faces.

The final diagram shows the net partially folded.



More information

A partially folded net diagram is shown. The diagram features a series of diamond shapes, each outlined in red, connected at intervals along the interior. The net folds upwards, creating a series of angular peaks and valleys, suggesting a three-dimensional structure at a midway folding stage. The diamonds likely denote connection points or folds critical for understanding the folding process. Additional text or labeling is not present in the image, focusing on the structural representation.

[Generated by AI]



Student
view




✓ **Important**

Total surface area of a right pyramid is the sum of the areas of the triangular faces and the base area.

Example 1



Pyramid ABCDE is square-based, as shown below. The length of the side of the square is 12 cm. The perpendicular height of the pyramid is 8 cm. Find the surface area of the pyramid.



More information






Overview
(/study/ap
aa-
hl/sid-
134-
cid-
761926/o

The image displays a square-based pyramid labeled as pyramid ABCDE. The base of the pyramid, ABCD, is a square with each side measuring 12 cm. The apex of the pyramid is point E, which is vertically above the center of the square base. The perpendicular height from the apex E to the base is indicated at point D and measures 8 cm. The vertices of the square are labeled as A, B, C, and D, each marked with a small circle at these points. Angles at the base are marked as right angles, emphasizing the geometric structure. The image is titled "Pyramid ABCDE" and illustrated in a purple and blue gradient tone.

[Generated by AI]

Examine the figure to see what information you have been given for triangle ABE.



Student
view



Overview
(/study/ap
aa-
hl/sid-
134-
cid-
761926/o

You know that the length $AB = 12 \text{ cm}$ since the base of the pyramid is a square.

The dimension 8 cm is not on triangle ABE since 8 is the distance OE – the perpendicular height of the pyramid.

Introduce points O (the centre of the base) and M (the midpoint of the edge AB).


The length EM is the height of triangle ABE. It is also the hypotenuse of triangle MOE.

$$AM = \frac{1}{2} \text{ of } AB, \text{ so } AM = 6 \text{ cm.}$$

Therefore, using Pythagoras' theorem,

$$EM = \sqrt{6^2 + 8^2} = 10.$$

The diagram shows triangle AEB.



Student
view



Overview

(/study/ap

aa-

hl/sid-

134-


cid-

761926/o

The area of the face ABE is

$$\text{Area}_{\triangle ABE} = \frac{1}{2}bh = \frac{1}{2} \times 12 \times 10 = 60 \text{ cm}^2.$$

The area of the base of the pyramid = $12 \times 12 = 144 \text{ cm}^2$.



Therefore, the total surface area is $4 \times 60 + 144 = 384 \text{ cm}^2$.

Student
view

Volume of a right pyramid

Overview
(/study/ap

aa-
hl/sid-
134-
cid-
761926/o

✓ Important

The volume of a pyramid is $\frac{1}{3}$ of the volume of the prism with the same height and base. Therefore, the formula for the volume of the pyramid is

$$V = \frac{1}{3} (\text{base area}) \times \text{height}$$

You can watch a demonstration of finding the volume of a square-based right pyramid in the following video.

Demo volume of pyramid



⌚ Exam tip

The formula booklet gives the following formula:

Volume of a right pyramid $V = \frac{1}{3} A \times h$, where A is the area of the base, h is the height.



Student
view



Example 2

Overview

(/study/ap

aa-

hl/sid-

134-


cid-

761926/o Pyramid ABCDE is square based, as shown in the diagram below. The length of the

side of the square is 12 cm. The perpendicular height of the pyramid is 8 cm.



Find the volume of the pyramid.



More information

The image depicts a geometric diagram of a pyramid with a square base labeled ABCD. The vertices are labeled A, B, C, D, and the apex E. The base ABCD is a square with sides measuring 12 units each, as indicated by labels on the side AB and side BC. The height of the pyramid is represented by a dotted line from the apex E to a point D on the base ABCD, marked as 8 units. Angles at vertices A, B, and C appear to be right angles, indicated by small squares.

[Generated by AI]



Student view



Overview
(/study/ap

aa-
hl/sid-
134-
cid-
761926/o

To find the volume of the pyramid, use the formula

$$V = \frac{1}{3}Ah,$$

where A is the area of the base and h is the perpendicular height from the base to the apex (vertex E).

The base is a square. Therefore, $A = 12 \times 12 = 144$.

$$V = \frac{1}{3} \times 144 \times 8 = 384$$

So the volume of the pyramid is 384 cm^3 .


3 section questions ^

Question 1

Difficulty:



★★☆



Student
view



Overview
(/study/aa-hl/sid-134-cid-761926/o)

More information

- aa-hl/sid-134-cid-761926/o
- A right pyramid with regular hexagonal base has a perpendicular height of 5 cm.
- Each side of the hexagon is 4 cm.

Find the area of the base of the pyramid correct to 3 significant figures.

You do not need to give the units with your answer.


41.6



Accepted answers

41.6, 41,6

Explanation



More information

When you draw the diagonals of the hexagon you get six equilateral triangles with the side length 4 cm.

You can find the height of these triangles using Pythagoras' theorem.



Student view

$$h = \sqrt{4^2 - 2^2} = \sqrt{12}$$

We can use that the area of a triangle is half of the product of the base and the height.



Overview
 (/study/ap
 aa-
 hl/sid-
 134-
 cid-
 761926/o

$$A = \frac{1}{2} \times 4 \times \sqrt{12} = 2\sqrt{12}$$

Each triangle has area $2\sqrt{12} \text{ cm}^2$


There are six triangles, so the area of the hexagon is six times this value.

total area = $6 \times 2\sqrt{12} = 41.56 \dots = 41.6 \text{ cm}^2$ (3 significant figures)

Question 2



★★☆



More information

A right pyramid with regular hexagonal base has a vertical height of 5 cm.

Each side of the hexagon is 4 cm.

Find the volume of the pyramid.

Give your answer as a decimal correct to 3 significant figures.



Student view

You do not need to give the units with your answer.



69.3

Overview
 (/study/ap
 aa-
 hl/sid-
 134-
 cid-
 761926/o

Accepted answers

69.3, 69,3

Explanation

Using the formula area of triangle = $\frac{1}{2}abs\sin C$, each of the triangles in the base has an area of

$$\frac{1}{2} \times 4 \times 4 \times \sin 60^\circ = 4\sqrt{3} \text{ cm}^2,$$

Therefore the total area is

$$A = 6 \times 4\sqrt{3} = 24\sqrt{3} \text{ cm}^2$$

Using the formula for the volume of a right pyramid:

$$V = \frac{1}{3}A \times h, \text{ where } A \text{ is the area of the base and } h \text{ is the height, gives}$$

$$V = \frac{1}{3} \times 24\sqrt{3} \times 5 = 40\sqrt{3} = 69.28\dots \text{ cm}^3$$

So the volume is 69.3 cm³ (to 3 significant figures).

Question 3


The pyramid in the diagram has a pentagonal base.



Student
view



Overview
(/study/app/math-aa-hl/sid-134-cid-761926/o)



More information

How many faces does this pyramid have?

1 6



2 5

3 10

4 12

Explanation

The base has 5 edges. On each of these edges is a different triangular face.

There are 5 triangular faces and 1 pentagonal face, which is a total of 6 faces.

3. Geometry and trigonometry / 3.1 Three-dimensional space

The cone



Student
view



Overview
(/study/ap)


aa-
hl/sid-
134-
cid-

761926/o

★ Definition

A cone is a solid with a circular base and an apex. It has two faces, the circular base and a sector of a circle which has the circumference of the base as its arc.

The diagram shows a cone with base radius r and perpendicular height h along with its net. The length $AB = l$ is called the slant height of the cone. AGB is a right triangle. The circumference of the base of the circle is equal to the length of the arc BE.



More information



Student
view



Overview
(/study/ap
aa-
hl/sid-
134-
cid-
761926/o

The diagram illustrates a three-dimensional cone with a circular base. The cone is shown alongside its net, which is used to represent the cone as a two-dimensional shape. The cone itself has a base with radius ' r ' and a perpendicular height ' h ', with its apex labeled as point A, the base center as point G, and the base perimeter intersecting at point B. The slant height of the cone, denoted by ' l ', is the hypotenuse of the right triangle AGB.

The net of the cone is displayed below the cone. It consists of a circle with radius ' r ', centered at point G, representing the base of the cone, and an adjoining circular sector that forms the lateral surface of the cone when folded. The sector has a radius equal to the slant height ' l ' of the cone. The arc BE of this sector is equal in length to the circumference of the cone's base. The labeling on the diagram includes the points A, B, G, and E, and the values for radius ' r ', slant height ' l ', and perpendicular height ' h '. This illustrates the geometric relationship between the physical cone and its geometric representation as a net.

[Generated by AI]

Area of the curved surface of a cone

! Exam tip

The formula booklet gives you the following formula:

Area of the curved surface of a cone $A = \pi r l$, where r is the radius, l is the slant height.

Example 1



Student
view



Overview
(/study/ap
aa-
hl/sid-
134-
cid-
761926/o



A piping nozzle

turbojet Getty Images

The photograph shows a piping nozzle, which is used to apply icing onto cakes in a decorative manner. It is the curved surface of a cone. There is no base.

A piping nozzle has the following dimensions. The radius of the base is 15 mm and the radius of the top is 3 mm. It is formed by cutting a small cone from the top of a larger cone. The perpendicular height of the large cone is 36 mm. The perpendicular height of the cone that is removed from the top is 7.2 mm.


Find the surface area of the piping nozzle, correct to 3 significant figures.



Student
view

Home
Overview
(/study/app/math-aa-hl/sid-134-cid-761926/o)

aa-
hl/sid-
134-
cid-
761926/o



Steps	Explanation
Call the curved surface area of the larger cone A_1 , the curved surface area of the smaller cone A_2 and the area required A_{nozzle} .	You are only interested in surface area, as the pipe does not have a base. You need to find the curved surface area of the larger cone, then the curved surface area of the smaller cone.
Using Pythagoras' theorem $l = \sqrt{15^2 + 36^2} = 39$.	From the formula booklet the surface has area $A = \pi r l$ the value of h but not l . Use Pythagoras' theorem to find l .
$A_1 = \pi \times 15 \times 39 = 1837.83 \dots \text{mm}^2$	Use the formula $A = \pi r l$.
The slant height of the smaller cone is $l = \sqrt{3^2 + 7.2^2} = 7.8$.	Use Pythagoras' theorem.
So $A_2 = \pi \times 3 \times 7.8 = 73.51 \dots \text{mm}^2$.	Use $A = \pi r l$.

Home
Overview
(/study/ap
aa-
hl/sid-
134-
cid-
761926/o

Steps	Explanation
$\begin{aligned} A_{\text{nozzle}} &= 1837.83 - 73.51\dots \\ &= 1764.31\dots \\ &= 1760 \text{ (3 significant figures)} \end{aligned}$	Use $A_{\text{nozzle}} = A_1 - A_2$.
So the surface area of the piping nozzle is 1760 mm^2 to 3 significant figures.	Remember to give the answer.

Volume of a right cone

Finding the volume of a cone is similar to the volume of a pyramid.

$$V = \frac{1}{3} (\text{base area}) \times \text{height.}$$

In this case, the base area is a circle $A = \pi r^2$. Therefore, $V = \frac{1}{3} \pi r^2 h$.

① Exam tip

The formula booklet gives you the following formula:

Volume of a right cone $V = \frac{1}{3} \pi r^2 h$, where r is the radius and h is the height.

Example 2



Find the exact volume of the cone with base radius 3 cm and perpendicular height 7 cm .



Student view

↪
 Overview
 (/study/ap
 aa-
 hl/sid-
 134-
 cid-
 761926/o

Using the volume formula $V = \frac{1}{3}\pi r^2 h$

$$\begin{aligned} V &= \frac{1}{3}\pi(3)^2 7 \\ &= 21\pi \end{aligned}$$

So the volume is $V = 21\pi \text{ cm}^3$.

① Exam tip

When the question is asking for an exact value, and π is involved in the calculation, leave your answer in terms of π . Do not round your answer, and make sure to include units.

2 section questions ^

Question 1



A waffle ice cream cone has diameter 5 cm and slant height 12 cm.

Which of the following is the correct calculation for finding the volume of the cone?

1 $V = \frac{1}{3}\pi \times 2.5^2 \times \sqrt{(12^2 - 2.5^2)}$



2 $V = \frac{1}{3}\pi \times 5^2 \times \sqrt{(12^2 - 5^2)}$

3 $V = \frac{1}{3}\pi \times 2.5^2 \times 12$

4 $V = \frac{1}{3} \times A \times 12$



Student
view

Explanation

The volume of a cone can be found using the formula



Overview
 (/study/ap
 aa-
 hl/sid-
 134-
 cid-
 761926/o

$$V = \frac{1}{3}\pi r^2 h.$$

However, h is the perpendicular height and you are given the slant height.

The radius of the circle is 2.5 cm.

From Pythagoras' theorem: $h = \sqrt{(12^2 - 2.5^2)}$.

You are told that $r = 2.5$ and $h = \sqrt{(12^2 - 2.5^2)}$.

Therefore, substituting these into the volume formula gives

$$V = \frac{1}{3}\pi r^2 h = \frac{1}{3}\pi \times 2.5^2 \times \sqrt{(12^2 - 2.5^2)}.$$

Question 2



A solid cone has slant height 2 m and radius 1.6 m. Find the total surface area of the cone in metres. Give your answer correct to three significant figures without the unit.

18.1



Accepted answers

18.1, 18,1

Explanation

According to the formula, area of the curved surface of a cone $A = \pi r l$, where r is the radius and l is the slant height.

$$\begin{aligned} \text{Area of curved surface} &= \pi \times 1.6 \times 2 \\ &= 3.2\pi \end{aligned}$$

$$\begin{aligned} \text{Area of base} &= \pi r^2 \\ &= 2.56\pi \end{aligned}$$



Student
view


Total surface area $= 3.2\pi + 2.56\pi = 18.1 \text{ m}^2$ to three significant figures



Overview
(/study/app/math-aa-hl/sid-134-cid-761926/o)

3. Geometry and trigonometry / 3.1 Three-dimensional space

The sphere and hemisphere



More information

The image is a Venn diagram consisting of two overlapping circles. The larger circle on the right is blue, representing the superset of items labeled as "universal set." Inside this circle, near the center, is a smaller purple circle labeled "B," likely representing a subset. On the left, partially overlapping the blue circle, is another smaller purple circle labeled "A," representing another subset. The overlap area between "A" and the blue section indicates the common elements shared by sets "A" and "B." This Venn diagram visually depicts the relationships between different sets or groups, facilitating understanding of how the subsets intersect with the universal set.

[Generated by AI]



Section

Student... (0/0)

Feedback



Print (/study/app/math-aa-hl/sid-134-cid-761926/book/the-right-pyramid-id-25409/print/)

Assign

Student view



Overview

(/study/ap)

aa-
hl/sid-
134-
cid-

761926/o



International Mindedness


'When a semicircle with fixed diameter is carried round and restored again to the same position from which it began to be moved, the figure so comprehended is a sphere.'

This is how Euclid defined a sphere in his *Elements, Book 11 Definition 14*. It was written c. 300 BCE. *The Elements* was one of the very first text books. People used it to study mathematics for more than 2000 years!

Hemisphere

A hemisphere is one half of a sphere. In world affairs, you often refer to the northern hemisphere or the southern hemisphere.

If you rotate an object, say the Earth, clockwise, then as seen from above the plane the rotation is clockwise, but if you look underneath the plane ...



More information

The image is a diagram illustrating the perspective of rotation. It shows a semicircular shape that represents an object, such as the Earth, viewed from a side angle. On the left side of the semicircle, a dot is marked, indicating the viewpoint from above, where the rotation appears clockwise. On the opposite side, another dot represents the viewpoint from below, where the rotation appears counterclockwise. The diagram effectively visualizes how the direction of rotation can seem different depending on the observer's perspective, as described in the surrounding text.

Student
view



[Generated by AI]

Overview
 (/study/app/math-aa-hl/sid-134-cid-761926/o)

... down here the rotation is counterclockwise! Here is a simple experiment you might like to try. Take a plate. Hold it up between you and a friend. Rotate it clockwise. They will tell you that it is rotating counterclockwise. It's a matter of perspective!

Surface area of a sphere

① Exam tip

The formula booklet gives you the following formula:

Surface area of a sphere $A = 4\pi r^2$, where r is the radius.

Example 1



Find the surface area of a sphere with radius 3.4 cm. Round your answer to the nearest cm^2 .

Using the area formula $A = 4\pi r^2$

$$\begin{aligned} A &= 4\pi(3.4)^2 \\ &= 145.267 \dots \end{aligned}$$

So the area is 145 cm^2 to the nearest cm^2 .

Section

Student... (0/0)

Feedback

Print (/study/app/math-aa-hl/sid-134-cid-761926/book/the-cone-id-25410/print/)

Assign



Student view

Volume of a sphere

Overview
(/study/ap

aa-
hl/sid-
134-
cid-
761926/o

Exam tip

The formula booklet gives you the following formula:

$$\text{Volume of a sphere } V = \frac{4}{3}\pi r^3, \text{ where } r \text{ is the radius.}$$

Example 2



Find the volume of a sphere with radius 3.4 cm. Round your answer to nearest cm^3 .

Using the volume formula $V = \frac{4}{3}\pi r^3$

$$\begin{aligned} V &= \frac{4}{3}\pi(3.4)^3 \\ &= 164.636\dots \end{aligned}$$

So the volume is 165 cm^3 to the nearest cm^3 .

2 section questions ^

Question 1



A sphere has radius 3.5 cm.



Student
view



Overview

(/study/app

aa-

hl/sid-

134-

cid-

761926/o

Which of the following is the correct calculation for finding the **volume** of the sphere?

1 $V = \frac{4}{3}\pi(3.5)^3$



2 $V = 3\pi(3.5)^2$

3 $V = 3\pi(3.5)^3$

4 $V = \frac{4}{3}\pi(3.5)^2$

Explanation

The formula for the volume of a sphere is

$$V = \frac{4}{3}\pi r^3$$

Substituting $r = 3.5$, gives

$$V = \frac{4}{3}\pi(3.5)^3 \text{ (cm}^3\text{)}$$

Question 2



A hemisphere has diameter 20 cm.

Which of the following is the correct calculation for finding the total surface area?

1 $3\pi(10)^2$



2 $3(10)^2$

3 $3\pi(20)^2$

4 $4\pi(10)^2$

Student
view

Explanation



The formula for the curved surface area of a sphere is $A = 4\pi r^2$ so for a hemisphere it is $A = 2\pi r^2$.

Overview
(/study/ap
aa-
hl/sid-
134-
cid-
761926/o

You are given that the diameter of the hemisphere is 20 cm. Therefore, the radius is 10 cm.

Substituting $r = 10$ gives curved surface area $2\pi(10)^2$.

You are asked for the total surface area so need to add the area of the circular base.

Area of circle = πr^2 .

As the radius is 10 cm, the area of the base = $\pi(10)^2$.

So the total surface area = $\pi(10)^2 + 2\pi(10)^2 = 3\pi(10)^2$.

3. Geometry and trigonometry / 3.1 Three-dimensional space

Combined solids

Combining shapes in design and architecture

Many everyday objects and many buildings are made up of a combination of several 3D geometrical shapes. The photograph shows a row of historical town houses. Can you identify some of the 3D shapes you have studied so far in these buildings? How are they combined?



Student
view




Overview
 (/study/app/math-aa-hl/sid-134-cid-761926/print)
 aa-
 hl/sid-
 134-
 cid-
 761926/o



Town houses

Credit: amedved Getty Images

The diagram below shows a spinning top made up of several different sized spheres, and the bottom part is a cone. Why do you think the centre is a sphere and the bottom part is a cone?



Example 1

Section

Student... (0/0)

Feedback

Print (/study/app/math-aa-hl/sid-134-cid-761926/book/the-sphere-and-hemisphere-id-25411/print/)



Student view


A plastic model of an ice-cream cone comprises a cone with height 8 cm and a hemisphere with radius 4 cm.



Calculate the total volume and the total surface area of the model.

Overview
 (/study/ap
 aa-
 hl/sid-
 134-
 cid-
 761926/o

Give your answer correct to 3 significant figures.



volume of the combined solids = volume of the cone + volume of the l

$$\begin{aligned}
 V &= \frac{1}{3}\pi r^2 h + \frac{1}{2} \left(\frac{4}{3}\pi r^3 \right) \\
 &= \frac{1}{3}\pi(4)^2(8) + \frac{1}{2} \left(\frac{4}{3}\pi(4)^3 \right) \\
 &= 268.082\dots
 \end{aligned}$$



Student
view

So the volume of the combined solid is 268 cm³ correct to 3 significant figures.



You can use the surface area formulas to find the combined surface area.

Overview
 (/study/ap
 aa-
 hl/sid-
 134-
 cid-
 761926/o

- The area of the curved surface of a hemisphere is $2\pi r^2$.
- The area of the curved surface of a cone is $\pi r l$, where the slant height can be calculated using Pythagoras' theorem, $l = \sqrt{h^2 + r^2}$.

You can find the total surface area combining these formulae.

$$\begin{aligned} A &= 2\pi r^2 + \pi r \sqrt{h^2 + r^2} \\ &= 2\pi(4^2) + \pi(4)\sqrt{8^2 + 4^2} \\ &= 212.929\dots \end{aligned}$$

So the surface area of the combined solid is 213 cm^2 correct to 3 significant figures.



Theory of Knowledge

As seen in the video below, there are many people who sincerely believe that the earth is flat. Is it possible to use trigonometry or geometry to ‘prove’ that the earth is not flat? Consider the knowledge question, ‘How much knowledge must one have in regard to something before we can say for “certain” that it is true?’



Student
view


Home
Overview
(/study/ap...
aa-
hl/sid-
134-
cid-
761926/o

Inside a Flat Earth convention, where nearly everyone belie...



2 section questions ^

Question 1



More information

A wooden spinning toy is made up of a cone and a hemisphere with a cylinder in between.



Student view

The height of the cone is 4 cm and it has a base radius of 2 cm.



What is the radius of the hemisphere?

Overview
 (/study/ap
 aa-
 hl/sid-
 134-
 cid-
 761926/o


- 1 2 cm
- 2 4 cm
- 3 1 cm
-
- 4 6 cm



Explanation

The radius of the base of the cone is the same of the radius of the hemisphere, therefore it is 2 cm.

Question 2



More information

A wooden spinning toy is made up of a cone and hemisphere with a cylinder in between.



The height of the cone is 4 cm and it has a base radius of 2 cm.

Student
view

What is the total volume of the toy correct to 3 significant figures?

Home
Overview
(/study/app)



aa-
hl/sid-
134-
cid-
761926/o

1 96.3 cm^3

2 113 cm^3

3 130 cm^3

4 92.5 cm^3

Explanation

As the radius of the base of the cone is the same of the radius of the hemisphere, it is 2cm.

Total volume is

$$\begin{aligned} V_{\text{total}} &= V_{\text{cone}} + V_{\text{cylinder}} + V_{\text{hemisphere}} \\ V_{\text{total}} &= \frac{1}{3}\pi(2)^24 + \pi(2)^25 + \frac{1}{2}\left(\frac{4}{3}\pi2^3\right) \\ &= 96.342\dots \end{aligned}$$

So the total volume is 96.3 cm^3 to 3 significant figures.

3. Geometry and trigonometry / 3.1 Three-dimensional space

Angles between lines

Right triangles in 3D shapes

As you have seen in previous sections, you can draw right-angled triangles within 3D shapes. In the diagram below, you can see a right triangle formed by the diagonal of the base DB, DL and BL. Using these right triangles, you can calculate the angles between lines, such as angle LBD = β in the diagram.

Student view




Overview
(/study/app/math-aa-hl/sid-134-cid-761926/o)

aa-
hl/sid-
134-
cid-
761926/o

Section

Student...

(0/0) 5 cm



Feedback

Print

134-cid-761926/book/combined-solids-id-25412/print/

Assign

More information

The image is a 3D diagram of a cuboid with points labeled A, B, C, D, I, J, K, and L at each corner. A right-angled triangle is highlighted within the cuboid, with points D, B, and L forming the vertices of the triangle. The base of the triangle is the diagonal DB of the cuboid's bottom face, with a length of 8 cm marked. The perpendicular side DL (vertical leg) of the triangle is 5 cm long, and the horizontal leg DA is 3 cm. The angle at B, labeled as β , is the angle of interest and forms part of the triangle with a right angle marked at D. The cuboid's vertical faces and additional dimensions are depicted but not labeled. The diagram illustrates how a 3D shape can be analyzed through its 2D triangular cross-section to calculate angles.


[Generated by AI]

Suppose three points are drawn on the edges of a cuboid and joined to form a triangle. Sometimes the triangle will be right-angled; sometimes it will not be right-angled.

Student view

Use the applet below to move the points on the edges of the cuboid to get a sense of when a triangle will be right-angled.

Home
Overview
(/study/ap-
aa-
hl/sid-
134-
cid-
761926/o



Interactive 2. Right Triangles in 3D Shapes.

Credit: [GeoGebra](https://www.geogebra.org/m/jMXT6f1P) ↗ (<https://www.geogebra.org/m/jMXT6f1P>) Lee W Fisher

🔗 More information for interactive 2


The image shows a 3D cuboid with labeled vertices: A, B, C, D, E, F, G, and H. The edges of the cuboid are highlighted in dark blue, and purple points are placed along its edges. These points can be moved along the edges to form different triangles.

This interactive allows users to explore the properties of triangles formed within a cuboid by moving its eight vertices. By adjusting the positions of these points along the edges, users can observe how different triangles are created and identify when a triangle becomes right-angled. As the points are moved, the dimensions of the cuboid change, altering the relationships between its edges and diagonals. Users can experiment by aligning points to form right-angled triangles along faces, space diagonals, and edge connections.



Student
view

Home
Overview
(/study/ap
aa-
hl/sid-
134-
cid-
761926/o



Interactive 3. Formation of Right Triangles in 3D Shapes.

Credit: GeoGebra [\(https://www.geogebra.org/m/zagcpBoZ\)](https://www.geogebra.org/m/zagcpBoZ) Lee W Fisher

 More information for interactive 3

This interactive allows users to explore the formation of right-angled triangles within a cuboid by moving three points along the edges.

As the user moves the points, a continuous angle measurement is shown, updating in real-time to indicate whether the triangle remains right-angled or not.


As users adjust the points, the triangle dynamically updates, and a 90° angle is displayed when the conditions for a right-angled triangle are met. A right-angled triangle appears when one side is parallel to an edge of the cuboid, and the second shorter side lies within one of the cuboid's faces. No matter where the third point is placed along an edge, as long as these conditions are satisfied, the triangle remains right-angled. Hence, the interactive clears the understanding of the conditions required to get a right angled triangle.

Notice that when one edge of the triangle is parallel to an edge of the cuboid, (forming the first shorter side) and the second shorter side lies on a face of the cuboid, the resulting triangle is right-angled, regardless of where on an edge the third point is.

 Student view

Example 1

Overview
 (/study/ap-
 aa-
 hl/sid-
 134-
 cid-
 761926/o)



More information

The diagram shows a cuboid with vertices labeled A, B, C, D, I, J, K, and L. The edges of the cuboid have the following lengths: $AB = 8 \text{ cm}$, $AD = 3 \text{ cm}$, and $AI = 5 \text{ cm}$. The diagram appears to illustrate spatial relationships and angles. Particular emphasis is placed on the triangle formed by vertices D, I, and B, which is highlighted in orange. Vertex D has a perpendicular angle marked, and the angle at vertex B is labeled as beta (β). The diagram helps in understanding the geometrical representation of the cuboid's dimensions and the relationships between points and angles.

[Generated by AI]

In the diagram, the cuboid has lengths $AB = 8 \text{ cm}$, $AD = 3 \text{ cm}$ and $AI = 5 \text{ cm}$.

Calculate the angle between the diagonal DB and LB.




Student
view

🏠
 Overview
 (/study/ap
 aa-
 hl/sid-
 134-
 cid-
 761926/o

Consider the right-angled triangle drawn in the cuboid.

First, you need to find the length of the diagonal of the base ABCD.



You need to find the length of the diagonal of the base ABCD using Pythagoras' theorem:

$$\begin{aligned} DB^2 &= 3^2 + 8^2 \\ &= \sqrt{73} \end{aligned}$$

So the length of the diagonal is $\sqrt{73}$ cm.


❗ Exam tip

Do not round your answer at this stage. Leave it in surd form.

✖
 Student
 view



Overview
 (/study/app/math-aa-hl/sid-134-cid-761926/o)



Now that you have the two of the sides of the right-angled triangle, you can use trigonometric ratios. You know the opposite and adjacent sides to the angle you need to find, so use the tangent ratio:

$$\tan \beta = \left(\frac{5}{\sqrt{73}} \right)$$

$$\begin{aligned} \beta &= \tan^{-1} \left(\frac{5}{\sqrt{73}} \right) \\ &= 30.336\dots^\circ \end{aligned}$$

So $\beta = 30.3^\circ$ to 3 significant figures.

① Exam tip

Only round the final answer.



Student
view




Example 2



Overview
(/study/ap
aa-
hl/sid-
134-
cid-
761926/o)

Cuboid ABCDEFGH has dimensions 8 cm, 6 cm and 4 cm, as shown in the diagram below.

Calculate the angle between line segments FA and AE .



More information

This is a diagram of a 3D geometric shape with eight points labeled A, B, C, D, E, F, G, and H. The shape appears to resemble a cuboid with additional lines connecting the labeled points. The face AEFC is highlighted in orange, suggesting it's the area of interest. The lengths of some of the line segments are given: segment AE is 8 cm, and segments AD and EF are 6 cm and 4 cm, respectively. The task is to calculate the angle between the line segments FA and AE. Each segment represents an edge of the shape and is part of the overall geometric configuration.

[Generated by AI]



Student
view




Overview
(/study/ap
aa-
hl/sid-
134-
cid-
761926/o

Steps	Explanation
<p>Calculate the length of AE:</p> $(AE)^2 = 8^2 + 4^2 = 80$ <p>Therefore, $AE = \sqrt{80}$.</p>	<p>Before you isolate triangle, make sure you know which side is the hypotenuse. The angle is at E; therefore, AE is the hypotenuse.</p>



Student view

Home
Overview
(/study/ap
aa-
hl/sid-
134-
cid-
761926/o

Steps	Explanation
	The final step is $\angle FAE$.

① Exam tip

Make sure your calculator mode is set to degrees when you are solving this type of problem.

3 section questions ^



Student view

Question 1





Overview

/study/ap

aa-


hl/sid-

134-

cid-

761926/o

The diagram below shows the cuboid ABCDEFGH.



More information

The length of AC is \sqrt{a} cm.

Find the value of a .

Give your answer as an integer. You do not need to include units.

244



Accepted answers

244, a=244

Explanation

The triangle required is triangle ABC.

It is right-angled at B. $AB = 12 \text{ cm}$ and $BC = 10 \text{ cm}$.

Therefore, by Pythagoras' theorem: $AC = \sqrt{12^2 + 10^2} = \sqrt{144 + 100} = \sqrt{244}$.


Therefore, $a = 244$.



Student
view

**Question 2**

The diagram below shows the cuboid ABCDEFGH.



More information

Find the length of AH.

Give your answer to the nearest centimetre. You do not need to include the units in your answer.

18

**Accepted answers**

18

Explanation**Method 1**

This is a two-step question where first you calculate the base diagonal AC. The second step is to calculate the space diagonal AH.

$$\text{Step 1: } AC = \sqrt{12^2 + 10^2} = \sqrt{244}$$

$$\text{Step 2: } AH = \sqrt{(AC)^2 + (CH)^2} = \sqrt{\sqrt{244})^2 + 8^2} = \sqrt{244 + 64} = \sqrt{308} \approx 18 \text{ cm}$$



Overview
(/study/ap

aa-
hl/sid-
134-
cid-
761926/o

Method 2

A commonly used formula for a space diagonal d is $d^2 = a^2 + b^2 + c^2$.

In this formula, d is the space diagonal, and a , b and c are the length, width and height in any order.


Using this formula, $d^2 = 12^2 + 10^2 + 8^2 = 308$.

Therefore $d = \sqrt{308} \approx 18$ cm.

Question 3



The cuboid ABCDEFGH is drawn below.



More information

M is the midpoint of GH, and N is the midpoint of BC.

Find the angle MDN in degrees, correct to 3 significant figures.

Do not write the degree sign in your answer.



31.6



Student
view

Accepted answers



31.6


Overview
 (/study/ap-
 aa-
 hl/sid-
 134-
 cid-
 761926/o)

Explanation

Calculate the length of DN.

Use right-angled triangle NCD on the base of the cuboid.


Note that $CN = 5$, because N is the midpoint of BC.



More information

Therefore, $DN = \sqrt{5^2 + 12^2} = \sqrt{169} = 13$.

You are now ready to isolate triangle DNM.



More information



Student
view



Overview
(/study/ap

aa-
hl/sid-
134-
cid-
761926/o

$$\tan\theta = \left(\frac{8}{13}\right)$$

$$\begin{aligned}\theta &= \tan^{-1}\left(\frac{8}{13}\right) \\ &= 31.607\dots^\circ\end{aligned}$$

So $\theta = 31.6^\circ$ (3 significant figures).

3. Geometry and trigonometry / 3.1 Three-dimensional space

Angle between a line and a plane

To calculate the angle between a line and a plane, or the face of a solid, you use a three-step calculation. The first step is simply to determine the correct line to use on the face of the solid. The next two steps are the same as in the previous example.


This time, you need to calculate the angle between the line GN and the base ABCD. N is the midpoint of AD.



Student
view



Overview
 (/study/ap
 aa-
 hl/sid-
 134-
 cid-
 761926/o



More information

The image shows a 3D diagram of a cuboid labeled with various points: A, B, C, D, E, F, G, H, and N. The points form the vertices of a rectangular cuboid. The line GN connects point G at the top with point N on the base, given that N is the midpoint of the line segment AD on the cuboid's base ABCD. The base ABCD is marked with side lengths, with side AB equaling 8 and side AD equaling 6. Additionally, the height from the base to the top of the cuboid, represented by points such as G and F, is 4. Line GN needs to be analyzed to determine the angle between it and the base of the cuboid.


[Generated by AI]

Step 1 Project the upper point to the base. In other words, drop a line from G to the base. That takes you to the point B. Your triangle of study is GBN. Because GB is perpendicular to the base ABCD, the triangle GBN is right-angled at B. Click on the link below to enable you to rotate the cuboid so that you can see it better.



Student
view

Home
Overview
(/study/app/math-aa-hl/sid-134-cid-761926/o)



Interactive 1. Angle Between a Line and a Plane.

More information for interactive 1

This interactive applet demonstrates a step-by-step method to calculate the angle between a line and a plane, specifically between the line GN and the base ABCD of a cuboid. Here, N is the midpoint of AD. Users can manipulate the cuboid's vertices to explore different configurations, allowing them to practice and gain a deeper understanding of this concept. The process involves projecting a point onto the base plane to form a right-angled triangle, calculating the necessary side lengths using geometric principles such as the Pythagorean theorem, and then applying trigonometric functions, such as tangent, to determine the angle between the line and the plane.

They learn how to apply the Pythagorean theorem to determine unknown side lengths and use trigonometric ratios to calculate the desired angle.

For example, to find the angle between GN and the base ABCD, the user must first determine at least two sides of the right-angled triangle. Given that BG = 4 cm, they need to calculate BN. With AB = 6 cm and AN = 4 cm, applying the Pythagorean theorem gives

$$BN^2 = 6^2 + 4^2 = 36 + 16 = 52, \text{ leading to } BN = \sqrt{52}.$$

Using trigonometric functions, such as $\tan\theta = \frac{\text{opposite}}{\text{Adjacent}}$, they can then compute the angle.

Since the cuboid can be rotated and viewed from different angles in 3D, users can interactively explore how the angle between the line and the plane changes with different perspectives. This hands-on




Student view



Overview
(/study/app/math-aa-hl/sid-134-cid-761926/o)

exploration enhances spatial reasoning skills and provides a clearer understanding of geometric transformations in three-dimensional space, making abstract mathematical concepts more tangible and engaging.

**Section**

Student... (0/0)

Feedback

Print (/study/app/math-aa-hl/sid-134-cid-761926/book/angles-between-lines-id-25413/print/)

Assign

More information

The image is a 3D geometric diagram representing a rectangular prism. The diagram is labeled with points A to H on the prism's vertices. Lines are drawn between these points to form rectangles and triangles, showing different two-dimensional shapes on the prism's surfaces. There are specific labels such as $BG = 4 \text{ cm}$, $AB = 6 \text{ cm}$, $AN = 4 \text{ cm}$, and BN is calculated as the square root of 52. The image includes a highlighted orange triangle within the blue prism, demonstrating a right angle of 90° at point B. The labels 8, 6, and 4 along the edges signify the dimensions of the prism. This structure is useful for solving geometric problems related to angles and sides within 3D shapes.

[Generated by AI]

Section

Student... (0/0)

Feedback

Print (/study/app/math-aa-hl/sid-134-cid-761926/book/angle-between-a-line-and-a-plane-id-25414/print/)

Assign




Student
view

Overview
(/study/ap
aa-
hl/sid-
134-
cid-
761926/o

Step 2 To calculate the required angle, you need to know at least two sides of the triangle. You know that $BG = 4 \text{ cm}$. Next you need to calculate BN . Now $AB = 6$ and $AN = 4$, so $(BN)^2 = 6^2 + 4^2 = 52$; therefore, $BN = \sqrt{52}$.

Step 3 Now you are ready to calculate $\angle GNB$, which is the angle between GN and the base ABCD. This angle is marked as θ in the diagram below.



More information

The image depicts a right triangle with vertices labeled G, N, and B. The side GN is the hypotenuse. The side GB is labeled with a length of 4 units, and BN with a length of $(\sqrt{52})$. There is a right angle marked at vertex B. The angle (θ) , formed between GN and BN, is highlighted as well. This diagram is used to illustrate the calculation of the angle $(\angle \rm{GNB})$, based on trigonometric relationships, specifically using the tangent function $(\tan \theta = \frac{4}{\sqrt{52}})$. An angle (θ) is marked near vertex N.

[Generated by AI]

Student view

$$\tan \theta = \left(\frac{4}{\sqrt{52}} \right)$$



Overview

(/study/ap

aa-

hl/sid-

134-

cid-

761926/o

$$\theta = \tan^{-1} \left(\frac{4}{\sqrt{52}} \right)$$

$$= 29.017\dots^\circ$$

So $\theta = 29.0^\circ$ (to 3 significant figures).

Example 1



The diagram below shows a square-based right pyramid ABCDE. The height of the pyramid is 6 cm and each of the sides of the base measures 10 cm. M is the midpoint of the line segment CD.

Find the following angles. Give your answers correct to 3 significant figures.

a) $\angle EMG$

b) $\angle ECM$


c) $\angle ECG$



Student
view



Overview
(/study/ap
aa-
hl/sid-
134-
cid-
761926/o



Interactive 2. Square-Based Right Pyramid ABCDE.

More information for interactive 2

This interactive tool provides an engaging and dynamic way to explore the geometric properties of a **square-based right pyramid ABCDE**. The pyramid has a **height of 6 cm**, and each side of its square base measures 10 cm. The midpoint of the base edge **CD** is labeled as **M**, serving as a key reference point for angle calculations.

The interactivity allows for **rotation of the 3D pyramid**, offering multiple perspectives to better understand its structure. Additionally, **points A and B can be moved**, enabling adjustments to the base dimensions. As the base expands or compresses, the internal angles of the pyramid change, demonstrating how different elements of the shape interact.

This tool focuses on guiding through the calculation of three important angles:

- $\angle E M G$: This angle is part of the right-angled triangle **EGM**, where **G** is the foot of the **perpendicular from E to the base**. Since this triangle is right-angled at **G**, trigonometry can be used to determine the other angles.
- $\angle E C M$: This angle lies within the triangular face **ECM**, which is formed by the slant height of the pyramid and a segment of the base.
- $\angle E C G$: This angle illustrates the relationship between the apex and different sections of the base.

By allowing direct manipulation of the pyramid's structure, this tool provides an interactive experience



Student
view




Overview
(/study/ap
aa-
hl/sid-
134-
cid-
761926/o

that enhances the understanding of **3D geometry, trigonometry, and spatial relationships**. The ability to rotate the shape and modify its dimensions makes it easier to explore geometric concepts in a more intuitive and accessible way.

To calculate $\angle EMG$, isolate triangle EGM, which is right-angled at G. This angle may also be described as the angle between EM and the base ABCD.

To calculate $\angle ECM$, isolate triangle EMC, which is right-angled at M.

To calculate $\angle ECG$, isolate triangle EGC, which is right-angled at G. This angle may also be described as the angle between EC and the base.



a) Let $\angle EMG = \theta$.



Student
view

So $\tan \theta = \frac{6}{5}$.




Overview
(/study/ap)

aa-
hl/sid-
134-
cid-
761926/o

Therefore, $\theta = \tan^{-1} \frac{6}{5} = 50.2^\circ$ (to 3 significant figures).

In triangle EMG you can also find the length of [EM], which will be used later.

$$EM = \sqrt{5^2 + 6^2} = \sqrt{61}$$



b) Let $\angle ECM = \alpha$.

$$\text{So, } \tan \alpha = \frac{\sqrt{61}}{5}.$$

Therefore, $\alpha = \tan^{-1} \frac{\sqrt{61}}{5} = 57.4^\circ$ (to 3 significant figures).

In triangle EMC you can also find the length of [EC], which will be used later.




Student
view



Overview
(/study/ap)

aa-
hl/sid-
134-
cid-
761926/o

$$EC = \sqrt{5^2 + \sqrt{61}^2} = \sqrt{86}$$



c) Relative to the angle marked β (which is $\angle ECG$), you know the opposite side and the hypotenuse of the triangle. So you can use the sine ratio:

$$\sin \beta = \frac{6}{\sqrt{86}}$$

Therefore, $\beta = \sin^{-1} \frac{6}{\sqrt{86}} = 40.3^\circ$ (to 3 significant figures).

You can also form right-angled triangles in cuboids and use these to find the angles between lines.

3 section questions ^




Student
view

Question 1



🏠
 Overview
 (/study/ap
 aa-
 hl/sid-
 134-
 cid-
 761926/o

The following diagram shows a pyramid ABCD, where the base ABC is an equilateral triangle. The point marked E is directly below the apex.



ⓘ More information

Which right-angled triangle should you use to calculate the angle between DA and the base?

1 ΔDEA



2 ΔEAC

3 ΔDAC

4 ΔDAB

Explanation

To find the angle between the line DA and the base, you drop a line from the point D to intersect the base at 90° . This is the line DE.

You can now identify the angle between the line and the base as the angle $\angle DAE$.

The triangle ΔDEA is right-angled at E, and can be used to calculate the angle $\angle DAE$.




Student
view



Overview
 (/study/ap
 aa-
 hl/sid-
 134-
 cid-
 761926/o

Question 2

The right pyramid ABCD has base ABC which is an equilateral triangle. E is directly below the point D.



More information

Calculate the angle AEC.

1 120°

2 60°

3 45°

4 90°

Explanation


Here is the pyramid from above:



Student
view



Overview
 (/study/app/math-aa-hl/sid-134-cid-761926/o)
 aa-
 hl/sid-
 134-
 cid-
 761926/o



More information


Note that the three angles α , β and γ are all equal and together sum to 360° . Therefore, $\gamma = \angle AEC = 120^\circ$.

Question 3



The square-based pyramid ABCDE is shown below.

The height is 12 cm. The length of each side of the base is 8 cm.



Student
view

More information

Find the angle between EA and the base ABCD.



Overview
(/study/ap
aa-
hl/sid-
134-
cid-
761926/o

1 64.8°



2 45°


3 71.6°

4 72.5°

Explanation

Step 1: The triangle required is EHA. However, only one side is currently known, $EH = 12 \text{ cm}$.

Step 2: Calculate the length of AH.



More information

From the diagram, $AH = \sqrt{4^2 + 4^2} = \sqrt{32}$.


Step 3: Now you are ready to calculate the angle.



Student
view



Overview
 (/study/app/
 aa-
 hl/sid-
 134-
 cid-
 761926/o)



More information

$$\tan \theta = \left(\frac{12}{\sqrt{32}} \right)$$

$$\begin{aligned} \theta &= \tan^{-1} \left(\frac{12}{\sqrt{32}} \right) \\ &= 64.760\dots^\circ \end{aligned}$$

So the angle between EA and the base ABCD is 64.8° (3 significant figures).

3. Geometry and trigonometry / 3.1 Three-dimensional space

Checklist

Section

Student... (0/0)



Feedback



Print (/study/app/math-aa-hl/sid-134-cid-761926/book/checklist-id-25415/print/)

Assign



What you should know

By the end of this subtopic you should be able to:

- find the midpoint and the distance of any two points in 3D
- select the correct formulae for finding areas and volumes of 3D shapes from the formula booklet
- count the faces on a pyramid or prism



Student
view



Overview
(/study/app/math-aa-hl/sid-134-cid-761926/o)

- calculate the total surface area of a solid by adding together the area of each of its faces
- calculate the volume of pyramids, cones, spheres and combinations of these solids.

3. Geometry and trigonometry / 3.1 Three-dimensional space

Investigation

Section

Student... (0/0)

Feedback

Print (/study/app/math-aa-hl/sid-134-cid-761926/book/investigation-id-25416/print/)

Assign

As you saw at the start of this subtopic, many geometric shapes form the bases of architectural designs. Here you have two examples from two different cities in the world.

The first one is the Petronas twin towers, Kuala Lumpur, Malaysia and the second one is the Kuwait towers of Kuwait City.



Petronas twin towers, Kuala Lumpur, Malaysia

Student view

Credit: fotoVoyager GettyImages



Overview
(/study/ap
aa-
hl/sid-
134-
cid-
761926/o



Kuwait towers, Kuwait City, Kuwait

Credit: Anson Fernandez Dionisio GettyImages

Your task is to model one of these two pairs of towers to estimate the height, surface area and volume of the building.

Consider the following questions to complete this task.

- What are the relevant elements? What information do you need to solve this problem?
- What mathematical strategies will you use to solve the problem? What mathematics do you need to do? What technology will you use to help you?
- Do your solutions make sense in real-life?
- Consider the accuracy of your solutions – justify the accuracy and explain any reasons why inaccuracy exists.
- Write a report showing how you solved the problem. Include reasoning and justification for all your decisions and the final solutions, as well as reflection.



Student
view



Overview
(/study/ap

aa-
hl/sid-
134-
cid-
761926/o

Rate subtopic 3.1 Three-dimensional space

Help us improve the content and user experience.



Student
view