

Overview  
(/study/apaa-  
hl/sid-  
134-  
cid-  
761926/o

Teacher view



(https://intercom.help/kognity)

**Index**

The big picture  
Circular functions revisit  
Symmetries  
Checklist  
Investigation

Table of  
contents

Notebook




Glossary

Reading  
assistance

3. Geometry and trigonometry / 3.11 Further circular functions

# The big picture

In previous subtopics, you studied properties of trigonometric functions and their graphs. In this subtopic you will have a closer look at the graphs of trigonometric functions and their relationships.



More information

Student  
view

The image is a graph displaying the trigonometric functions sine, cosine, and tangent. The X-axis represents the angle in radians while the Y-axis represents the function values. The range of the Y-axis is from  $-2$  to  $2$ .



Overview  
 (/study/ap  
 aa-  
 hl/sid-  
 134-  
 cid-  
 761926/o

1.  **$g(x) = \cos(x)$** : This is represented by a magenta curve that oscillates between -1 and 1. It starts from the maximum value of 1 when  $x = 0$ .
2.  **$f(x) = \sin(x)$** : This function is shown as a blue curve that also oscillates between -1 and 1. It starts from 0 when  $x = 0$  and reaches its peaks (1) and troughs (-1) as  $x$  increases.
3.  **$h(x) = \tan(x)$** : This is represented by a cyan curve. The tangent function has vertical asymptotes where the function is undefined, which occurs at  $x = \pi/2, 3\pi/2$ , etc., and it crosses the x-axis at multiples of  $\pi$ .

The graph also includes vertical lines indicating key angles, such as  $x = \pi/2$ . There are gridlines for easy determination of values.

[Generated by AI]



## Concept

Trigonometric functions allow you to model cyclic phenomena.

The study of trigonometric functions provides you with the tools to analyse, measure and transform the quantities, movements and relationships of cyclic (periodic) functions.

3. Geometry and trigonometry / 3.11 Further circular functions

# Circular functions revisit




Student  
view



Overview  
(/study/app/  
math-aa-hl/  
sid-134-  
cid-761926/o

In subtopic 3.5 (/study/app/math-aa-hl/sid-134-cid-761926/book/the-big-picture-id-27741/), you studied the unit circle and trigonometric ratios beyond acute angles. The symmetry of the unit circle helped you to find trigonometric ratios of any angle using its relationship to the principal angle in the first quadrant.

## Making connections



More information

This image is a diagram illustrating the unit circle, a central concept in trigonometry. The circle is centered at the origin  $(0,0)$  on a Cartesian coordinate plane, with both  $x$  and  $y$  axes clearly labeled. Various points on the circle are labeled with their trigonometric identities, including  $(1,0)$ ,  $(0,1)$ ,  $(-1,0)$ , and  $(0,-1)$ . Additional points are labeled with expressions of cosine and sine functions:  $(\cos \theta, \sin \theta)$ ,  $(-\cos \theta, \sin \theta)$ ,  $(-\cos \theta, -\sin \theta)$ , and  $(\cos \theta, -\sin \theta)$ . The diagram illustrates angles  $\theta$  in standard position, originating from the center of the circle and sweeping outwards. Dotted lines indicate the triangle formed by the radius, helping to visualize sine and cosine as coordinates. The central angle  $\theta$  is shown multiple times inside the circle. The radius of the circle is consistently marked as 1, which is the defining feature of the unit circle.

[Generated by AI]  
Student... (0/0)

Feedback

Print (/study/app/math-aa-hl/sid-

Assign

Section



Student  
view

134-cid-761926/book/the-big-picture-id-27915/print/)



Overview

(/study/ap

aa-

hl/sid-

134-

cid-

761926/o

In [subtopic 3.5 \(/study/app/math-aa-hl/sid-134-cid-761926/book/the-big-picture-id-27741/\)](#), you studied the unit circle and used the following identities


$$\cos(\pi - \theta) = -\cos \theta$$

$$\sin(\pi - \theta) = \sin \theta$$

$$\tan(\pi - \theta) = -\tan \theta$$

Although  $f(x) = \sin x$  and  $h(x) = \sin(\pi - x)$  are two different functions, when you graph them they would look exactly the same because  $\sin(\pi - x) = \sin x$ .

Consider the two functions  $f(x) = \sin x$  and  $q(x) = \sin(\pi + x)$ , which are shown in the diagram below. The graph of  $y = q(x)$  is the reflection of  $y = f(x)$  in the  $y$ -axis.



More information

The graph displays the functions  $f(x) = \sin x$  and  $q(x) = \sin(\pi - x)$ . The X-axis represents values ranging from  $-(2\pi)$  to  $(2\pi)$ , labeled at intervals of  $(\pi/2)$ ,  $(\pi)$ , and  $(2\pi)$ . The Y-axis ranges from -1 to 1.

The blue curve represents  $f(x) = \sin x$ , exhibiting a typical sinusoidal wave starting at  $(0,0)$ , peaking at  $(\pi/2, 1)$ , crossing zero at  $(\pi)$ , troughing at  $(3\pi/2, -1)$ , and completing the cycle at  $(2\pi, 0)$ .

Student  
view



Overview

(/study/ap

aa-

hl/sid-

134-

cid-

761926/o

The red curve corresponds to  $q(x) = \sin(\pi - x)$ , which is a reflection of the blue curve across the Y-axis. It begins at (0,0), peaks down to (-1) at  $(\pi/2, -1)$ , returns to zero at  $(\pi, 0)$ , peaks up to (1) at  $(3\pi/2, 1)$ , and then returns to zero at  $(2\pi, 0)$ .

Both curves overlay such that as one descends, the other ascends, showcasing their reflectional relationship across the Y-axis.

[Generated by AI]

## Example 1



Prove that  $\cos\left(\frac{3\pi}{2} + x\right) = \sin x$ .

Use the compound angle identity for cosine:

$$\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta$$

$$\cos\left(\frac{3\pi}{2} + x\right) = \cos \frac{3\pi}{2} \cos x - \sin \frac{3\pi}{2} \sin x$$

$$\cos \frac{3\pi}{2} = 0 \text{ and } \sin \frac{3\pi}{2} = -1$$

$$\cos\left(\frac{3\pi}{2} + x\right) = 0 \times \cos x - (-1) \sin x$$

Simplify

$$\cos\left(\frac{3\pi}{2} + x\right) = \sin x$$

Therefore,



Student  
view




Overview  
 (/study/ap  
 aa-  
 hl/sid-  
 134-  
 cid-  
 761926/o

$$\cos\left(\frac{3\pi}{2} + x\right) = \sin x$$

## Example 2



Sketch the graph of  $y = f(x)$  where  $f(x) = \tan x$  for  $0 \leq x \leq 2\pi$ . Sketch the graph of  $y = g(x)$ , where  $g(x) = f(\pi - x)$ . Describe the relationship between the graphs of  $y = f(x)$  and  $y = g(x)$ .

Steps	Explanation
	<p>Sketch the graph for the domain making sure that vertical asymptotes and are marked clearly.</p>




Student  
view



Overview  
(/study/ap  
aa-  
hl/sid-  
134-  
cid-  
761926/o

## Steps

## Explanation



Use the result that

$$\tan(\pi - x) =$$

to sketch the graph of  $y$



The graph of  $y = g(x)$  is the reflection of  $y = f(x)$  in the  $x$ -axis.

### ⚠ Be aware

In IB examinations, if the question starts with ‘hence’ it means you have to use the results from previous work you have done. If you do not do so, even if your alternative method is correct it will not be accepted.

If the question starts with ‘hence’ followed by ‘or otherwise’ then you can use an alternative method.

## Example 3




Student  
view

Sketch the graph of  $y = f(\theta)$  where  $f(\theta) = \sin(\pi + 2\theta)$ ,  $0 \leq \theta \leq \pi$ .



Overview  
(/study/ap  
aa-  
hl/sid-  
134-  
cid-  
761926/o

Hence, sketch the graph of  $g(\theta) = \operatorname{cosec}(\pi + 2\theta)$ , clearly marking the coordinates of any local maximum/minimum points and the equations of any asymptotes.

Steps	Explanation
	<p><math>\sin(\pi + 2\theta) = -\sin 2\theta</math></p> <p>which has period of <math>\pi</math>. The local minimum at <math>A\left(\frac{\pi}{4}, -1\right)</math>, local maximum at <math>B\left(\frac{3\pi}{4}, 1\right)</math></p> <p><math>x</math>-intercepts are <math>(0, 0)</math>, <math>(\pi, 0)</math>.</p>



Student  
view






Overview  
(/study/ap  
aa-  
hl/sid-  
134-  
cid-  
761926/o

## Steps

## Explanation



As  $\operatorname{cosec}(\pi + 2\theta) = \frac{1}{\sin(\pi + 2\theta)}$

its graph will have asymptotes where  $f(\theta) = 0$

$y = g(\theta)$  will have a local maximum when  $f(\theta)$  has a minimum

$y = g(\theta)$  will have a local minimum when  $f(\theta)$  has a maximum

Vertical asymptotes of  $y = g(\theta)$  are  $x = 0$ ,  $x = \frac{\pi}{2}$  and  $x = \pi$

The local maximum of  $y = g(\theta)$  is at  $B\left(\frac{3\pi}{4}, 1\right)$  and the local minimum is at  $A\left(\frac{\pi}{4}, -1\right)$ .

## Example 4



Determine whether  $f(x) = \cos x^2 - x$  is even, odd or neither.



Student  
view



Overview

(/study/ap

aa-

hl/sid-

134-

cid-

761926/o

$$f(-x) = \cos(-x)^2 - (-x)$$

$$f(-x) = \cos x^2 + x$$

$$f(-x) \neq -f(x)$$

And

$$f(-x) \neq f(x)$$

Therefore,  $f(-x)$  is neither odd nor even.

### ! Exam tip

In IB examinations, if you are asked to sketch a graph in multiple steps, do not erase your work to show your final answer. You should label your final answer clearly making sure all the critical points like maxima, minima, axes intercepts and asymptotes are marked and their values or equation are written down.

## 2 section questions ^

### Question 1



★☆☆

Sketch the graph of  $y = f(x)$ , where  $f(x) = \cos(\pi - x)$ , for  $-\pi \leq x \leq \pi$ .

Hence, find the number of solutions to the equation  $f(x) = 0.2$ .

Two solutions




Student  
view




Overview  
(/study/ap  
aa-  
hl/sid-  
134-  
cid-  
761926/o

1



More information

Two solutions



More information

Two solutions




Student  
view



Overview  
(/study/ap  
aa-  
hl/sid-  
134-  
cid-  
761926/o


3



More information

Four solutions

4



More information

### Explanation

$$\cos(\pi - x) = -\cos x$$




Student  
view

Sketch the graph, draw the line  $y = 0.2$  and mark the intersections with the curve.



Overview  
(/study/ap  
aa-  
hl/sid-  
134-  
cid-  
761926/o



More information

Therefore there are two solutions in the given domain.

## Question 2



★★☆

Let  $f(x) = \tan x$ ,  $-\frac{3\pi}{2} < x < \frac{-\pi}{2}$ . The graph of  $y = f(x)$  is reflected in the x-axis and then translated by the vector  $\begin{pmatrix} \pi \\ 1 \end{pmatrix}$ .

Which of the following represents the graph and the equation of the transformed function?




Student  
view



Overview  
(/study/ap  
aa-  
hl/sid-  
134-  
cid-  
761926/o

1



👁 More information

$$h(x) = -\tan x + 1$$




Student  
view



Overview  
 (/study/ap  
 aa-  
 hl/sid-  
 134-  
 cid-  
 761926/o


2



🔍 More information

$$h(x) = \tan x + 1$$

3



🔍 More information



Student  
view



Overview

(/study/ap

aa-


hl/sid-

134-

cid-

761926/o

$$h(x) = -\tan x - 1$$



4

More information

$$h(x) = -\tan x - 1$$

### Explanation

Reflection in x-axis:  $g(x) = -f(x)$

This reflection does not change the domain of the graph.

Translation by vector  $\begin{pmatrix} \pi \\ 1 \end{pmatrix}$ :  $h(x) = g(x - \pi) + 1$

### Section

Student (0/0)

Feedback

Print

Assign

This translation changes the domain of the graph from  $-\frac{\pi}{2} < x < \frac{\pi}{2}$  to  $\frac{\pi}{2} < x < \frac{3\pi}{2}$ , so all

answer options are graphed on the correct domain.

Moreover,

$$h(x) = -f(x - \pi) + 1 = -\tan(x - \pi) + 1 = -\tan x + 1.$$

So the correct answer is




Student  
view





Overview  
 (/study/ap  
 aa-  
 hl/sid-  
 134-  
 cid-  
 761926/o



🔍 More information

3. Geometry and trigonometry / 3.11 Further circular functions

## Symmetries

As trigonometric functions are periodic, you can find many symmetries in each graph.

In this section, you will look into symmetries in relation to even and odd functions.

### 🔗 Making connections

In [subtopic 2.14 \(/study/app/math-aa-hl/sid-134-cid-761926/book/the-big-picture-id-26756/\)](#), you studied even and odd functions.


A function  $f(x)$  is odd if  $f(-x) = -f(x)$  for each  $x$  in the given domain.



Student  
view



Overview  
 (/study/ap  
 aa-  
 hl/sid-  
 134-  
 cid-  
 761926/o



More information

The image shows a graph representing an odd function. The graph is plotted on a Cartesian coordinate system with  $x$  and  $y$  axes. The  $x$ -axis is labeled from negative to positive, and similarly, the  $y$ -axis is labeled. The function intersects the origin  $(0,0)$  and is symmetrical about this point, demonstrating the property of an odd function where  $f(-x) = -f(x)$ . Points are marked on the graph: one on the negative side of the  $x$ -axis labeled ' $-x$ ' and a corresponding  $y$  value, and one on the positive side of the  $x$ -axis labeled ' $x$ ' and its corresponding negative  $y$  value, emphasizing the function's symmetry. Two points are marked on the  $y$ -axis, indicating specific values at ' $a$ ' and ' $-a$ '.

[Generated by AI]


A function  $f(x)$  is even if  $f(-x) = f(x)$  for each  $x$  in the given domain.



Student  
view



Overview  
 (/study/ap  
 aa-  
 hl/sid-  
 134-  
 cid-  
 761926/o



More information

A graph depicts an even function demonstrating symmetry about the y-axis. The x-axis and y-axis are clearly labeled. The curve of the function is a parabola, opening upwards, with its vertex at the origin (0, 0). The y-value at the point labeled "a" remains the same for both  $x = -x$  and  $x = x$ , illustrating the concept of even functions, where  $f(x) = f(-x)$ . The parabola is symmetric, meaning that for every point on the curve at  $x$ , there is a corresponding point at  $-x$  with the same y-value.

[Generated by AI]



## Activity

In this activity, you will investigate whether trigonometric functions are even or odd.

For each of the trigonometric functions

$$y = \sin x, y = \cos x, y = \tan x, y = \sec x \text{ and } y = \operatorname{cosec} x$$

1. determine whether it is even, odd or neither



Student  
view



Overview

(/study/ap

aa-

hl/sid-


134-

cid-

761926/o

## 2. by sketching their graphs, describe the symmetries you observe.

The sine graph has rotational symmetry with respect to the origin, as you can see in the applet below, each point on the right-hand side of the origin is rotated  $\pi$  radians.



### Interactive 1. Rotational Symmetry of the Sine Graph.

Credit: [GeoGebra](https://www.geogebra.org/m/vezpm7qt) (https://www.geogebra.org/m/vezpm7qt) Nuriye Sirinoglu Singh

More information for interactive 1

This interactive explores the rotational symmetry of the sine function about the origin. The sine curve maintains symmetrical relationships when rotated by  $\pi$  radians, demonstrating the function's odd symmetry property through visual representation and dynamic point manipulation.

The display shows a graph with xy axes, with x-axis measured in radians ranging from -2 to 2 And y-axis ranging from -2 to 1. A purple sine graph, is centered at (0,0), representing the function  $f(x) = \sin(x)$ . Two points A (movable) and B (automatically updated) appear on the curve, connected by a line, with their coordinates visible. Point A's position can be adjusted along the sine wave.

For example, as users drag point A to  $(\pi/2, 1)$ , point B automatically moves to  $(-\pi/2, -1)$ , showing the sign change. When A is placed at  $(\pi, 0)$ , B moves to  $(-\pi, 0)$ , maintaining the origin symmetry. These movements visually confirm the mathematical relationship  $f(-x) = -f(x)$  for sine.

Through this exploration, users gain concrete understanding of odd functions' rotational symmetry, seeing how sine's properties manifest graphically. They learn to predict symmetrical point pairs and verify the function's behavior algebraically through coordinate observation, reinforcing fundamental trigonometric concepts essential for advanced mathematics and physics applications.



Student  
view



Overview

(/study/ap

aa-

hl/sid-

134-

cid-

761926/o

## ✓ Important

The graph of an odd function is symmetric with respect to the origin (rotational symmetry) and the graph of an even function is symmetric with respect to the  $y$ -axis (line symmetry).

## Example 1



Sketch the graph of  $y = \cos 2x$  for  $-\pi \leq x \leq \pi$ .

Given that  $\cos 2x = 0.8$ , use your sketch to find

$$\cos(-2\pi + 2x) \quad \cos(-2x) \quad \cos(2\pi - 2x).$$




Student  
view




Overview  
(/study/ap  
aa-  
hl/sid-  
134-  
cid-  
761926/o

## Steps

## Explanation



Sketch the graph of  $y = \cos(2x)$  marking the end points, and minima.




Draw the line  $y = 0.8$  and points of intersection with  $y = \cos(2x)$ .

Use the symmetry of the graph to write down the relationship between the coordinates of the point of intersection.



Student  
view

  
Overview  
(/study/ap  
aa-  
hl/sid-  
134-  
cid-  
761926/o


Steps	Explanation
<p>Therefore:</p> $\cos (-2\pi + 2x) = 0.8$ $\cos (-2x) = 0.8$ $\cos (2\pi - 2x) = 0.8$	

Example 2



Let  $m = \cos n, 0 < n < \frac{\pi}{2}$


Find the solution(s) of  $\cos 2x = -m, 0 \leq x \leq \pi$ , in terms of  $n$ .

Steps	Explanation
$\cos 2x = -\cos n$	<p>Use the symmetry properties of the cosine</p> 

  
Student  
view



Overview  
(/study/ap  
aa-  
hl/sid-  
134-  
cid-  
761926/o

Steps	Explanation
$\cos 2x = \cos (\pi - n)$ or $\cos 2x = \cos (\pi + n)$	<p>Use the symmetry properties of the cosine</p> 
$2x = \pi - n \Rightarrow x = \frac{\pi}{2} - \frac{n}{2}$ or $2x = \pi + n \Rightarrow x = \frac{\pi}{2} + \frac{n}{2}$	Solve the equations.
<p>Therefore, the answer is</p> $x = \frac{\pi}{2} - \frac{n}{2} \text{ or } x = \frac{\pi}{2} + \frac{n}{2}$	<p>Both results satisfy the initial condition</p> $0 \leq x \leq \pi$

### ❗ Exam tip

In the IB examination, you will not be told whether a trigonometric function is even or odd, nor will you be given information about the symmetry. You should memorise that, in a given domain,



Student  
view





Overview

(/study/ap

aa-

hl/sid-

134-

cid-

761926/o

odd functions have the property that  $f(-x) = -f(x)$ , i.e. they have rotational symmetry about the origin

even functions have the property that  $f(-x) = f(x)$ , i.e. they have line symmetry (with respect to the  $y$ -axis)

and be able to sketch the graphs to see the symmetries.



## International Mindedness

Each year, as many as 10 000 containers fall off the ships which carry goods across the seas. In 1992, a container full of rubber ducks, which were going to Seattle from China, fell off the ship. After almost a decade they appeared off the coast of New England having travelled the oceans around the globe.

Along with these containers, millions of tons of plastic end up in the oceans, so polluting the water and threatening the sea life.

Understanding currents and tides could support cleaning efforts. In this article

(<https://plus.maths.org/content/os/issue32/features/elwell/index#ref1>),

Frances Elwell discusses how currents and tides behave and how studying these could help to protect the environment.

## 3 section questions ^

### Question 1



Select, from the given options, which of the following is the graph of an even function for the domain  $-\frac{\pi}{2} \leq x \leq \frac{\pi}{2}$ .




Student  
view




Overview  
(/study/ap  
aa-  
hl/sid-  
134-  
cid-  
761926/o

1



More information

2



More information




Student  
view


3



Overview  
 (/study/ap  
 aa-  
 hl/sid-  
 134-  
 cid-  
 761926/o



👁 More information



👁 More information

### Explanation


Even functions have line symmetry with respect to the  $y$ -axis. Therefore, the correct answer is



Student  
view



Overview  
(/study/ap  
aa-  
hl/sid-  
134-  
cid-  
761926/o



More information

## Question 2



★☆☆

Select, from the given options, an odd function in the domain  $-\frac{\pi}{2} \leq x \leq \frac{\pi}{2}$ .

1  $y = \cos x \sin x$



2  $y = \cos x + \sin x$

3  $y = 2 \cos x + \sin x$

4  $y = 2 \cos x + 3 \sin x$

## Explanation

Odd functions have the property that  $f(-x) = -f(x)$

$$\cos(-x) \sin(-x) = \cos x (-\sin x) = -\cos x \sin x$$



Therefore, the correct answer is

$$y = \cos x \sin x$$

Student  
view



Overview  
 (/study/app/  
 math-aa-  
 hl/sid-  
 134-  
 cid-  
 761926/o

**Question 3**

★☆☆

Which of the following is an even function in the domain  $-\pi \leq x \leq \pi$ ?

1  $y = \sec 2x$



2  $y = \tan 2x$

3  $y = \operatorname{cosec} 2x$

4  $y = \sin 2x$

**Explanation**

Even functions have the property that  $f(-x) = f(x)$

$$\sec(-2x) = \frac{1}{\cos(-2x)} = \frac{1}{\cos 2x} = \sec 2x$$

Therefore, the correct answer is  $y = \sec 2x$ .

3. Geometry and trigonometry / 3.11 Further circular functions

# Checklist

**Section**

Student... (0/0)



Feedback



Print (/study/app/math-aa-hl/sid-

134-cid-761926/book/checklist-id-  
27918/print/)

**Assign****What you should know**

By the end of this subtopic you should be able to:

- identify whether a trigonometric function is odd, even or neither



Student  
view



Overview

(/study/app

aa-

hl/sid-

134-

cid-

761926/o

- use properties of even and odd functions to solve equations involving trigonometry.

3. Geometry and trigonometry / 3.11 Further circular functions

# Investigation

Section

Student... (0/0)



Feedback



Print (/study/app/math-aa-hl/sid-

Assign

134-cid-761926/book/investigation-id-

27919/print/)

You will investigate the relationship between  $y = \sin x$  and  $y = \sin \left( \frac{\pi}{2} \pm x \right)$ .

Use Geogebra and draw the graphs of  $y = \sin x$ ,  $y = \sin \left( \frac{\pi}{2} + x \right)$  and  $y = \sin \left( \frac{\pi}{2} - x \right)$ .

Describe the relationship between graphs.

Use compound angle identities to prove these relationships.

## Rate subtopic 3.11 Further circular functions

Help us improve the content and user experience.

Student  
view