Individual Assignment 5

Zelda Reif

2026-05-10

1. Set Up

Packages

```
# general use
library(tidyverse)
library(here)
library(janitor)
library(snakecase)

# wrap axis labels
library(scales)

# reading .xlsx files
library(readxl)

# visualize missing data
library(naniar)

# arrange plots
library(patchwork)
```

Data

```
# taxonomic info
taxon_list <- read_csv(here("data","taxon_list.csv"))
```

```r
# aquatic invertebrates
aq_ins <- read_xlsx(
  here("data", "Aquatic Sampling Data-2026-03-10.xlsx"),
  sheet = "Aquatic Insects"
)

# site info
sites <- read_xlsx(
   here("data", "Aquatic Sampling Data-2026-03-10.xlsx"),
   sheet = "sites"
)

# water quality
water_quality <- read_xlsx(
  here("data", "Aquatic Sampling Data-2026-03-10.xlsx"),
  sheet = "Water Quality",
  na = c("", "n/a", "N/A", "over", "173+")
)
```

2. Cleaning

`sites` object

```r
# creating new object
ncos_sites <- sites |>
  # filtering to only include ncos quarterly sampling sites
  filter(sector == "NCOS" & sampling_frequency == "Quarterly")
```

clean `taxon_list`

```r
taxon_list_clean <- taxon_list |>
  # convert taxon names to snakecase
  mutate(verbatim_name = to_any_case(verbatim_name,
                                     case = "snake"))
```

clean `water_quality`

```
water_quality_clean <- water_quality |>
  clean_names() |>
  semi_join(ncos_sites, by = "site") |>
  unite("date_site", date, site, remove = FALSE) |>
  group_by(date, site, date_site) |>
  summarize(med_ph = median(p_h, na.rm = TRUE),
            med_sal = median(salinity_ppt, na.rm = TRUE),
            med_do = median(dissolved_oxygen_mg_l, na.rm = TRUE)) |>
  ungroup() |>
  filter(date_site != "2022-08-12_NVBR")
```

clean `aq_ins`

```
# creating new clean object from aquatic inverts
aq_ins_clean <- aq_ins |>
  # clean column names
  clean_names() |>
  # filtering join: only include sites occurring in ncos_sites
  semi_join(ncos_sites, by = c("site")) |>
  # renaming columns
  rename(date = date_on_vial) |>
  # select columns of interest
  select(date, sample_type, site,
         ostracod,
         copepod,
         hemiptera_corixidae_boatman,
         cladocera,
         nematode,
         diptera_ceratopogonidae,
         annelida_oligochaete,
         diptera_chironomid,
         annelida_polychaete) |>
  # filter to only include "filtered beaker" samples
  filter(sample_type %in% c("FB 250", "FB250")) |>
  # convert the data frame to long format
  pivot_longer(cols = ostracod:annelida_polychaete,
               names_to = "taxon",
               values_to = "abundance") |>
```

```
  # group by date, site, taxon
  group_by(date, site, taxon) |>
  # calculate average abundance per liter (sum abundance divided by 7.5
   ↪  liters)
  summarize(ave_lit = sum(abundance, na.rm = TRUE)/7.5) |>
  # ungroup data frame
  ungroup() |>
  # create a new column called `date_site` to join with water quality
  unite("date_site", date, site, remove = FALSE) |>
  # join with water quality
  left_join(
  select(water_quality_clean,
         date_site,
         med_ph,
         med_sal,
         med_do),
  by = "date_site"
) |>
  # join with taxon list
  left_join(taxon_list_clean, by = c("taxon" = "verbatim_name")) |>
  # add full names of sites
  mutate(site_full = case_when(
    site == "NVBR" ~ "North of Venoco Bridge",
    site == "NEC" ~ "South of Venoco Bridge",
    site == "NMC" ~ "Main Channel",
    site == "NPB" ~ "South of Phelps Bridge",
    site == "NPB1" ~ "North of Phelps Bridge",
    site == "NPB2" ~ "Phelps Road",
    site == "NWP" ~  "West Pond",
    site == "NDC" ~ "Devereux Creek"
  )) |>
  # setting factor levels for site
  mutate(site_full = fct_reorder(site_full, med_sal, .fun = "median", .na_rm
   ↪  = TRUE),
         site_full = fct_rev(site_full))
```

Problems

1. Visualizing differences across sites in major taxon abundance

a. Plan your figure

x-axis: site y-axis: ostracod abundance geometry to represent average abundance/L: `geom_jitter` geometry to represent medians: `stat_summary()`

b. Wrangle your data

c. Code your figure

d. Create a multipanel figure

e. Interpret your figure

differences between sites in copepod and ostracod abundance how ostracod abundance may relate (or not) to site-level differences in salinity

2. Visualizing total abundance as a function of salinity

a. Plan your figure

x-axis: y-axis: color: geometry: facets:

b. Wrangle your data

c. Code your figure

d. Interpret your figure

In 3-5 sentences, describe differences or similarities between taxa in relationships between abundance and salinity. Be specific using the visual components of the plot.