

Individual Assignment 5

Zelda Reif

2026-05-10

1. Set Up

Packages

```
# general use
library(tidyverse)
library(here)
library(janitor)
library(snakecase)

# wrap axis labels
library(scales)

# reading .xlsx files
library(readxl)

# visualize missing data
library(naniar)

# arrange plots
library(patchwork)
```

Data

```
# taxonomic info
taxon_list <- read_csv(here("data","taxon_list.csv"))
```

```

# aquatic invertebrates
aq_ins <- read_xlsx(
  here("data", "Aquatic Sampling Data-2026-03-10.xlsx"),
  sheet = "Aquatic Insects"
)

# site info
sites <- read_xlsx(
  here("data", "Aquatic Sampling Data-2026-03-10.xlsx"),
  sheet = "sites"
)

# water quality
water_quality <- read_xlsx(
  here("data", "Aquatic Sampling Data-2026-03-10.xlsx"),
  sheet = "Water Quality",
  na = c("", "n/a", "N/A", "over", "173+")
)

```

2. Cleaning

sites object

```

# creating new object
ncos_sites <- sites |>
  # filtering to only include ncos quarterly sampling sites
  filter(sector == "NCOS" & sampling_frequency == "Quarterly")

```

clean taxon_list

```

taxon_list_clean <- taxon_list |>
  # convert taxon names to snakecase
  mutate(verbatim_name = to_any_case(verbatim_name,
    case = "snake"))

```

clean water_quality

```
water_quality_clean <- water_quality |>
  clean_names() |>
  semi_join(ncos_sites, by = "site") |>
  unite("date_site", date, site, remove = FALSE) |>
  group_by(date, site, date_site) |>
  summarize(
    med_ph = median(p_h, na.rm = TRUE),
    med_sal = median(salinity_ppt, na.rm = TRUE),
    med_do = median(dissolved_oxygen_mg_l, na.rm = TRUE)) |>
  ungroup() |>
  filter(date_site != "2022-08-12_NVBR")
```

clean aq_ins

```
# creating new clean object from aquatic inverts
aq_ins_clean <- aq_ins |>
  # clean column names
  clean_names() |>
  # filtering join: only include sites occurring in ncos_sites
  semi_join(ncos_sites, by = c("site")) |>
  # renaming columns
  rename(date = date_on_vial) |>
  # select columns of interest
  select(
    date, sample_type, site,
    ostracod,
    copepod,
    hemiptera_corixidae_boatman,
    cladocera,
    nematode,
    diptera_ceratopogonidae,
    annelida_oligochaete,
    diptera_chironomid,
    annelida_polychaete) |>
  # filter to only include "filtered beaker" samples
  filter(sample_type %in% c("FB 250", "FB250")) |>
  # convert the data frame to long format
  pivot_longer(
    cols = ostracod:annelida_polychaete,
    names_to = "taxon",
    values_to = "abundance") |>
```

```

# group by date, site, taxon
group_by(date, site, taxon) |>
# calculate average abundance per liter (sum abundance divided by 7.5
  ↳ liters)
summarize(ave_lit = sum(abundance, na.rm = TRUE)/7.5) |>
# ungroup data frame
ungroup() |>
# create a new column called `date_site` to join with water quality
unite("date_site", date, site, remove = FALSE) |>
# join with water quality
left_join(
  select(water_quality_clean,
        date_site,
        med_ph,
        med_sal,
        med_do),
  by = "date_site"
) |>
# join with taxon list
left_join(taxon_list_clean, by = c("taxon" = "verbatim_name")) |>
# add full names of sites
mutate(site_full = case_when(
  site == "NVBR" ~ "North of Venoco Bridge",
  site == "NEC" ~ "South of Venoco Bridge",
  site == "NMC" ~ "Main Channel",
  site == "NPB" ~ "South of Phelps Bridge",
  site == "NPB1" ~ "North of Phelps Bridge",
  site == "NPB2" ~ "Phelps Road",
  site == "NWP" ~ "West Pond",
  site == "NDC" ~ "Devereux Creek"
)) |>
# setting factor levels for site
mutate(site_full = fct_reorder(site_full, med_sal, .fun = "median", .na_rm
  ↳ = TRUE),
  site_full = fct_rev(site_full))

```

clean aq_ins_clean

```

# creating new object from clean aquatic inverts
copepods <- aq_ins_clean |>

```

```
# filter to only include copepods
filter(taxon == "copepod") |>
# log + 1 transform mean abundance per liter
mutate(log_ave_lit = log(ave_lit + 1))
```

Problems

1. Visualizing differences across sites in major taxon abundance

a. Plan your figure

- x-axis: site
- y-axis: ostracod abundance
- geometry to represent average abundance/L: `geom_jitter`
- geometry to represent medians: `stat_summary()`

b. Wrangle your data

```
# create new object from aq_ins_clean
ost_abundance <- aq_ins_clean |>
# filter to only ostracods
filter(taxon == "ostracod") |>
# remove missing abundance/L values
filter(!is.na(ave_lit)) |>
# group by site
group_by(site_full) |>
# log + 1 transform mean abundance per liter
mutate(log_ave_lit = log(ave_lit + 1)) |>
# ungroup data frame
ungroup()

slice_sample(ost_abundance, n = 10)
```

A tibble: 10 x 24

	date_site	date	site	taxon	ave_lit	med_ph	med_sal	med_do
	<chr>	<dtm>	<chr>	<chr>	<dbl>	<dbl>	<dbl>	<dbl>
1	2022-08-10_NPB	2022-08-10 00:00:00	NPB	ostr~	2.67	8.28	8.84	9.81
2	2024-01-28_NPB	2024-01-28 00:00:00	NPB	ostr~	3.33	7.43	3.08	6.6

```

3 2022-05-13_NVBR 2022-05-13 00:00:00 NVBR ostr~ 0 9.72 74.2 11.4
4 2023-09-15_NMC 2023-09-15 00:00:00 NMC ostr~ 0.4 NA NA 14.2
5 2022-11-19_NDC 2022-11-19 00:00:00 NDC ostr~ 6 NA NA NA
6 2023-11-11_NEC 2023-11-11 00:00:00 NEC ostr~ 0 NA 38.6 0.25
7 2023-02-28_NPB1 2023-02-28 00:00:00 NPB1 ostr~ 0 NA NA NA
8 2022-02-23_NVBR 2022-02-23 00:00:00 NVBR ostr~ 0 8.86 45.2 10.2
9 2021-08-25_NPB 2021-08-25 00:00:00 NPB ostr~ 581. 7.52 4.57 4.5
10 2020-02-21_NVBR 2020-02-21 00:00:00 NVBR ostr~ 4.8 7.34 4.1 6.1
# i 16 more variables: taxonRank <chr>, scientificName <chr>, kingdom <chr>,
#   Phylum <chr>, Class <chr>, Subclass <chr>, Superorder <chr>, Order <chr>,
#   Infraorder <chr>, Family <chr>, Subfamily <chr>, Tribe <chr>, Genus <chr>,
#   comments <chr>, site_full <fct>, log_ave_lit <dbl>

```

c. Code your figure

```

# base layer
ostracods_plot <- ggplot(data = ost_abundance,
  # x-axis: site
  # y-axis: mean abundance/L
  mapping = aes(x = site_full,
    y = log_ave_lit)) +
# first layer: points to represent observations
geom_jitter(height = 0,
  shape = 21,
  width = 0.1) +
# second layer: points to represent medians
stat_summary(geom = "point",
  fun = "median",
  size = 4,
  color = "darkgreen") +
# wrapping x-axis text
scale_x_discrete(labels = label_wrap(15)) +
# relabelling x- and y-axis, adding title
labs(x = "Site",
  y = "log + 1 transformed average abundance/L",
  title = "Ostracod Abundance") +
# different theme
theme_bw()

# displaying object
ost_abundance

```

```
# A tibble: 51 x 24
```

	date_site <chr>	date <dtm>	site <chr>	taxon <chr>	ave_lit <dbl>	med_ph <dbl>	med_sal <dbl>	med_do <dbl>
1	2020-01-21_NDC	2020-01-21 00:00:00	NDC	ostr~	2.4	6.53	0.3	7.6
2	2020-02-21_NVBR	2020-02-21 00:00:00	NVBR	ostr~	4.8	7.34	4.1	6.1
3	2021-04-15_NPB	2021-04-15 00:00:00	NPB	ostr~	185.	8.09	3.57	8.24
4	2021-08-25_NPB	2021-08-25 00:00:00	NPB	ostr~	581.	7.52	4.57	4.5
5	2021-09-01_NPB2	2021-09-01 00:00:00	NPB2	ostr~	0	NA	NA	NA
6	2021-11-12_NPB	2021-11-12 00:00:00	NPB	ostr~	79.2	NA	NA	NA
7	2021-11-16_NPB	2021-11-16 00:00:00	NPB	ostr~	72.1	7.3	2.38	12.1
8	2021-11-23_NPB2	2021-11-23 00:00:00	NPB2	ostr~	0.267	7.59	1.99	0.65
9	2022-01-21_NEC	2022-01-21 00:00:00	NEC	ostr~	0	NA	NA	NA
10	2022-01-24_NEC	2022-01-24 00:00:00	NEC	ostr~	0.267	NA	NA	NA

```
# i 41 more rows  
# i 16 more variables: taxonRank <chr>, scientificName <chr>, kingdom <chr>,  
#   Phylum <chr>, Class <chr>, Subclass <chr>, Superorder <chr>, Order <chr>,  
#   Infraorder <chr>, Family <chr>, Subfamily <chr>, Tribe <chr>, Genus <chr>,  
#   comments <chr>, site_full <fct>, log_ave_lit <dbl>
```

d. Create a multipanel figure

```
# base layer  
salinity_plot <- ggplot(data = aq_ins_clean,  
  # x-axis: site  
  # y-axis: mean salinity  
  mapping = aes(x = site_full,  
    y = med_sal)) +  
  # first layer: salinity measurements at each survey  
  geom_jitter(height = 0,  
    width = 0.1,  
    shape = 21) +  
  # wrapping x-axis text  
  scale_x_discrete(labels = label_wrap(width = 15)) +  
  # second layer: points to represent means  
  stat_summary(fun = median,  
    color = "red",  
    size = 1) +  
  # relabelling x- and y-axes  
  labs(x = "Site",  
    y = "Median salinity (ppt)",  
    title = "Site Salinity") +
```

```

# different theme
theme_bw()

# base layer
copepods_plot <- ggplot(data = copepods,
                        # x-axis: site
                        # y-axis: mean abundance/L
                        mapping = aes(x = site_full,
                                      y = log_ave_lit)) +
# first layer: points to represent observations
geom_jitter(height = 0,
            shape = 21,
            width = 0.1) +
# second layer: points to represent medians
stat_summary(geom = "point",
            fun = "median",
            size = 4,
            color = "blue") +
# wrapping x-axis text
scale_x_discrete(labels = label_wrap(15)) +
# relabelling x- and y-axis, adding title
labs(x = "Site",
     y = "Average abundance/L",
     title = "Copepod Abundance") +
# different theme
theme_bw()

# base layer
ostracods_plot <- ggplot(data = ostr_abundance,
                        # x-axis: site
                        # y-axis: mean abundance/L
                        mapping = aes(x = site_full,
                                      y = log_ave_lit)) +
# first layer: points to represent observations
geom_jitter(height = 0,
            shape = 21,
            width = 0.1) +
# second layer: points to represent medians
stat_summary(geom = "point",
            fun = "median",
            size = 4,
            color = "darkgreen") +
# wrapping x-axis text

```




```

scale_x_discrete(labels = label_wrap(15)) +
# relabelling x- and y-axis, adding title
labs(x = "Site",
      y = "Average abundance/L",
      title = "Ostracod Abundance") +
# different theme
theme_bw()

(salinity_plot) / (copepods_plot + ostracods_plot) +
plot_annotation(title = "Comparing Salinity, Copepod Abundance, and
  ↪ Ostracod Abundance across sites at NCOS")

```

Comparing Salinity, Copepod Abundance, and Ostracod Abundance across sites at NCOS



e. Interpret your figure

South of Phelps Bridge and Devereux Creek had the highest Average abundance/L measurements across sites for Ostracods. The other sites had pretty consistent lower average abundance/L measurements. North of Venoco Bridge and South of Phelps Bridge had the highest Average abundance/L measurements across sites for Copepods. Main Channel and North of Phelps Bridge had the lowest average abundance/L measurements for Copepods. I could determine

these comparisons by observing how high up on the y axis data points lie, and compare across sites on the x axis.

There does not appear to be a strong correlation between site salinity and ostracod abundance. The site with the highest average salinity (ppt) measurements, South of Vencoco Bridge, does not have a notably different average abundance/L measurements than the other sites. Similarly, the site with the highest average abundance/L measurements, South of Phelps Bridge, did not have comparably high of low salinity (ppt) points along the y axis.

2. Visualizing total abundance as a function of salinity

a. Plan your figure

x-axis: med_sal y-axis: log_ave_lit color: taxon geometry: geom_point() facets: facet_wrap(~taxon, scales = "free")

b. Wrangle your data

```
# create new object from aq_ins_clean
abundance_salinity <- aq_ins_clean |>
  # create log-transformed abundance column
  mutate(log_ave_lit = log(ave_lit + 1)) |>
  # make taxon names title case
  mutate(taxon = str_to_title(taxon)) |>
  # group by taxon
  group_by(taxon) |>
  # calculate max abundance
  mutate(max_abundance = max(log_ave_lit, na.rm = TRUE)) |>
  # un group data frame
  ungroup() |>
  # reorder taxa by maximum abundance
  mutate(taxon = fct_reorder(taxon,
                             max_abundance,
                             .desc = TRUE)) |>
  # keep needed columns
  select(taxon,
         med_sal,
         log_ave_lit)

# display 10 random rows
abundance_salinity |>
```

```
slice_sample(n = 10)
```

```
# A tibble: 10 x 3
```

	taxon	med_sal	log_ave_lit
	<fct>	<dbl>	<dbl>
1	Cladocera	NA	0.125
2	Annelida_polychaete	NA	0
3	Ostracod	2.47	1.10
4	Hemiptera_corixidae_boatman	NA	0.236
5	Annelida_polychaete	NA	0
6	Ostracod	3.08	1.47
7	Cladocera	NA	0
8	Nematode	2.00	0
9	Diptera_chironomid	NA	0
10	Annelida_oligochaete	NA	0

c. Code your figure

d. Interpret your figure

In 3-5 sentences, describe differences or similarities between taxa in relationships between abundance and salinity. Be specific using the visual components of the plot.