

NLA: Golub for $k = 1 : m, n$: $u_k = (\text{sgn}(b_{k,k}) \|b_{k:m,k}\| e_1 + b_{k:m,k})$; $u_k := \hat{u}_k$; $U_k := I - 2u_k u_k^T$;
 $B_{k:m,k:n} := U_k B_{k:m,k:n}$; $U = [I_{k-1,k-1}, 0, 0, U_k]$; **for** $j = 1 : m, n - 1$: $v_k^T := \text{sgn}(b_{k,k+1}) \|b_{k,k+1:n}\| e_1 +$
 $b_{k:m,k}$; $V_k := I - 2v_k v_k^T$; $B_{1:m,k+1:n} = B_{1:m,k+1:n} V_k$; $V = [I_{k,k}, 0, 0, V_k]$ **endfor endfor**; $2 \cdot (2mn^2 - 2n^3/3)$
Householder for $k = [1, n]$: $x = A_{k:m,k}$; $v_k = \text{sgn}(x) \|x\| e_k + x$; $v_k = \frac{v_k}{\|v_k\|}$ **for** $j = [k, n]$ $A_{k:m,j} =$
 $A_{k:m,j} - 2v_k [v_k^* A_{k:m,j}]$ **endfor endfor**. $2mn^2 - \frac{2n^3}{3}$. **MG-S** $V = A$; **for** $i = [1, n]$: $r_{ii} = \|v_i\|$; $q_i = \frac{v_i}{r_{ii}}$; **for**
 $j = [i + 1, n]$ $v_j = v_j - (q_i^T v_j) q_i$; $r_{ij} = q_i^T v_j$ **endfor endfor**. $2mn^2$. **Arnoldi**: $q_1 := \hat{b}$; $q_{k+1} h_{k+1,k} =$
 $A q_k - \sum_{i=1}^k q_i h_{ik}$; $h_{ik} = q_i^T (A q_k)$; $h_{k+1,k} := \|v\| \rightarrow A Q_k := Q_k H_k + q_{k+1} [0 \dots h_{k+1,k}]$. **Givens**
 $3mn^2$ **SVD**: $= \sum_{i=\min(m,n)}^r u_i \sigma_i v_i^T$. **QR Algo**: $A_{k+1} = Q_k^T A_k Q_k \rightarrow A_{k+1} = (Q^{(k)})^T A Q^{(k)}$ $\mathcal{E} A^k =$
 $(Q_1 \dots Q_k) (R_k \dots R_1) := Q^{(k)} R^{(k)}$, via induction **GMRES**: $\min \|A Q_k y - b\| \rightarrow \min \|H_k y - \|b\| e_1\|$ **CG**
Bound: With $c = x - x_0$, $c_k = x_k - x_0$ s.t. $r_k = A(c - c_k)$ we have $r_k^T v = 0 \forall v \in \mathcal{K}_k$ so $v^T A(c - c_k) = 0$,
s.t. $y = c_k = \arg \min \|c - y\|_A$. WTS $e_k = e_0 p_k(A)$ with $p(0) = 1$, and write $e_0 := \sum \gamma_i v_i$ with
 $A v_i = \lambda_i v_i \rightarrow \|e_k\|_A = \min_{p_k, p(0)=1} \max |p(\lambda_i)| \|e_0\|_A$ **CG Convergence**: $\|e_k\| = \min_{p(0)=1} \|p_k(A) e_0\| =$
 $\min_{p_k(A)} \max |p_k(\lambda)| \|e_0\| \rightarrow \leq 2 \left(\frac{(\sqrt{\kappa_2} - 1)}{(\sqrt{\kappa_2} + 1)} \right)^k$; need $\alpha := 2(\lambda_1 + \lambda_2)$ **Cheb**: $T_k(x) = \frac{1}{2}(z^k +$
 $z^{-k})$; $2xT_k = T_{k+1} + T_{k-1}$ **CG Conditions**: To show $r_{k+1}^T r_k = 0$ first show $p_k^T A p_k = p_k^T A r_k$ via β
then show $p_k^T r_k = r_k^T r_k$ via $p_{k-1}^T r_k = 0$. **MP**: $\sigma(G) \in [\sqrt{m} - \sqrt{n}, \sqrt{m} + \sqrt{n}] \rightarrow k_2 = O(1)$ **Sketch**:
with $GA\hat{x} = Gb$, and via $C - F \|G[A, b][v, -1]^T\| \leq (s + \sqrt{n+1}) \|R[v, -1]^T\|$, similar for lower bound
via MP $\rightarrow \|A\hat{x} - b\| \leq (\sqrt{s} + \sqrt{n+1}) / (\sqrt{s} - \sqrt{n+1}) \|Ax - b\|$ **Blend**: solve $\|(A\hat{R}^{-1})y - b\| = 0$ via
CG; $k_2(A\hat{R}^{-1}) = O(1)$ with $GA = \hat{Q}\hat{R}$ **PROOF**: $A = QR$; $GA = GQR = \hat{G}R$. Let $\hat{G} = \hat{Q}\hat{R}$ so
 $GA = \hat{Q}\hat{R}R \rightarrow \hat{R}^{-1} = R^{-1}\hat{R}^{-1} \rightarrow k_2(A\hat{R}^{-1}) = k_2(\hat{R}^{-1}) = O(1)$ by MP. $O(mn)$ to solve via normal
Bounds: $\|ABB^{-1}\| \geq \|AB\| \|B^{-1}\| \rightarrow \|A\| / \|B^{-1}\| \geq \|AB\|$. **Weyls**: $\sigma_i(A+B) = \sigma_i(A) + [-\|B\|, \|B\|]$
Rev Δ Ineq: $\|A - B\| \geq \| \|A\| - \|B\| \|$ **Courant Application**: $\sigma_i([A_1; A_2]) \geq \max(\sigma_i(A_1), \sigma_i(A_2))$
Schur: Take $Av_1 = \lambda_1 v_1$; construct $U_1 = [v_1, V_\perp] \rightarrow AU_1 = U_1[e_1, X]$. Repeat. **Conditioning**
 $\kappa_2(A) = \sigma_1/\sigma_n = \|A\| \|A^{-1}\|$ **Similarity**: $A \rightarrow P^{-1}AP$, same λ .
CO: G-N: $\vec{x}_{k+1} = \vec{x}_k - \frac{\nabla f_k}{J^T J}$, with $J :=$ Jacobian of $r(x)$ **SD**: $\|x_{k+1} - x_*\| \leq (k_2(H) - 1) / (k_2(H) +$
 $1) \|x_k - x_*\|$ with H hessian **bArm**: w/ $\phi(\alpha) = f(x_k + \alpha s_k)$, $\psi(\alpha) = \phi(\alpha) - \phi(0) - \beta \alpha \phi'(0) \leq 0$, show
 $\psi'(0) = (1 - \beta)\phi'(0) \leq 0 \rightarrow \psi(\alpha) \downarrow$ with α . **BFGS**: To show $H_{k+1} \geq 0$ nec. $\gamma^T \delta > 0$. Suff via
 γ, δ LI \rightarrow use $\|\cdot\|_H \rightarrow \gamma^T \delta > 0$. **Quad Penalty Meth** With $y = -c/\sigma$, $\|\nabla_\sigma \Phi\| \leq \epsilon^k, \sigma^k \rightarrow 0, x \rightarrow$
 $x_*, \nabla c(x_*)$ LI, then $y \rightarrow y_*, x \rightarrow KKT$. **PROOF**: If $y_* := J_*^T \nabla f_* \rightarrow \|y_k - y_*\| = \|J_k^T \nabla f_k - I y_*\| \leq$
 $\|J_k^T\| \|\nabla_\sigma \Phi\| \rightarrow 0$. Also, $\nabla f_* - J_*^T y_* = 0$, and $c_{k \rightarrow *} = -\sigma^{k \rightarrow *} y_{k \rightarrow *} = 0$ so $x_* \rightarrow KKT$ **Quad**
Pen. Meth Newt Have $w = (J\Delta x + c)/\sigma$ so $[\nabla^2 f, J^T; J, -\sigma I][\Delta x, w]^T = -[\nabla f, c]$ **Trust Region**
Radius: $\rho_k := (f(x_k) - f(x_k + s_k)) / (f(x_k) - m_k(s_k))$ **TR-Method**: If $\rho \approx 1$ then double radius,
update step $x_{k+1} = x_k + s_k$. If $\rho \geq 0.1$ then same radius, update step. If ρ small shrink radius,
don't update step. **Cauchy**: Is the point on gradient which minimises the quadratic model within
TR. Want $m_k(s_k) \leq m_k(s_{kc})$, where $s_{kc} := -\alpha_{kc} \nabla f(x_k)$, and $\alpha_{kc} := \arg \min m_k(\alpha \nabla f(x_k))$ subject to
 $\|\alpha \nabla f\| \leq \Delta$, i.e. $\alpha_{max} := \Delta / \|\nabla f\|$. **Calculation of Cauchy**: We want to prove cauchy model decrease
i.e. $f(x_k) - m_k(s_k) \geq f(x_k) - m_k(s_{kc}) \geq 0.5 \|\nabla f_k\| \min \left\{ \Delta_k, \frac{\|\nabla f_k\|}{\|\nabla^2 f_k\|} \right\}$. First define $\Psi(\alpha) := m_k(-\alpha \nabla f)$
s.t. $\Psi := f_k - \alpha \|\nabla f_k\|^2 - 0.5 \alpha^2 H_k$, with $H_k := [\nabla f_k]^T [\nabla^2 f_k] [\nabla f_k]$. N.B. that $\alpha_{min} := \frac{\|\nabla f_k\|^2}{H_k}$ if $H_k > 0$,
from $\Psi'(0) < 0$. Now **A: If $H_k \leq 0$** then we have $\Psi(\alpha) \leq f_k - \alpha \|\nabla f_k\|^2 \rightarrow \alpha_{kc} = \alpha_{max}$. So, we have
 $f_k - m_{s_k} \geq f_k - m_{s_{kc}} \geq \|\nabla f_k\| \Delta_k \geq 0.5 \|\nabla f_k\| \min \{\Delta_k\}$. Now **B.i: If $H_k > 0 \rightarrow \alpha_{kc} = \alpha_{min}$** . Here
 $f_k - m_{s_{kc}} = \alpha_{kc} \|\nabla f\|^2 - 0.5 \alpha_{kc}^2 H_k = \frac{\|\nabla f\|^4}{2H_k} \geq \frac{\|\nabla f\|}{2} \min \left\{ \frac{\|\nabla f\|}{\|\nabla^2 f\|} \right\}$ via C-S. Now **B.ii: If $H_k > 0 \rightarrow$**
 $\alpha_{kc} = \alpha_{max}$. Here $\Delta / \|\nabla f\| \leq \|\nabla f\|^2 / H_k \rightarrow \alpha_{kc} H_k \leq \|\nabla f\|^2$. So $f_k - m_{kc} = -\alpha_{kc} \|\nabla f\|^2 + \frac{\alpha_{kc}^2}{2} H_k \geq$
 $\frac{\|\nabla f\|^2}{2} \alpha_{kc} \geq 0.5 \|\nabla f\| \min \{\Delta_k\}$ **TR-Global Convergence**: If $m_k(s_k) \leq m_k(s_{kc})$ then either $\exists k \geq 0$
s.t. $\nabla f_k = 0$ or $\lim \|\nabla f\| \rightarrow 0$. Further, require $f \in C^2$, bounded below and also ∇f L-cont. **PROOF**:
Using def of ρ , $f_k - f_{k+1} \geq \frac{0.1}{2} \|\nabla f_k\| \min \{\dots\}$ from above. Let $\|\nabla^2 f\| := L$, and assuming $\|\nabla f\| \geq \epsilon$
we have $f_k - f_{k+1} \geq 0.05 \frac{\epsilon}{L} \epsilon^2$ assuming TR has a lower bound $c\epsilon/L$. Then sum over all successful jumps
s.t. $f_0 - f_{lower} \geq \sum_{i \in S} f_i - f_{i+1} \geq |S| \frac{0.05 c \epsilon^2}{L}$ **KKT Feasibility**: Need $s^T J \geq 0$, $J_E^T s = 0$, and $s^T \nabla f < 0$.
KKT Conditions: **REMEMBER $c \geq 0, \lambda \geq 0$!** **First Order KKT (Equality)**: If we have x_* local
min, then let $x = x_* + \alpha s$. Then we have $c_i(x(\alpha)) \rightarrow 0 = c_i(x_*) + \alpha s^T J \rightarrow s^T J = 0$. Further, we
have $f(x) = f(x_*) + \alpha s^T \nabla f \rightarrow \alpha s^T \nabla f \geq 0$. Repeat for negative α s.t. $s^T \nabla f = 0$. By Rank-Nullity
(assuming $J_E(x_*)$ full rank), we have $\nabla f_* = J_*^T y + s_*$ for some y_* , which then implies (after s^T from
LHS) that $\|s_*\| = 0$, so $\nabla f_* = J_*^T y_*$. **KKT 2nd Order** If we have $\min f$ with $c(x) \geq 0$, 2nd order con-
ditions are that $s^T \nabla^2 \mathcal{L} s \geq 0$ for all $s \in \mathcal{A}$, with \mathcal{A} defined s.t. **EITHER** $s^T J_i = 0 \forall i$ s.t. $\lambda_i > 0, c_i = 0$,

1 **OR** $s^T J_i \geq 0 \forall i$ s.t. $\lambda_i = 0, c_i = 0$, for J, c, λ evaluated at x_* For EQUALITY constraints instead
 2 need positive definite $\forall s$ s.t. $J^T s = 0$ **Convex Problems** $\hat{x} = KKT \Rightarrow \hat{x} = \arg \min f(x)$. Proof via
 3 $f \geq f(\hat{x}) + \nabla f^T(x - \hat{x})$ so $f \geq f(\hat{x}) + \hat{y}^T A(x - \hat{x}) + \sum_{i \in I} \lambda_i J_i^T(x - \hat{x})$. Choose $Ax = b$, and note that
 4 c_i concave s.t. $\lambda_i J_i^T(\hat{x})(x - \hat{x}) \geq \lambda_i(c_i(x) - c_i(\hat{x})) = \lambda_i c_i(x) \geq 0 \rightarrow f(x) \geq f(\hat{x})$. **Log-Barrier Global**
 5 **Convergence:** With $f \in C^1, \lambda_{ik} = \frac{\mu_k}{c_{ik}}, \|\nabla f_u(x_k)\| \leq \epsilon_k, \mu_k \rightarrow 0, x_k \rightarrow x_*$. Also, $\nabla c(x_*)^T L_i \forall i \in \mathcal{A}$
 6 (active constraints). Then x_* KKT and $\lambda \rightarrow \lambda_*$. **PROOF:** Have $J_A^\dagger(x_*) = (J_A(x_*)J_A(x_*)^T)^{-1}J_A(x_*)$.
 7 Also, $c_A = 0, c_I > 0$. So $\lambda = \mu/c \rightarrow 0$ so $\lambda_I = 0$ as $c_I > 0$. Next $\|\nabla f_k - J_{Ak}\lambda_{Ak}\| \leq \|\nabla f_k - J_k^T \lambda_k\| +$
 8 $\|\lambda_I\| \dots = \|\nabla f_{\mu k}\| \rightarrow 0$. Now $\left\| J_A^\dagger \nabla f_k - \lambda_{kA} \right\| \leq \left\| J_A^\dagger \right\| \|\nabla f_k - J_{Ak}^T \lambda_{Ak}\| \rightarrow 0$. So with triangle ineq
 9 $\left\| \lambda_{kA} - J_{Ak}^\dagger \nabla f_k + J_{Ak}^\dagger \nabla f_k - \lambda_{A*} \right\| \rightarrow 0$, via cont. of ∇f and J^\dagger . Thus $\lambda_{Ak} \rightarrow \lambda_{A*} \geq 0$. Combine s.t.
 10 $\nabla f_k - J_{Ak}^T \lambda_{Ak}$ with $k \rightarrow *$ so get KKT. **Primal-Dual Newton:** Have $\nabla f = J^T \lambda, C(x)\lambda = \mu e$ so
 11 $[\nabla^2 \mathcal{L}, -J^T; \Lambda J, C][dx, d\lambda]^T = -[\nabla f - J^T \lambda, C\lambda - \mu e]^T$