

NS: Classifications: Node: $\lambda_i \in \mathbb{R}, \Pi \lambda_i > 0$ Centre: $\lambda_i = \pm ib$ Focus: $\lambda_i = a \pm ib$ Hyperbolic: $\text{Re}(\lambda) \neq 0 \rightarrow$ hyperbolic. If all $\lambda < 0$ for $\text{Spec}(Df(x_0))$ then A-Stable Invariant Set: $\phi_t(S) \subseteq S$ Lim Pts: ω pt. if $\lim_{t \rightarrow \infty} \phi(x) = p$, i.e. flows tend to p . α pt. if $\lim_{t \rightarrow -\infty} \phi(x) = p$. Attracting Set: A set $A \subseteq S$ if \exists neighbourhood U s.t. $\phi(U) \subseteq U \forall t \geq 0$, and $A = \bigcap_{t > 0} \phi(U)$ Dense Orbits: If $\forall \epsilon > 0, x \in A$ with A an attracting set, $\exists \tilde{x} \in \Gamma$ s.t. $|x - \tilde{x}| < \epsilon$. I.e. a dense orbit goes as close as needed to any point within A Attractor: An attracting set with a dense orbit. Lyapunov Stable: If $\forall \epsilon > 0, \exists \delta > 0$ s.t. $\forall x \in B_\delta, t \geq 0, \phi(t) \in B_\delta$ (i.e. points stay close within region). Asymptotically Stable: If L-Stable and $\exists \delta > 0$ s.t. $\phi(x) \rightarrow x_0 \forall x \in B_\delta$ Lyapunov F'n: $V(x_0) = 0, V(x) > 0 \forall x \neq x_0$. Then if $\dot{V} < 0 \rightarrow$ A-Stable, or if $\dot{V} \leq 0$ L-Stable. Stable Manifold: If spectrum of $Df(x_0)$ has k eigvals with positive real parts, and $n - k$ with negative, then \exists an $n - k$ dim manifold tangent to E^s s.t. for all $t > 0 \phi(W_{loc}^s) \subseteq W_{loc}^s$, and $\forall x \in W_{loc}^s \phi(x) \rightarrow x_0$ as time increases. Repeat for k -dim unstable manifold but for negative time. Then, define e.g. global stable manifold by $W^s(x_0) := \bigcup_{t \leq 0} \phi_t(W_{loc}^s)$. Note that we search backwards in time for stable, and forwards for unstable! Centre Manifold: If x_0 not hyperbolic (0 real part), then E^c is the centre subspace. Then $\exists W^c$ parallel to E^c , of class C^r , and invariant under flow. Want bifurcation at $\mu = 0$, so with change of variables first find eigvecs v_1, v_2 . Then, construct $P := [v_1, v_2]$ s.t. $\tilde{x} = P\tilde{\xi}$. Solve for $\tilde{\xi}$ and then expand with $\eta = h(\tilde{\xi}, \tilde{\mu})$ Transcritical Bifurcation: Always two points, change type at origin. E.g. $\dot{x} = \mu x - x^2$ Saddle-Node: E.g. $\dot{x} = \mu - x^2$ Bifurcation begins to exist at origin. Supercritical: E.g. $\dot{x} = \mu x - x^3$, where stable $\rightarrow 2 \times$ stable and one unstable. Subcritical: E.g. $\dot{x} = -\mu x + x^3$, where unstable $\rightarrow 2 \times$ unstable and one stable. General co-dim 1: If $\dot{x} = f$ then $\dot{x} = \mu f_u + 0.5x^2 f_{xx} + x\mu f_{x\mu} + 0.5\mu^2 f_{\mu\mu}$. Generally this is a saddle-node but if $f_u = 0$ we have $\dot{x} = x\mu f_{x\mu} + 0.5x^2 f_{xx}$, which is a transcritical. However if $x = -x$ then $\dot{x} = x(\mu f_{x\mu} + \dots) + x^3(f_{xxx}/6 + \dots) \rightarrow$ pitchfork. Saddle-node stable under perturbations! HomoClinic Orbits Sum of roots of cubic = - coeff. of x^2 FPDE: Types: 1^{st} : \exists scale s.t. solution found, not so for 2^{nd} . Heat: $\hat{T} = u(\hat{T}_\infty - \hat{T}_{-\infty}) + \hat{T}_{-\infty}$ Oil Spread: Dims: $x = x_f + \varepsilon \xi, t = \tau$ Ground Spread: $(1-s)\phi h_t + Q_x = 0; Q \sim -hh_x, 0 < x_s < x_f$. Have $h(x_f) = 0, h_t(x_s) = 0$, and $hh_x|_{x=0, x_f} = 0$ (i.e. no flux at centre and front), and h, hh_x cont. at joint. Expansions: Let $\xi = z + \varepsilon \eta$ for perturbations Scale: Try $x = x_f + \varepsilon \xi$ for groundwater Stefan: $S_0 = C(T_1 - T_m)/L$, condition = $\rho L \dot{s} = kT_x|_{s_-}^{s_+}$ 1ph Stefan: Bar = $T_h |liq|_s sol|INS$. Use $T = T_m + (T_1 - T_m)u$ s.t. $S_0 u_t = u_{xx}, u = 1 @ x = 0, \{\dot{s} = -u_x, u = 0\} @ x = s, s(0) = 0$. Sim. sol is $s = \beta \sqrt{t}, f = f(x/\sqrt{t})$ 2ph Stefan: Use $T = T_m + (T_1 - T_m)u$ s.t. $S_0 u_t = u_{xx} @ 0 < x < s, (S_0/\kappa)u_t = u_{xx} @ s < x < 1, u = 1 @ x = 0, u_x = 0 @ x = 1, \{\dot{s} = Ku_x|_{s_+} - u_x|_{s_-}, u = 0\} @ x = s, \{s = 0, u = -\theta\} @ x = 0$. Here $\theta := (T_m - T_0)/(T_1 - T_m), \kappa := c_1 k_1/(c_2 k_2), K := k_2/k_1$ Sim. sol is $s = \beta \sqrt{t}, f = f(x/\sqrt{t})$ 2-Dim: $U_n = \hat{n} \cdot u = K(u_2)_n - (u_1)_n$. If $x = f(y, t)$ then $\hat{n} := \nabla(x - f) = [1, -f_y]^T / \sqrt{1 + f_y^2}$ Welding: Have $0 < s_2 < s_1$. Have cold $x = a$, no flux $x = 0$. $\theta = 1$ in liquid. In mush $\rho L \theta_t = J^2/\sigma$, CoE $\rightarrow \theta \rho L \dot{s} + kT_x|_{s_-}^{s_+} = 0$. Have θ cont. (=0) at s_1 . I.e. we have $S_0 u_t = u_{xx} + q, u_x = 0 @ x = 0, u = -1 @ x = 1, \theta = 0 @ x = s_1$. Also $\theta_t = q$ in mush. FMM: Integral Constraint If $J[y] = \int F dx$ with $\int G dx = C$ then $\tilde{J}[y] = \int F - \lambda G dx$ Hamiltonian: $H := y' F_{y'} - F \rightarrow H' = -F_x$. If $F = F(y, \dot{y})$ then $H = C$ Hamilton's Eqs: $p := F_{y'}, q = y$ and so $p' = -H_q, q' = H_p$ Free Boundary: $J[y, b] = \int_a^b F(x, y, y') dx$ where b free. Expand with $y + \varepsilon \eta, b + \varepsilon \beta \rightarrow J = J_0 + \varepsilon \left\{ \int_a^b \eta F_y + \eta' F_{y'} dx + \beta F(b, y(b), y'(b)) \right\}$ If $y(b) = d \rightarrow d = y(b + \varepsilon \beta) + \varepsilon \eta(b + \varepsilon \beta) = y(b) + \varepsilon(\beta y'(b) + \eta(b))$ so $\eta(b) = -\beta y'(b)$. IVP on integral so $\beta [F - y' F_{y'}]_{x=b} + \int (\dots) = 0$ so $F = y' F_{y'}$ at free boundary. Control: Have $\int \xi h_x + \eta h_u dt = 0, \dot{\xi} = \xi f_x + \eta f_u$. Sub for η , IVP s.t. $\frac{d}{dt} \frac{h_u}{f_u} = h_x - f_x \frac{h_u}{f_u}$ and $\dot{x} = f$ Hamiltonian (Control): $H := f \frac{h_u}{f_u} - h$ s.t. $\dot{H} = \frac{h_u}{f_u} f_t - h_t \rightarrow$ autonomous if $h_t = f_t = 0$. Fredholm Alt Integ Eqs. For $y = f + \int K(x, t)y(t)dt$ we have ONE (N) has a unique sol $y = 0$ if $f = 0$, and adjoint has unique sol, or TWO (H) as sols $y_1 \dots y_r$ iff \forall solutions to H^*, z_i , we have $\langle f, z_i \rangle = 0$. EX: Solve $y = f + \lambda \int \sin(x + t)y(t)dt$. Unique sol iff (H) has trivial sol $\rightarrow X_1 = \int y \cos(t) = \int \cos(t)y_H(t) \rightarrow$ solve $[1, -\lambda\pi; -\lambda\pi, 1][X_1, X_2]^T = [0, 0]^T \rightarrow$ unique sol if $\lambda \neq \pm 1/\pi$. In this case $X_1 = \int \cos(x)y_N(x) = \lambda\pi X_2 + \int f(x)\cos(x)$, and similar for X_2 . Invert matrix and solve. If non-unique sol, then find sols to (H) first. If $\lambda = 1/\pi$ then $X_1 = X_2 = X$ so $Ly = y - \pi^{-1}(\sin(x) + \cos(x)) \int \cos(x)y(x)dx$ with sols $y = c_1(\sin(x) + \cos(x))$ by inspection. Problem self adjoint so $Ly = 0 = L^*w$ so need $\int f(x)w(x) = 0$ i.e. $\int f(x)(\sin(x) + \cos(x)) = 0$, repeat for $\lambda = -1/\pi$. Then $y = y_p(x) + \sum_i y_{h,i}(x)$ Fred Diff Eq. For nonunique sol to exist, need $\langle Ly, w \rangle = \langle f, w \rangle \forall w$ s.t. $L^*w = 0$