

Impact of Environmental Factors

on Acute Myocardial Infarction

Gabriela Zemenčíková

June, 2024



UNIVERSITAT DE
BARCELONA

Dataset & Data Preparation

- The environmental dataset and census data taken from the **Statistical Institute of Catalonia**.

Dataset & Data Preparation

- The environmental dataset and census data taken from the **Statistical Institute of Catalonia**.
- AMI dataset taken from ten main hospitals across Catalonia.

Dataset & Data Preparation

- The environmental dataset and census data taken from the **Statistical Institute of Catalonia**.
- AMI dataset taken from ten main hospitals across Catalonia.
- 22,812 hospital admissions at the scale of the 948 municipalities in Catalonia stratified by province, sex and age

Dataset & Data Preparation

- The environmental dataset and census data taken from the **Statistical Institute of Catalonia**.
- AMI dataset taken from ten main hospitals across Catalonia.
- 22,812 hospital admissions at the scale of the 948 municipalities in Catalonia stratified by province, sex and age
- **Response variable:** Age Standardised Incidence Rate (ASIR)

Dataset & Data Preparation

- The environmental dataset and census data taken from the **Statistical Institute of Catalonia**.
- AMI dataset taken from ten main hospitals across Catalonia.
- 22,812 hospital admissions at the scale of the 948 municipalities in Catalonia stratified by province, sex and age
- **Response variable:** Age Standardised Incidence Rate (ASIR)
- A total of **5** predictors - Humidity, Temperature, Ozone levels, Particulate Matter, Public holidays

Choice of Age-Standardized Incidence Rates (ASIR)

- Ensures comparability of events across populations.

Choice of Age-Standardized Incidence Rates (ASIR)

- Ensures comparability of events across populations.
- Ensures comparability across different age distributions.


Choice of Age-Standardized Incidence Rates (ASIR)

- Ensures comparability of events across populations.
- Ensures comparability across different age distributions.
- Adjusts for age as a confounding factor, providing a more accurate representation of AMI incidence.


Choice of Age-Standardized Incidence Rates (ASIR)

- Ensures comparability of events across populations.
- Ensures comparability across different age distributions.
- Adjusts for age as a confounding factor, providing a more accurate representation of AMI incidence.
- Allows comparisons between regions or over time periods.


Target Variable - ASIR




Target Variable - ASIR



Target Variable - ASIR



Target Variable - ASIR



SARIMAX

- **S**: Seasonal component that captures periodic patterns at fixed intervals.
- **AR**: Autoregressive part that models the dependency between an observation and a number of lagged observations.
- **I**: Integrated part representing the differencing of raw observations to make the time series stationary.
- **MA**: Moving Average component that models the dependency between an observation and a residual error from a moving average model applied to lagged observations.
- **X**: Exogenous variables that are external factors influencing the time series.

SARIMAX

- **S**: Seasonal component that captures periodic patterns at fixed intervals.
- **AR**: Autoregressive part that models the dependency between an observation and a number of lagged observations.
- **I**: Integrated part representing the differencing of raw observations to make the time series stationary.
- **MA**: Moving Average component that models the dependency between an observation and a residual error from a moving average model applied to lagged observations.
- **X**: Exogenous variables that are external factors influencing the time series.

By incorporating exogenous variables, SARIMAX can model the influence of external factors on ASIR.

LSTM

- a type of recurrent neural network (RNN) architecture specifically designed to model sequence data while addressing the vanishing gradient problem.

LSTM

- a type of recurrent neural network (RNN) architecture specifically designed to model sequence data while addressing the vanishing gradient problem.
- The LSTM cell has a memory cell and three gates: input gate, forget gate, and output gate


LSTM

- a type of recurrent neural network (RNN) architecture specifically designed to model sequence data while addressing the vanishing gradient problem.
- The LSTM cell has a memory cell and three gates: input gate, forget gate, and output gate
- LSTM networks are capable of learning long-term dependencies in sequential data, making them suitable for time series forecasting tasks.

LSTM

- a type of recurrent neural network (RNN) architecture specifically designed to model sequence data while addressing the vanishing gradient problem.
- The LSTM cell has a memory cell and three gates: input gate, forget gate, and output gate
- LSTM networks are capable of learning long-term dependencies in sequential data, making them suitable for time series forecasting tasks.
- can capture complex patterns and relationships to forecast future ASIR.

Results - SARIMAX




Results - SARIMAX

- Maximum temperature (`max_temp`): A decrease of 1°C in maximum temperature is associated with a decrease of approximately 0.345 units in the Age-Standardized Incidence Rate (ASIR) of AMI.


Results - SARIMAX

- Maximum temperature (`max_temp`): A decrease of 1°C in maximum temperature is associated with a decrease of approximately 0.345 units in the Age-Standardized Incidence Rate (ASIR) of AMI.
- Mean PM10 concentration (`mean_PM10`): An increase of $1 \mu\text{g}/\text{m}^3$ in mean PM10 concentration is associated with an increase of approximately 0.146 units in ASIR of AMI.

Diagnostics - SARIMAX



Results - LSTM



Comparison

Table 1: Results

	SARIMAX	LSTM
MAE	0.12	0.15
MSE	0.02	0.03
RMSE	0.15	0.19
MAPE	26.60	14.53

Comparison

Table 1: Results

	SARIMAX	LSTM
MAE	0.12	0.15
MSE	0.02	0.03
RMSE	0.15	0.19
MAPE	26.60	14.53

- SARIMAX outperforms LSTM in terms of both MAE and MSE, making it a more suitable choice for this forecasting task.

Comparison

Table 1: Results

	SARIMAX	LSTM
MAE	0.12	0.15
MSE	0.02	0.03
RMSE	0.15	0.19
MAPE	26.60	14.53

- SARIMAX outperforms LSTM in terms of both MAE and MSE, making it a more suitable choice for this forecasting task.
- However, the LSTM model exhibits substantially lower AIC and BIC values compared to SARIMAX, indicating a potentially better fit to the data and superior long-term forecasting capabilities

Thank you for your attention