

# Impact of Environmental Factors

## on Acute Myocardial Infarction

Gabriela Zemenčíková

June, 2024



UNIVERSITAT DE  
BARCELONA

# Dataset & Data Preparation

---

- The environmental dataset and census data taken from the **Statistical Institute of Catalonia**.

# Dataset & Data Preparation

---

- The environmental dataset and census data taken from the **Statistical Institute of Catalonia**.
- AMI dataset taken from ten main hospitals across Catalonia.

# Dataset & Data Preparation

---

- The environmental dataset and census data taken from the **Statistical Institute of Catalonia**.
- AMI dataset taken from ten main hospitals across Catalonia.
- 22,812 hospital admissions at the scale of the 948 municipalities in Catalonia stratified by province, sex and age

# Dataset & Data Preparation

---


- The environmental dataset and census data taken from the **Statistical Institute of Catalonia**.
- AMI dataset taken from ten main hospitals across Catalonia.
- 22,812 hospital admissions at the scale of the 948 municipalities in Catalonia stratified by province, sex and age
- **Response variable:** Age Standardised Incidence Rate (ASIR)

# Dataset & Data Preparation

---


- The environmental dataset and census data taken from the **Statistical Institute of Catalonia**.
- AMI dataset taken from ten main hospitals across Catalonia.
- 22,812 hospital admissions at the scale of the 948 municipalities in Catalonia stratified by province, sex and age
- **Response variable:** Age Standardised Incidence Rate (ASIR)
- A total of **5** predictors - Humidity, Temperature, Ozone levels, Particulate Matter, Public holidays

# Target Variable - ASIR



# Target Variable - ASIR


---






# Target Variable - ASIR

---



# Target Variable - ASIR

---



# SARIMAX

---

- **S**: Seasonal component that captures periodic patterns at fixed intervals.
- **AR**: Autoregressive part that models the dependency between an observation and a number of lagged observations.
- **I**: Integrated part representing the differencing of raw observations to make the time series stationary.
- **MA**: Moving Average component that models the dependency between an observation and a residual error from a moving average model applied to lagged observations.
- **X**: Exogenous variables that are external factors influencing the time series.

# SARIMAX

---

- **S**: Seasonal component that captures periodic patterns at fixed intervals.
- **AR**: Autoregressive part that models the dependency between an observation and a number of lagged observations.
- **I**: Integrated part representing the differencing of raw observations to make the time series stationary.
- **MA**: Moving Average component that models the dependency between an observation and a residual error from a moving average model applied to lagged observations.
- **X**: Exogenous variables that are external factors influencing the time series.

By incorporating exogenous variables, SARIMAX can model the influence of external factors on ASIR.

# LSTM

---

- a type of recurrent neural network (RNN) architecture specifically designed to model sequence data while addressing the vanishing gradient problem.

# LSTM

---

- a type of recurrent neural network (RNN) architecture specifically designed to model sequence data while addressing the vanishing gradient problem.
- The LSTM cell has a memory cell and three gates: input gate, forget gate, and output gate

# LSTM

---

- a type of recurrent neural network (RNN) architecture specifically designed to model sequence data while addressing the vanishing gradient problem.
- The LSTM cell has a memory cell and three gates: input gate, forget gate, and output gate
- LSTM networks are capable of learning long-term dependencies in sequential data, making them suitable for time series forecasting tasks.


# LSTM

---

- a type of recurrent neural network (RNN) architecture specifically designed to model sequence data while addressing the vanishing gradient problem.
- The LSTM cell has a memory cell and three gates: input gate, forget gate, and output gate
- LSTM networks are capable of learning long-term dependencies in sequential data, making them suitable for time series forecasting tasks.
- can capture complex patterns and relationships to forecast future ASIR.



# Results - SARIMAX



# Results - SARIMAX

---


- Maximum temperature (`max_temp`): A decrease of 1°C in maximum temperature is associated with a decrease of approximately 0.345 units in the Age-Standardized Incidence Rate (ASIR) of AMI.

# Results - SARIMAX


---

- Maximum temperature (`max_temp`): A decrease of  $1^{\circ}\text{C}$  in maximum temperature is associated with a decrease of approximately 0.345 units in the Age-Standardized Incidence Rate (ASIR) of AMI.
- Mean PM10 concentration (`mean_PM10`): An increase of  $1 \mu\text{g}/\text{m}^3$  in mean PM10 concentration is associated with an increase of approximately 0.146 units in ASIR of AMI.

# Diagnostics - SARIMAX



# Results - LSTM



# Comparison

---

Table 1: Results

	SARIMAX	LSTM
MAE	<b>0.12</b>	0.15
MSE	<b>0.02</b>	0.03
RMSE	<b>0.15</b>	0.19
MAPE	26.60	14.53

# Comparison

---

Table 1: Results

	SARIMAX	LSTM
MAE	<b>0.12</b>	0.15
MSE	<b>0.02</b>	0.03
RMSE	<b>0.15</b>	0.19
MAPE	26.60	14.53

- SARIMAX outperforms LSTM in terms of both MAE and MSE, making it a more suitable choice for this forecasting task.

# Comparison

---

Table 1: Results

	SARIMAX	LSTM
MAE	<b>0.12</b>	0.15
MSE	<b>0.02</b>	0.03
RMSE	<b>0.15</b>	0.19
MAPE	26.60	14.53

- SARIMAX outperforms LSTM in terms of both MAE and MSE, making it a more suitable choice for this forecasting task.
- However, the LSTM model exhibits substantially lower AIC and BIC values compared to SARIMAX, indicating a potentially better fit to the data and superior long-term forecasting capabilities



Thank you for your attention