

**Exercise 3.2**

Let  $R$  be a UFD. Show that any prime ideal of height one is principal.

*Solution:* Let  $\mathfrak{p}$  be a prime ideal in  $R$ . The zero ideal  $(0)$  is properly contained in  $\mathfrak{p}$ , so there exists a nonzero element  $x \in \mathfrak{p}$ . Since  $R$  is a UFD,  $x$  can be written as

$$x = ux_1 \cdots x_n$$

where  $u \in R$  is a unit and  $x_1, \dots, x_n$  are irreducible elements in  $R$ . We know that  $\mathfrak{p}$  is a prime ideal, so at least one  $x_i \in \mathfrak{p}$  for  $1 \leq i \leq n$ . Without loss of generality, we can assume  $x_1 \in \mathfrak{p}$ . The principal ideal  $(x_1) \subset \mathfrak{p}$  is also prime because  $x_1$  is irreducible, and since  $\mathfrak{p}$  has height 1,  $(x_1) = \mathfrak{p}$ , namely the prime ideal  $\mathfrak{p}$  is principal.

**Exercise 3.19**

Consider the curve  $C$  in  $\mathbb{A}^2$  whose equation is  $y^2 = x^3$ . Show that  $C$  can be parametrized by the map

$$\begin{aligned} \phi : \mathbb{A}^1 &\rightarrow \mathbb{A}^2, \\ t &\mapsto (t^2, t^3). \end{aligned}$$

Describe the map  $\phi^* : A(C) \rightarrow A(\mathbb{A}^1)$ . Show that  $\phi$  is bijective but not an isomorphism. Show that the function field of  $C$  equals  $k(t)$ .

*Solution:* For all  $t \in \mathbb{A}^1$ , it is easy to see that the point  $\phi(t) = (t^2, t^3)$  is a point on  $C$ , so  $\text{Im } \phi \in C$ . Moreover,  $\phi$  is injective because

$$\begin{cases} t_1^2 = t_2^2 \\ t_1^3 = t_2^3 \end{cases}$$

implies that  $t_1 = t_2$ . Conversely, suppose  $(a, b)$  is a point on  $C$ . If  $a = b = 0$ , choose  $t = 0$  and  $\phi(0) = (a, b)$ . If  $a \neq 0$  and  $b \neq 0$ , the equation  $x^2 = a$  has two different solutions in  $\mathbb{C}$ . Suppose  $t$  is such a solution. Note that

$$t^6 = (t^2)^3 = a^3 = b^3.$$

This implies that either  $t^3 = b$  or  $t^3 = -b$ . Choose the solution  $t$  satisfying  $t^3 = b$ . Thus, we find a preimage  $t \in \mathbb{A}^1$ . This proves that  $C$  can be parametrized by the map  $\phi$  which is a bijective map.

The map  $\phi^* : A(C) \rightarrow A(\mathbb{A}^1)$  is given by

$$\phi^* : k[x, y]/(y^2 - x^3) \rightarrow k[t], \quad (1)$$

$$x \mapsto t^2, \quad (2)$$

$$y \mapsto t^3. \quad (3)$$

This map  $\phi^*$  is not surjective as  $t \in k[t]$  is not in the image. Hence,  $\phi$  is not an isomorphism of affine varieties.

The image of  $\phi^*(A(C))$  is isomorphic to the subring of  $k[t]$  with only degree  $\geq 2$  part. It is an integral domain. Let  $F$  be the field of fractions for this subring, which is isomorphic to the function field of  $C$ . We claim that it is isomorphic to  $k(t)$ . Indeed, note that

$$t = t^3 \cdot (t^2)^{-1}, \quad t^{-1} = (t^3)^{-1} \cdot t^2.$$

This implies  $F$  is subfield of  $k(t)$  containing  $t$  and  $t^{-1}$ , so  $F = k(t)$ .

---