Trixe (T) Dynamics: A Mathematical Framework for Information Transformation from Probabilistic Chaos to Absolute Determinism

Muhammad Rifat Qolby Almusawa

Independent Researcher

Abstract

The Trixe model introduces a tripartite architecture connecting the probabilistic domain ($< Xe$), the stabilization mechanism (Xe via the Orbital Variable O_v), and the information singularity ($> Xe$) through the transition pathway $0Xe$. This paper formulates the evolution equations, stabilizer operators, non-linear coupling mechanisms, and singularity operators, rendering the model mathematically verifiable [9, 8]. Quantitatively, the model identifies a critical bifurcation point at $\alpha = \beta$ as the boundary of asymptotic stability, which dictates the successful transformation of information from stochastic fluctuations into an absolute deterministic output.

Index Terms: Absolute Determinism; Bifurcation Analysis; Orbital Variable; Probabilistic Chaos; Trixe Model.

---

1 Introduction and Scientific Positioning

The transition from probabilistic to deterministic systems is a fundamental problem that has long been explored through various lenses, ranging from dissipative structures far from equilibrium [6] to *Synergetics*, which formulates order parameters in self-organization [2]. In the domain of stochastic control and signal processing, Bayesian filters are frequently employed to incrementally reduce uncertainty [3].

However, traditional approaches generally view deterministic transitions as a byproduct of slow thermodynamic stabilization. The Trixe (T) Model distinguishes itself by proposing an operational structural hypothesis: it positions the *information singularity* not as a point of mathematical failure, but as a mechanistic operator that actively processes and collapses entropy.

It must be emphasized that within the context of this framework, the term "Absolute Determinism" is not proposed as a philosophical or epistemological claim. Rather, it is defined strictly by operational boundaries as the limit condition where the variance of the system's probability distribution decays to zero ($\sigma^2 \to 0$) as $t \to \infty$, resulting in a single phase trajectory that is entirely predictable mathematically.

This paper does not aim to challenge classical statistical mechanics, but rather to expand the stochastic control framework. By introducing a tripartite architecture ($<$

Xe, $0Xe$, $> Xe$), the Trixe Model provides a formulation demonstrating how low-level probabilistic fluctuations can be forced to converge asymptotically into a deterministic output through the intervention of orbital variables and non-linear operators.

2 Mathematical Structure and Formulation

2.1 Notation and Basic Assumptions

- **Spatial domain:** $\Omega \subset \mathbb{R}^n$.
- **Time:** $t \in [0, \infty)$.
- **Function assumption:** All functions are assumed to be sufficiently smooth (class C^2).
- **Parameters:** $\alpha, \beta, \lambda_k, \kappa_{ji}$ operate with the sign conditions necessary for dynamical system stability [1].

2.2 $< Xe$ Probabilistic Chaos Model

The $< Xe$ domain is defined through the probability density $P(x,t)$. The stochastic representation is modeled via the Langevin process [4, 5]:

$$dx_t = a(x_t, t)dt + b(x_t, t)dW_t \tag{1}$$

where W_t denotes the Wiener process. Its density evolution counterpart is the Fokker-Planck equation [7]:

$$\frac{\partial P(x,t)}{\partial t} = -\nabla \cdot (a(x,t)P(x,t)) + \nabla^2(D(x,t)P(x,t)) \tag{2}$$

Where $D(x,t) = \frac{1}{2}b(x,t)b(x,t)^T$ represents the local diffusion matrix.

2.3 Stabilizer and Orbital Variable O_v

The system introduces a stabilizer vector $S(t) = (S_1(t), ..., S_m(t))$ that satisfies the Ordinary Differential Equation (ODE):

$$\frac{dS_j}{dt} = -\alpha_j S_j + \beta_j G_j[P(\cdot, t)] + \sum_i \kappa_{ji} H_{ji}(S_i, P) \tag{3}$$

Where $\alpha_j > 0$ is the asymptotic damping coefficient. The orbital variable is defined as the limit:

$$O_V := \lim_{t \to \infty} S(t) \tag{4}$$

2.4 Inter-Component Coupling and Operational Dimensions

Coupling between components necessitates dimensional care. Suppose there exists a deterministic field parameter u_S and a weighting function $w(x)$. To maintain dimensional consistency within the coupling equation, $w(x)$ is defined as a dimensionless scalar (representing a normalized probability density), while u_S and its transfer function possess spatial dimensions congruent with the domain $x \in \Omega$.

2.5 Explicit Formulation: 1D Toy-Model

To render this model *verifiable*, we consider a 1D "Toy-Model" case with a single spatial variable x and one stabilizer $S(t)$. The functional $G[P]$ is explicitly defined as the first moment (expectation value) of the distribution:

$$G[P] = \int_\Omega xP(x,t)dx = \mu(t) \tag{5}$$

The non-linear coupling term $H(S,P)$ is formulated to respond to spatial variance:

$$H(S,P) = S(t)\left(\int_\Omega x^2 P(x,t)dx - \mu(t)^2\right) = S(t)\sigma^2(t) \tag{6}$$

Substitution into Equation (3) yields a closed system. This step involves a high-degree moment simplification assumption (the *moment closure problem*), where the third moment (*skewness*) and fourth moment (*kurtosis*) are assumed not to contribute significantly to the coupling dynamics when the system is near equilibrium:

$$\frac{dS}{dt} = -\alpha S + \beta\mu(t) + \kappa S\sigma^2(t) \tag{7}$$

This explicit formulation allows for the analytic computation of fixed points and the evaluation of the 2×2 Jacobian matrix to test stability boundaries.

2.6 $0Xe$ - Imperfection Dynamics

The transition pathway is modeled as the variable $R(t)$:

$$\frac{dR_k}{dt} = -\lambda_k R_k + \Gamma_k[P,S] + \sum_l \mu_{kl}\Phi_{kl}(R_l) \tag{8}$$

The condition $\lambda_k > 0$ is enforced to prevent divergence.

2.7 $> Xe$ Information Singularity and Operator h

The singularity is viewed as a transformation operator h that converts the distribution P into a deterministic output.

- **Kernel collapse (regularization):**

$$h[P](y) = \lim_{\sigma\to 0}\int_\Omega K_\sigma(y,z)P(z)dz \approx \delta(y-y^*) \tag{9}$$

3 Numerical Verification and Stability

Numerical Implementation Methodology

The integration of system variables requires a strict mixed approach due to the presence of stochastic variables (the Wiener process).

- The macroscopic probability evolution (Fokker-Planck Equation) is integrated using the Finite Difference Method via the *Crank-Nicolson* scheme to guarantee unconditional numerical stability and mass conservation $\int P = 1$.
- The stochastic probability sample realizations (SDE) are executed using the **Euler-Maruyama** scheme to ensure the validity of strong convergence within Ito calculus.
- The coupled deterministic ODE system (evolution of variable $S(t)$) is integrated using the **Fourth-Order Runge-Kutta (RK4)** scheme.

Numerical Simulation Formulation (Euler-Maruyama and RK4)

Based on the test scenario parameters, the following discrete differential equation systems are executed simultaneously with a time resolution of Δt:

1. Probabilistic Sample Evolution (Euler-Maruyama):
The Langevin equation is simulated for an ensemble of N probability agents (representing the phase space distribution X) using the Euler-Maruyama scheme:

$$X_{n+1}^{(i)} = X_n^{(i)} - k_{pull}(X_n^{(i)} - S_n)\Delta t + \sigma\sqrt{\Delta t} Z_n^{(i)} \tag{10}$$

where $Z_n^{(i)} \sim \mathcal{N}(0,1)$ is a standard Gaussian random variable, k_{pull} is the singularity pull strength toward determinism, and σ is the initial chaos intensity (noise). At each step n, the first moment (expectation) of the probability is extracted empirically:

$$\mu_n = \frac{1}{N}\sum_{i=1}^{N} X_n^{(i)} \tag{11}$$

2. Stabilizer Evolution (4th-Order Runge-Kutta):
The deterministic variable $S(t)$ is integrated based on the response function $f(S,\mu) = -\alpha S + \beta\mu$ using the RK4 scheme:

$$S_{n+1} = S_n + \frac{\Delta t}{6}(k_1 + 2k_2 + 2k_3 + k_4) \tag{12}$$

with the following gradient evaluation stages:

$$\begin{aligned}
k_1 &= f(S_n, \mu_n) \\
k_2 &= f\left(S_n + \frac{\Delta t}{2}k_1, \mu_n\right) \\
k_3 &= f\left(S_n + \frac{\Delta t}{2}k_2, \mu_n\right) \\
k_4 &= f(S_n + \Delta t k_3, \mu_n)
\end{aligned}$$

Numerical Simulation Results

Numerical testing was conducted under two primary parameter scenarios.



Figure 1: Stability Scenario ($\alpha > \beta$). Macrodynamics (left) demonstrate smooth convergence toward the Orbital Variable $O_v = 0$. Microdynamics (right) demonstrate the projective function of the $> Xe$ singularity.



Figure 2: Trixe Bifurcation Diagram. Confirms the existence of a runaway point (critical threshold) at $\alpha = \beta = 1.5$. Beyond this boundary, the stabilizer deviation explodes exponentially.



Figure 3: Failure Scenario and Critical Boundary ($\beta > \alpha$). Macrodynamics (left) and Microdynamics (right) show that when the response force exceeds the damping power, the system undergoes a runaway feedback divergence toward infinity.

4 Conceptual Analysis and Discussion

The singularity within the Trixe framework functions as a functional engine that processes probabilistic information into a deterministic output. Numerical verification proves that the transition toward a deterministic state ($> Xe$) is not a natural equilibrium of the system, but rather a conditional state heavily dependent on damping efficiency.

There exists a critical threshold (bifurcation point $\alpha = \beta$) that acts as a **runaway point (asymptotic stability boundary)**. If the probabilistic entropy exerts a feedback force that exceeds the stabilizer's damping power metric ($\beta > \alpha$), the Trixe functionality will experience infinite divergence. The stability modulus will fracture, aborting convergence and reverting the system to pure chaos. This demonstrates that certainty in dynamical systems is a functional product of a strictly calibrated control architecture.

5 Conclusion

The Trixe Model offers a structured mathematical framework for controlling the transformation of information from probabilistic chaos to deterministic convergence. The integration of the Orbital Variable (O_v) and the Information Singularity mechanism provides a measurable operational transformation pathway. Stability analysis confirms the existence of a critical boundary for component interactions at $\alpha = \beta$.

Model Limitations: As a note of academic objectivity, the current model formulation is largely based on the assumption of Gaussian white noise perturbations (standard Wiener process). Analysis of Trixe's robustness against colored noise or jump processes (Lévy flights) within the $< Xe$ domain remains an open subject. For further development, this functional framework has pragmatic application potential in low-level computational noise control, complex network synchronization, and non-linear control optimization.

Data Availability Statement

The Python source code implementing the coupled numerical integration (Euler-Maruyama and RK4 schemes), along with datasets and high-resolution figures supporting the findings in this manuscript, are publicly available in an open repository. Data can be accessed via the following link: `https://github.com/zerghirsaw/Trixe-Theory/tree/main/test`.

References

[1] Guckenheimer, J., & Holmes, P. (1983). *Nonlinear Oscillations, Dynamical Systems, and Bifurcations of Vector Fields*. Springer-Verlag.

[2] Haken, H. (1983). *Synergetics: An Introduction and Advanced Topics*. Springer-Verlag.

[3] Jazwinski, A. H. (1970). *Stochastic Processes and Filtering Theory*. Academic Press.

[4] Langevin, P. (1908). Sur la théorie du mouvement brownien. *Comptes Rendus de l'Académie des Sciences*, 146(1), 530–533.

[5] Lemons, D. S., & Gythiel, A. (1997). Paul Langevin's 1908 paper on the theory of Brownian motion. *American Journal of Physics*, 65(11), 1079–1081.

[6] Prigogine, I. (1977). Time, Structure and Fluctuations. *Nobel Lecture*, 261–275.

[7] Risken, H. (1989). *The Fokker-Planck Equation: Methods of Solution and Applications* (2nd ed.). Springer.

[8] Shannon, C. E. (1948). A Mathematical Theory of Communication. *Bell System Technical Journal*, 27(3), 379–423.

[9] Strogatz, S. H. (1994). *Nonlinear Dynamics and Chaos: With Applications to Physics, Biology, Chemistry, and Engineering*. Westview Press.