

# Emergenet: Fast Scalable Pandemic Risk Assessment of Influenza A Strains Circulating In Non-human Hosts

Kevin Wu<sup>1</sup>, Jin Li<sup>1</sup>, Timmy Li<sup>1</sup>, Aaron Esser-Kahn<sup>2,3</sup>, and Ishanu Chattopadhyay<sup>1,4,5★</sup>

<sup>1</sup>Department of Medicine, University of Chicago, IL, USA

<sup>2</sup>Pritzker School of Molecular Engineering, University of Chicago, Chicago, IL, USA

<sup>3</sup>Committee on Immunology, University of Chicago, Chicago, IL, USA

<sup>4</sup>Committee on Genetics, Genomics & Systems Biology, University of Chicago, IL, USA

<sup>5</sup>Committee on Quantitative Methods in Social, Behavioral, and Health Sciences, University of Chicago, IL, USA

\*To whom correspondence should be addressed: e-mail: ishanu@uchicago.edu.




**Abstract:** Animal Influenza A strains emerging into humans can introduce large antigenic shifts in the circulating population, and are suspected to have triggered devastating pandemics in the past<sup>1–4</sup>. Yet, our current ability to scalably evaluate the pandemic potential of individual strains that do not yet circulate in humans, remains limited. In this study we introduce the Emergenet, to computationally learn how viable genotypic variations are shaped by evolutionary constraints using only genomic sequences of key viral proteins. Analyzing Hemagglutinin (HA) and Neuraminidase (NA) amino acid sequences from nearly 100,000 unique Influenza A strains recorded in public databases, we estimate the likelihood of specific future mutations, ultimately yielding the numerical odds of one parent strain giving rise to a specific descendant via natural evolutionary processes. After validating our model on the problem of forecasting the dominant strain(s) in the upcoming flu season, with Emergenet-based forecasts significantly outperforming WHO recommended flu vaccine compositions almost consistently over the past two decades for H1N1/H3N2 subtypes, individually in the Northern/Southern hemispheres (HA: 34.8% improvement, NA: 12.5% improvement), we assess the pandemic potential of novel animal strains that do not yet circulate in humans. While the state-of-the-art Influenza Risk Assessment Tool (IRAT) from the CDC comprises multiple time-consuming experimental assays, our proposed E-risk score can be evaluated in  $\approx$  6 seconds for each new strain, while strongly correlating with published IRAT scores (correlation=0.703, p-value = 0.00026). This six orders of magnitude speedup (weeks vs seconds) in identifying risky strains is a necessary step to fully exploit the current surveillance capacity via scalably analyzing tens of thousands of strains collected every year. Applying our tool to all Influenza A strains recently collected in the wild (post-2020, 6066 in total), we identify a small number of strains of diverse subtypes, in diverse hosts and geo-locations, that pose the maximal risk (6 with estimated IRAT emergence score  $>$  6.5). Such scalable risk-ranking can enable follow-up targeted investigations, and meaningful preemptive pandemic mitigation strategies, which are especially relevant in the light of the recent pandemic.

## INTRODUCTION

Influenza viruses constantly evolve<sup>5</sup>, altering surface protein structures over a time scale of months to evade the prevailing host immunity, and cause the recurring seasonal epidemic. These periodic infection peaks claim a quarter to half a million lives<sup>6</sup> globally, and currently our response hinges on inoculating the population with a reformulated vaccine annually<sup>5,7</sup>. Among numerous factors that hinder optimal design of the seasonal flu shot, failing to correctly predict the future dominant strain dramatically reduces vaccine effectiveness<sup>8</sup>. Despite recent advances<sup>6,9</sup> such predictions remain imperfect. In addition to the seasonal epidemic, Influenza A strains spilling over into humans from animal reservoirs have triggered pandemics at least four times (1918 Spanish flu/H1N1, 1957 Asian flu/H2N2, 1968 hongkong flu/H3N2, 2009 swine flu/H1N1) in the past 100 years<sup>1</sup>. With the memory of the sudden emergence of COVID-19 fresh in our minds, a looming question is whether we can preempt and mitigate such events in the future. Influenza A, partly on account of its segmented genome and its wide prevalence in common animal hosts, can easily incorporate genes from multiple strains and (re)emerge as novel human pathogens<sup>3,10</sup>, thus harboring a high pandemic potential.


A possible approach to mitigating such risk is to identify animal strains that do not yet circulate in humans but is likely to spill-over and quickly achieve human-to-human (HH) transmission capability. Despite global surveillance efforts to



**Fig. 1. Emergenet inference and applications.** **a.** Variations of genomes for identical subgroups of Influenza A are analyzed to infer a recursive forest of conditional inference trees<sup>11</sup> – the Emergenet– which maximally captures the emergent dependencies between an a priori unspecified number of mutations, deletions and insertions. With these inferred dependencies we can estimate the numerical odds of specific mutations, and by extension, the numerical value of the probability of one strain giving rise to another in the wild, under complex selection pressures from the background. **b.** Snapshot of decision trees from the Emergenet constructed for H1N1 haemagglutinin 2018 sequences. Note that the decision tree predicting the bases at index 1274 uses the bases at 1064, 1445, 197 as features. These features are automatically selected, as being maximally predictive of the bases be at 1274. Then, we compute predictors for each of these feature indices, e.g. trees for index 1064, which involves index 1314 and 339 as features. Continuing, we find that the trees for index 1314 involves indices 1263, 636 and 21, and that for 1263 involves 1314, 667 and 313. The predictor for 1263 depends on 1314, and that for 1314 depends on 1263, revealing the recursive structure of Emergenet. **c.** First application: With Emergenet induced ability to quantify mutation probabilities, we forecast dominant strain(s) for the next flu season, using only sequences collected in the previous season (and the inferred Emergenet, using data from the past year). **d.** Second application: estimation of the risk of a global pandemic posed by individual animal strains that are still not known to circulate in humans.


collect wild specimens from diverse hosts and geo-locations, our current ability to objectively, reliably and scalably evaluate such risk posed to humans by individual strains is limited<sup>12</sup>. Despite recent progress towards understanding emergence risk<sup>13–15</sup>, scalable ranking of individual strains remains out of reach.

The Center for Disease Control's (CDC) current solution to preempt strain-specific emergence is the Influenza Risk




**Fig. 2. Seasonal predictions for Influenza A.** Relative out-performance of Qnet predictions against WHO recommendations for H1N1 and H3N2 sub-types for the HA and NA coding sequences over the both hemispheres. The negative bars (red) indicate the reduced edit distance between the predicted sequence and the actual dominant strain that emerged that year. Note that the recommendations for the north are given in February, while that for the south are given at the previous December, keeping in mind that the flu season in the south begins a few months early (e.g. for the 2021-2022 flu season, southern data in the table is labelled '2021' and northern is labelled '2022'). **Panels e, f, k, l** show further possible improvement in NA predictions if we return three recommendations instead of one each year.


## A. Predicted emergence risk vs published IRAT scores




## b. Estimating emergence



## C. Estimating impact



## d. Global prediction of IRAT scored for all Influenza A sequences collected since 2020



**Fig. 3. Emergenet based estimation of IRAT score.** Panel a. There is an approximate linear relationship between average q-distance from human circulating strains (geometric mean of the E-distance for HA and NA sequences) and the published IRAT emergence score calculated by CDC. Panel b. Estimation of the IRAT emergence score via fitting a GLM model to the E-distances estimated from the Emergenet. Panel c. Estimation of IRAT impact scores via fitting a separate GLM model to the the E-distances estimated from the Emergenet. Panel d. Identifying risky Influenza A strains amongst those collected between 2020 and 2022 April, via our IRAT emergence score estimation framework.

Assessment Tool (IRAT)<sup>16</sup>, which reflects evaluations by subject matter experts (SME) from the CDC, the Food and Drug Administration (FDA), the Animal and Plant Health Inspection Service (APHIS), and the Agricultural Research Service (ARS). Each SME scores 1-3 elements from a set of ten factors comprising the number of recorded human infections, transmission in laboratory animals, receptor binding characteristics, population immunity, animal infections, genomic analysis, antigenic relatedness, global prevalence, pathogenesis, and antiviral treatment options. The point estimates for each element are averaged across SMEs, scaled by predetermined weights, and summed to give an aggregate score for each of the two IRAT questions: 1) the potential risk for emergence in humans and 2) the potential public




Fig. 4. Standard phylogenetic tree constructed with edit distances, with all Influenza A strains collected between 2020 and 2022 April, with estimated IRAT emergence risk > 6.0, and collapsing leaves which differ by less than 15 edits in the HA, leaving the most risky strains in the leaves. The most risky strains are marked with a red arrowhead, which comes from diverse animal hosts, and geographic regions.

TABLE 1  
Influenza A Strains Evaluated by IRAT and Corresponding Qnet Computed Risk Scores

Influenza Virus	Subtype	IRAT Date	IRAT Emergence Score	IRAT Impact Score	HA Sample	NA Sample	HA Avg. E-distance	NA Avg. E-distance	Geom. Mean	Emergenet Emergence Score	Emergenet Impact Score
A/swine/Shandong/1207/2016	H1N1	Jul 2020	7.5	6.9	1000	1000	0.0941	0.0205	0.0440	6.0	6.2
A/Ohio/13/2017	H3N2	Jul 2019	6.6	5.8	1000	1000	0.0184	0.0306	0.0238	6.3	6.2
A/Hong Kong/125/2017	H7N9	May 2017	6.5	7.5	437	437	0.0296	0.0058	0.0131	6.6	6.5
A/Shanghai/02/2013	H7N9	Apr 2016	6.4	7.2	178	178	0.0055	0.0036	0.0044	6.7	6.6
A/Anhui-Lujiang/39/2018	H9N2	Jul 2019	6.2	5.9	31	30	0.0290	0.1681	0.0698	5.2	5.0
A/Indiana/08/2011	H3N2	Dec 2012	6.0	4.5	1000	1000	0.0523	0.0091	0.0218	6.4	6.5
A/California/62/2018	H1N2	Jul 2019	5.8	5.7	55	55	0.1089	0.0610	0.0815	5.4	5.5
A/Bangladesh/0994/2011***	H9N2	Feb 2014	5.6	5.4			0.2078	0.1823	0.1947	4.3	4.9
A/Sichuan/06681/2021	H5N6	Oct 2021	5.3	6.3	45	45	0.3616	0.0518	0.1369	5.2	6.4
A/Vietnam/1203/2004	H5N1	Nov 2011	5.2	6.6	258	246	0.1673	0.0111	0.0430	6.2	6.7
A/Yunnan/14564/2015**	H5N6	Apr 2016	5.0	6.6	344	331	0.3482	0.2987	0.3225	4.9	6.5
A/Astrakhan/3212/2020**	H5N8	Mar 2021	4.6	5.2	381	365	0.1603	0.3472	0.2359	3.9	4.4
A/Netherlands/219/2003	H7N7	Jun 2012	4.6	5.8	46	46	0.2757	0.3521	0.3115	4.6	5.8
A/American wigeon/South Carolina/AH0195145/2021	H5N1	Mar 2022	4.4	5.1	335	323	0.1722	0.5114	0.2967	4.0	4.7
A/Jiangxi-Donghu/346/2013***	H10N8	Feb 2014	4.3	6.0			0.2088	0.2101	0.2094	4.3	4.8
A/gyrfalcon/Washington/41088/2014**	H5N8	Mar 2015	4.2	4.6	341	328	0.1532	0.3424	0.2290	3.9	4.3
A/Northern pintail/Washington/40964/2014**	H5N2	Mar 2015	3.8	4.1	341	328	0.1529	0.3799	0.2410	3.9	4.3
A/canine/Illinois/12191/2015	H3N2	Jun 2016	3.7	3.7	1000	1000	0.0607	0.1509	0.0957	4.9	4.8
A/American green-winged teal/Washington/1957050/2014	H5N1	Mar 2015	3.6	4.1	326	314	0.1911	0.4482	0.2927	4.1	4.9
A/turkey/Indiana/1573-2/2016**	H7N8	Jul 2017	3.4	3.9	495	494	0.1130	0.7738	0.2957	3.4	3.9
A/chicken/Tennessee/17-007431-3/2017	H7N9	Oct 2017	3.1	3.5	496	495	0.1027	0.2569	0.1624	4.1	4.2
A/chicken/Tennessee/17-007147-2/2017	H7N9	Oct 2017	2.8	3.5	496	495	0.2095	0.2541	0.2307	4.2	4.8

\*\* Emergenet constructed using all human strains that match the HA sub-type, e.g., H5Nx for H5N6.

\*\*\* distance estimated averaging over those obtained by considering all Emergenets from other subtypes.

health impact if the virus gains HH capability. The scores are potentially subjective, and involve multiple experimental assays, possibly taking weeks to compile for a single strain or lineage. This results in a scalability bottleneck: with global efforts collecting thousands of sequences annually, IRAT assessment protocols are not fast enough to leverage the full capacity of current surveillance output.

Here we introduce a pattern recognition algorithm to automatically parse out emergent evolutionary constraints operating on Influenza A viruses in the wild, to numerically estimate the probability  $Pr(x \rightarrow y)$  of a strain  $x$  spontaneously giving rise to  $y$ . We show that this capability is key to preempting strains which are expected to be in future circulation, and thus 1) reliably forecast dominant strains of seasonal epidemics, and 2) approximate IRAT scores of non-human strains without experimental assays or direct involvement of SMEs.

To uncover relevant evolutionary constraints, we analyzed observed variations (point substitutions and indels) of the amino acid (AA) sequences of the two key proteins implicated in cellular entry and exit<sup>1,17</sup>, namely HA and NA respectively. By representing these constraints within a predictive framework – the Emergenet– we estimated the odds of a specific mutation to arise in a given strain, and consequently the probability of a specific strain spontaneously evolving into another. Such explicit calculations are difficult without first inferring the variation of mutational probabilities and the potential residue replacements from one positional index to the next along the residue sequence of a protein. The many well-known classical DNA substitution models<sup>18</sup> or standard approaches to phylogenetic tree inference which assume a constant mutation rate or some variation thereof, do not address these issues, and are not applicable

TABLE 2  
Influenza A Strains Evaluated by IRAT and Corresponding Qnet Computed Risk Scores

strain	sub-type	HA accession	NA accession	predicted IRAT impact	predicted IRAT emergence
A/swine/Missouri/A02524711/2020	H1N1	EPI1818121	EPI1818122	6.6318	6.7067
A/Camel/Inner_Mongolia/XL/2020	H7N9	EPI2026200	EPI2026202	6.6286	6.7026
A/swine/Indiana/A02524710/2020	H3N2	EPI1818137	EPI1818138	6.6070	6.6757
A/swine/North_Carolina/A02479173/2020	H1N1	EPI1780425	EPI1780426	6.5878	6.6517
A/swine/Tennessee/A02524414/2022	H1N1	EPI2149257	EPI2149258	6.5379	6.5893
A/swine/Minnesota/A02635976/2021	H1N1	EPI1912208	EPI1912209	6.4841	6.5220
A/swine/Chile/VN1401-5054/2020	H3N2	EPI1974975	EPI1974978	6.4578	6.4891
A/swine/Italy/56910/2020	H3N2	EPI2142217	EPI2142173	6.4537	6.4840
A/swine/Minnesota/A02245643/2020	H3N2	EPI1769178	EPI1769179	6.4370	6.4631
A/swine/Iowa/A02479005/2020	H1N1	EPI1777621	EPI1777622	6.4103	6.4297
A/swine/Iowa/A02524874/2020	H3N2	EPI1907838	EPI1907839	6.3836	6.3962
A/swine/Indiana/A02636638/2022	H1N1	EPI2153370	EPI2153371	6.3830	6.3954
A/swine/Illinois/A02479007/2020	H3N2	EPI1777629	EPI1777630	6.3769	6.3877
A/swine/Spain/44579-1/2020	H3N2	EPI1930744	EPI1930748	6.3737	6.3838
A/swine/Minnesota/A02248037/2021	H1N1	EPI1912188	EPI1912189	6.3694	6.3784
A/swine/Minnesota/A02245699/2020	H3N2	EPI1833007	EPI1833008	6.3692	6.3781
A/swine/Iowa/A02635917/2021	H1N1	EPI1911753	EPI1911754	6.3687	6.3774
A/swine/Minnesota/A02711801/2022	H1N1	EPI2153420	EPI2153421	6.3674	6.3759
A/swine/Illinois/A02635936/2021	H1N1	EPI1911791	EPI1911792	6.3671	6.3754
A/swine/South_Dakota/A02524453/2020	H1N1	EPI1765555	EPI1765556	6.3658	6.3738
A/swine/Minnesota/A02248061/2021	H1N1	EPI1912494	EPI1912495	6.3624	6.3696
A/swine/Iowa/A02636439/2022	H1N1	EPI2147475	EPI2147476	6.3616	6.3686
A/swine/Iowa/A02524875/2020	H1N1	EPI1907858	EPI1907859	6.3610	6.3678
A/swine/Minnesota/A02248060/2021	H1N1	EPI1912500	EPI1912501	6.3592	6.3656
A/swine/Nebraska/A02636117/2021	H1N1	EPI1932937	EPI1932938	6.3582	6.3644
A/swine/Iowa/A02524513/2020	H1N1	EPI1832647	EPI1832648	6.3580	6.3641
A/swine/Iowa/A02524724/2020	H1N1	EPI1818387	EPI1818388	6.3579	6.3640
A/swine/Iowa/A02635719/2021	H1N1	EPI1910907	EPI1910908	6.3579	6.3640
A/swine/Nebraska/A02479337/2020	H1N1	EPI1769116	EPI1769117	6.3579	6.3640
A/swine/Iowa/A02479383/2020	H1N1	EPI1771027	EPI1771028	6.3579	6.3640
A/swine/Nebraska/A02479212/2020	H1N1	EPI1775884	EPI1775885	6.3579	6.3640
A/swine/Minnesota/A02245424/2020	H1N1	EPI1780207	EPI1780208	6.3579	6.3640
A/swine/Minnesota/A02479051/2020	H1N1	EPI1778572	EPI1778573	6.3579	6.3640
A/swine/Missouri/A02525065/2021	H1N1	EPI1908581	EPI1908582	6.3579	6.3640
A/swine/Missouri/A02524951/2020	H1N1	EPI1908429	EPI1908430	6.3579	6.3640
A/swine/Iowa/A02524892/2020	H1N1	EPI1907881	EPI1907882	6.3579	6.3640
A/swine/Nebraska/A02524954/2020	H1N1	EPI1908393	EPI1908394	6.3579	6.3640
A/swine/Iowa/A02524994/2020	H1N1	EPI1908427	EPI1908428	6.3579	6.3640
A/swine/Iowa/A02525313/2021	H1N1	EPI1910761	EPI1910762	6.3579	6.3640
A/swine/Iowa/A02524646/2020	H1N1	EPI1817164	EPI1817165	6.3579	6.3640
A/swine/Nebraska/A02479186/2020	H1N1	EPI1774141	EPI1774142	6.3578	6.3638
A/swine/Iowa/A02479156/2020	H1N1	EPI1780249	EPI1780250	6.3574	6.3634
A/swine/Iowa/A02479229/2020	H1N1	EPI1775914	EPI1775915	6.3574	6.3633
A/swine/Iowa/A02479303/2020	H1N1	EPI1768639	EPI1768640	6.3567	6.3625
A/swine/Minnesota/A02710691/2021	H1N1	EPI2146090	EPI2146091	6.3561	6.3618
A/swine/Iowa/A02635881/2021	H1N1	EPI1911668	EPI1911669	6.3560	6.3616
A/swine/Iowa/A02525354/2021	H1N1	EPI1910789	EPI1910790	6.3560	6.3616
A/swine/Iowa/A02524739/2020	H1N1	EPI1818383	EPI1818384	6.3543	6.3595
A/swine/Iowa/A02635823/2021	H1N1	EPI1911263	EPI1911264	6.3543	6.3594
A/swine/Minnesota/A02711797/2022	H3N2	EPI2153382	EPI2153383	6.3537	6.3587
A/swine/Iowa/A02479141/2020	H1N1	EPI1780241	EPI1780242	6.3531	6.3579
A/swine/Iowa/A02635955/2021	H1N1	EPI1912240	EPI1912241	6.3516	6.3561
A/swine/Iowa/A02750621/2022	H1N1	EPI2161576	EPI2161577	6.3509	6.3552
A/swine/Illinois/A02525253/2021	H3N2	EPI1910375	EPI1910376	6.3506	6.3548
A/swine/Iowa/A02245587/2020	H1N1	EPI1775817	EPI1775818	6.3497	6.3537
A/swine/Iowa/A02636145/2021	H1N1	EPI1932055	EPI1932930	6.3487	6.3525
A/swine/Iowa/A02636114/2021	H1N1	EPI1931853	EPI1931854	6.3487	6.3525
A/swine/Iowa/A02525217/2021	H1N1	EPI1909087	EPI1909088	6.3487	6.3525
A/swine/Iowa/A02636496/2022	H1N1	EPI2148086	EPI2148087	6.3485	6.3522
A/swine/Iowa/A02635871/2021	H1N1	EPI1911656	EPI1911657	6.3473	6.3507
A/swine/Iowa/A02479067/2020	H1N1	EPI1778734	EPI1778735	6.3469	6.3501
A/swine/Minnesota/A02246459/2021	H1N1	EPI1912518	EPI1912519	6.3465	6.3496
A/swine/Minnesota/A02525348/2021	H1N1	EPI1910795	EPI1910796	6.3400	6.3415
A/swine/Iowa/A02479343/2020	H1N1	EPI1769114	EPI1769115	6.3387	6.3399
A/canine/Texas/21-011409-001/2021	H3N2	EPI1896555	EPI1896557	6.3093	6.3030
A/swine/Kansas/A02245381/2020(H1N1)	H1N1	EPI1777723	EPI1777724	6.2958	6.2860
A/swine/Iowa/A02246996/2021	H1N1	EPI2146133	EPI2146134	6.2817	6.2684
A/mink/China/chick_embryo/2020	H9N2	EPI2161544	EPI2161548	6.2787	6.2646

to the problem at hand. Similarly, recently reported algorithms such as FluLeap<sup>19</sup> which identifies host tropism from sequence data, or estimating risk posed by different viral species<sup>15</sup> do not allow strain-specific risk assessment.




TABLE 3  
Count of identified strains above estimated emergence risk threshold

subtype	No. with minimum estimated IRAT emergence score				
	6.0	6.2	6.3	6.4	6.5
H1N1	62 (83%)	57 (81%)	53 (82%)	5 (50%)	4 (67%)
H3N2	11 (15%)	11 (16%)	11 (17%)	4 (40%)	1 (17%)
H7N9	1 (1%)	1 (1%)	1 (1%)	1 (10%)	1 (17%)
H9N2	1 (1%)	1 (1%)	0	0	0

The genomic dependencies we uncover are shaped by a functional necessity of conserving/augmenting fitness in the wild. A strain must be sufficiently common to be recorded, implying that the sequences from public databases that we train with have high replicative fitness. Lacking kinetic proofreading in RNA-polymerase, Influenza A integrates faulty nucleotides at a relatively high rate ( $10^{-3} - 10^{-4}$ ) during replication<sup>20,21</sup>. However, few of these variations are actually viable, leading to emergent dependencies between such changes. Furthermore, these fitness constraints are not time-invariant. The background distribution of strains, and selection pressure from the evolution of cytotoxic T lymphocyte epitopes<sup>22-26</sup> in humans can change quickly. With a sufficient number of unique samples to train on for each flu season, the Emergenet (recomputed for each time-period) is expected to automatically reflect the effect of an evolving host immunity, and the current background strain distribution.

Structurally, an Emergenet comprises an interdependent collection of local predictors: each aiming to predict the residue at a particular index using as features (input variables) the residues appearing at other indices (Fig. 1b). Thus, an Emergenet comprises almost as many such position-specific predictors as the length of the sequence. These individual predictors are implemented as conditional inference trees<sup>11</sup>, in which nodal splits have a minimum pre-specified significance in differentiating the child nodes. Inferring residue predictors at each index as functions of the rest of the sequence yields an estimated conditional residue distribution at each index. The set of indices acting as features in each tree varies, *e.g.*, in the fragment of the H1N1 HA Emergenet (2020-2021, Fig 1b), the predictor for residue at index 63 is dependent on residue 155, and the predictor for 155 is dependent on 223, the predictor for 223 is dependent on 14, and the residue at 14 is dependent on 63, revealing a cyclic dependency. The complete Emergenet harbors a vast number of such relationships, wherein each internal node of a tree may be “expanded” to its own tree. Owing to this recursive expansion, a complete Emergenet is hard to visualize; nevertheless it captures the complexity of the rules guiding evolutionary change substantially better than earlier attempts, as evidenced by our validation results.

In this study we used HA and NA sequences from all unique Influenza A strains in the NCBI and GISAID databases recorded between 2000-2022 (98,299 strains in total). We construct Emergenets separately for H1N1 and H3N2 subtypes, and for each flu season, constructing in total 85 models for predicting seasonal dominance. We use no detailed phenotypic characteristics of the strains, which is advantageous since antigenic characterization tend to be substantially laborious and low-throughput compared to genome sequencing<sup>27</sup>; however, incorporation of deep mutational scanning (DMS) assays have been shown to improve prediction of seasonal strains<sup>6</sup>. Despite limiting ourselves to only genotypic information (and subtypes), our approach is able to distill emergent fitness-preserving constraints outperforming reported DMS augmented strategies for Influenza A.

Inference of the Emergenet components is the first step in our approach, which then induces an intrinsic distance metric between strains. The E-distance (Eq. (5) in Online Methods) is defined as the square-root of the Jensen-Shannon (JS) divergence<sup>28</sup> of the conditional residue distributions, averaged over the sequence. Unlike the classical approach of measuring the number of edits between sequences, the E-distance is informed by the Emergenet-inferred dependencies, and adapts to the specific subtype, allele frequencies, and environmental variations. Central to our approach is the theoretical result (Theorem 1 in Online Methods) that  $\log Pr(x \rightarrow y)$  may be approximated by the E-distance  $\theta(x, y)$ . The mathematical intuition relating E-distance to the loglikelihood of spontaneous change is similar to quantifying the odds of a rare biased outcome when we toss a fair coin<sup>29</sup>. Importantly, unlike the edit distance, the E-distance between two fixed sequences may change even if only the background strain population changes (SI-Table ?? illustrates that the distance between fixed sequences vary with collection years).

Determining the numerical odds of spontaneous jump  $Pr(x \rightarrow y)$  (Fig. 1) allows us to frame the problem of forecasting dominant strain(s), and that of estimating the pandemic potential of an animal strain as mathematical propositions (albeit with some simplifying assumptions), with approximate solutions (Fig. 1c-d). Thus, a dominant strain for an upcoming season may be identified as one which maximizes the joint probability of simultaneously arising from each (or most) of the currently circulating strains (Fig. 1c). This does not deterministically specify the dominant strain, but a strain satisfying this criterion has high odds of emerging as the dominant one. And, a pandemic risk score of a novel strain may be estimated by the probability of it giving rise to a well-adapted human strain.

In the context of forecasting future dominant strain(s), we derive a search criteria (See Online Methods) from the above proposition, to identify historical strain(s) that are expected to be close to the next dominant strain(s):

$$x_*^{t+\delta} = \arg \min_{y \in \cup_{\tau \leq t} H^\tau} \left( \sum_{x \in H^t} \theta^{[t]}(x, y) - |H^t| A \ln \omega_y \right) \quad (1)$$

---

where  $x_*^{t+\delta}$  is a predicted dominant strain at time  $t + \delta$ ,  $H^t$  is the set of currently circulating human strains at time  $t$  observed over the past year,  $\theta^{[t]}$  is the E-distance informed by the inferred Emergenet using sequences in  $H^t$ ,  $\omega_y$  is the estimated probability of strain  $y$  being generated by the Emergenet, and  $A$  is a constant dependent on the sequence length and significance threshold used (See Online Methods). The first term gets the solution close to the centroid of the current strain distribution (in the E-distance metric, which is different from the centroid if the standard edit distance is used), and the second term relates to how common the genomic patterns are amongst recent human strains.

Prediction of the future dominant strain as a close match to a historical strain allows out-of-sample validation of the approach with past World Health Organization (WHO) recommendations for the flu shot, which is currently reformulated about six months in advance based on a recommended cocktail of historical strains determined via global surveillance<sup>30</sup>. For each year of the past two decades, we calculated strain forecasts using Eq. (12) with data available six months before the target season. We measured forecast performance by the number of mutations by which the predicted HA/NA sequences deviated from the dominant strain. Our Emergenet-informed forecasts outperform WHO/CDC recommended flu vaccine compositions almost consistently over the past two decades, for both H1N1 and H3N2 subtypes, individually in the northern and the southern hemispheres (which have distinct recommendations). The results broken down by hemisphere/protein/subtypes is given in Table 1. Fig. 2 illustrates the relative gains computed for both subtypes and the two hemispheres. Additional improvement is possible if we recommend multiple strains every season for the vaccine cocktail (Fig. 2e,f,k,l). The details of the specific strain recommendations made by the Emergenet approach for two subtypes (H1N1, H3N2), for two genes (HA, NA) and for the northern and the southern hemispheres over the previous two decades are enumerated in the Supplementary Tables SI-Table ?? through SI-Table 8. While it is recognized that even well-matched strains can fail to induce a strong immune response due to previous infection history of vaccine recipients<sup>31</sup>, strain-matching is a crucial component to realizing high vaccine effectiveness<sup>32</sup>, and we outperform the current practice (WHO/CDC) as well as recently reported prediction strategies using more standard computational and/or experimental frameworks<sup>6,9</sup>.

Our primary claim, however, is the ability to estimate the pandemic potential of novel animal strains, via a time-varying E-risk score  $\rho_t(x)$  for a strain  $x$  not yet found to circulate in human hosts. We show that (See Online Methods):

$$\rho_t(x) \triangleq -\frac{1}{|H^t|} \sum_{y \in H^t} \theta^{[t]}(x, y) \quad (2)$$

scales as the average log-likelihood of  $Pr(x \rightarrow y)$  where  $y$  is any human strain of a similar subtype to  $x$ , and  $\theta^{[t]}$  is the E-distance informed by the Emergenet computed from recent human strains  $H_t$  at time  $t$  of the same subtype as  $x$ , observed over the past year. As before, the Emergenet inference makes it possible to estimate  $\rho_t(x)$  explicitly.

To validate our score against CDC-estimated IRAT emergence scores, we construct Emergenet models for HA and NA sequences using subtype-specific human strains, typically collected within the past year of the assessment date, *e.g.*, the IRAT assessment date for A/swine/Shandong/1207/2016 is July 2020, and we use human H1N1 strains collected between 1/7/2019-6/30/2020 for the Emergenet construction. For sub-types with very few recorded human strains (H1N2, H5N1, H5N6, H7N7, H9N2), we consider all subtype-specific human strains collected upto the IRAT assessment date to construct our Emergenet. We then compute the average E-distance between the animal strain of interest and the recent human strains for both HA and NA sequences (using Eq. (2)), finally reporting their geometric mean as our estimated risk. Considering IRAT scores of 22 strains published by the CDC, we find strong out-of-sample support (correlation of  $-0.704$ , pvalue  $< 0.00026$ , Fig. 3, see Online Methods) for this claim. Importantly, each E-risk score is computable in approximately 6 seconds as opposed to potentially weeks taken by IRAT experimental assays.

The time-dependence of the E-risk reflects the impact of the background (See Table ?? and SI-Fig. 3), and we show that recomputing the risks using Emergenets constructed from the recent circulating strains instead of using those from when the IRAT assessments took place at the CDC, worsens the correlation ( $-0.59$ , p-value 0.003).

To map the Emergenet distances to the more recognizable IRAT scores, we train a general linear model (GLM) from the HA/NA-based E-risk values (See Online Methods). Since the CDC-estimated IRAT impact scores are strongly correlated with their IRAT emergence scores (correlation of 0.8015), we also trained a separate GLM to estimate the impact score from the E-risk values, despite our theoretical intuition primarily supporting the emergence phenomenon. With these maps we estimate the IRAT scores of all distinct Influenza A strains collected between 2020 through 2022 April (6066 strains in total), and identify the ones posing maximal risk (Fig. 3c). 1,773 strains have a predicted emergence score  $> 6.0$ . However, many of these strains are highly similar, differing by only a few edits. To identify the sufficiently distinct risky strains, we constructed the standard phylogeny from HA sequences with score  $> 6$  (Fig. 4), and collapsed all leaves within 15 edits, showing only the most risky strain within a collapsed group. This leaves 75 strains as shown in Fig. 4 (68 with emergence risk  $> 6.25$ , and 6 with emergence risk above 6.5, see Table 2). A substantial number of risky strains are H1N1, followed by H3N2, with a small number of H7N9 and H9N2. Five risky strains with emergence score  $> 6.58$  are identified to be: A/swine/Missouri/A02524711/2020 (H1N1), A/Camel/Inner\_Mongolia/XL/2020 (H7N9), A/swine/Indiana/A02524710/2020 (H3N2), A/swine/North Carolina/A02479173/2020 (H1N1), and A/swine/Tennessee/A02524414/2022 (H1N1). Additionally, A/mink/China/chick embryo/2020 (H9N2), with a lower estimated emergence score (6.26) is also important, as the most risky H9N2 strain in our analysis. We compare the HA sequences along with two dominant human strains in 2021-2022 season (See Fig. 5), which shows substantial residue replacements, in and out of the receptor binding domain (RBD).

Swines are known to be efficient mixing vessels<sup>3,33?,34</sup>, and hence unsurprisingly host a large fraction of the risky strains (> 80% over 6.0, to over 50% over 6.5). Also, as expected, most of these swine strains are of H1N1 subtype, with the other subtypes having emerged into humans more recently. Our finding that a H7N9 poses substantial risk is likewise not surprising: HH transmission has been suspected in asian-lineage H7N9 strains, and are rated by IRAT as having the greatest potential to cause a pandemic<sup>35</sup>. The finding of the most risky H9N2 strain in a mink is again unsurprising, in the light of these hosts been recently suggested as efficient mixing vessels to breed human-compatible strains<sup>36</sup>. Thus, qualitatively results are well aligned with the current expectations; nevertheless the ability to quantitatively rank specific strains which pose maximal risk is a crucial new capability enabling proactive pandemic mitigation efforts.

In conclusion, while numerous tools exist for ad hoc quantification of genomic similarity<sup>9,18,37–40</sup>, a smaller distance i.e. a higher similarity of two strains in these frameworks is not sufficient to imply a high likelihood of a jump. To the best of our knowledge, the Emergenet algorithm is the first of its kind to learn an appropriate biologically meaningful comparison metric from data, without assuming any model of DNA or amino acid substitution, or a genealogical tree *a priori*. While the effect of the environment and selection cannot be inferred from a single sequence, an entire database of observed strains, processed through the right lens, can parse out useful predictive models of these complex interactions. Our results are aligned with recent studies demonstrating effective predictability of future mutations for different organisms<sup>41,42</sup>. Our approach is currently limited by the fact that the E-distance calculation is currently applicable to analogous sequences (such as point variations of the same protein from different viral subtypes), and the Emergenet inference requires a sufficient diversity of observed strains. A multi-variate regression analysis indicates that the most important factor for our approach to succeed is the diversity of the sequence dataset (see Supplementary Table 12), which would exclude applicability to completely novel pathogens with no related human variants, and ones that evolve very slowly. Nevertheless, the tools reported here can improve effectiveness of the annual flu shot, and perhaps allow for the development of preemptive vaccines to target risky animal strains before the first human infection in the next pandemic.

## ONLINE METHODS

Next, we briefly describe the details of the proposed computational framework.

## EMERGENET FRAMEWORK

We do not assume that the mutational variations at the individual indices of a genomic sequence are independent (See Fig 1a). Irrespective of whether mutations are truly random<sup>43</sup>, since only certain combinations of individual mutations are viable, individual mutations across a genomic sequence replicating in the wild appear constrained, which is what is explicitly modeled in our approach.

Consider a set of random variables  $X = \{X_i\}$ , with  $i \in \{1, \dots, N\}$ , each taking value from the respective sets  $\Sigma_i$ . Here each  $X_i$  is the random variable modeling the “outcome” i.e. the AA residue at the  $i^{th}$  index of the protein sequence. A sample  $x \in \prod_1^N \Sigma_i$  is an ordered  $N$ -tuple, which is a specific strain in this context, consisting of a realization of each of the variables  $X_i$  with the  $i^{th}$  entry  $x_i$  being the realization of random variable  $X_i$ .

We use the notation  $x_{-i}$  and  $x^{i,\sigma}$  to denote:

$$x_{-i} \triangleq x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_N \quad (3a)$$

$$x^{i,\sigma} \triangleq x_1, \dots, x_{i-1}, \sigma, x_{i+1}, \dots, x_N, \sigma \in \Sigma_i \quad (3b)$$

Also,  $\mathcal{D}(S)$  denotes the set of probability measures on a set  $S$ , e.g.,  $\mathcal{D}(\Sigma_i)$  is the set of distributions on  $\Sigma_i$ .

We note that  $X$  defines a random field<sup>44</sup> over the index set  $\{1, \dots, N\}$ .

**Definition 1** (Emergenet). *For a random field  $X = \{X_i\}$  indexed by  $i \in \{1, \dots, N\}$ , the Emergenet is defined to be the set of predictors  $\Phi = \{\Phi_i\}$ , i.e., we have:*

$$\Phi_i : \prod_{j \neq i} \Sigma_j \rightarrow \mathcal{D}(\Sigma_i), \quad (4)$$

where for a sequence  $x$ ,  $\Phi_i(x_{-i})$  estimates the distribution of  $X_i$  on the set  $\Sigma_i$ .

We use conditional inference trees as models for predictors<sup>11</sup>, although more general models are possible.

## Biology-Aware Distance Between Sequences

The mathematical form of our metric is not arbitrary; JS divergence is a symmetricised version of the more common KL divergence<sup>28</sup> between distributions, and among different possibilities, the E-distance is the simplest metric such that the likelihood of a spontaneous jump (See Eq. (8) in Methods) is provably bounded above and below by simple exponential functions of the E-distance.

---

**Definition 2** (E-distance: adaptive biologically meaningful dissimilarity between sequences). *Given two sequences  $x, y \in \prod_1^N \Sigma_i$ , such that  $x, y$  are drawn from the populations  $P, Q$  inducing the Emergenet  $\Phi^P, \Phi^Q$ , respectively, we define a pseudo-metric  $\theta(x, y)$ , as follows:*

$$\theta(x, y) \triangleq \mathbf{E}_i \left( \mathbb{J}^{\frac{1}{2}} \left( \Phi_i^P(x_{-i}), \Phi_i^Q(y_{-i}) \right) \right) \quad (5)$$

where  $\mathbb{J}(\cdot, \cdot)$  is the Jensen-Shannon divergence<sup>45</sup> and  $\mathbf{E}_i$  indicates expectation over the indices.

The square-root in the definition arises naturally from the bounds we are able to prove, and is dictated by the form of Pinsker's inequality<sup>28</sup>, ensuring that the sum of the length of successive path fragments equates the length of the path.

## Membership Degree

For our modeling to be reliable, we need a quantitative test of how well the Emergenet represents the data. Here, we formulate an explicit membership test to ascertain if individual samples may indeed be generated by the Emergenet with sufficiently high probability.

**Definition 3** (Membership probability of a sequence). *Given a population  $P$  inducing the Emergenet  $\Phi^P$  and a sequence  $x$ , we can compute the membership probability of  $x$ :*

$$\omega_x^P \triangleq \Pr(x \in P) = \prod_{j=1}^N (\Phi_j^P(x_{-j})|_{x_j}) \quad (6)$$

$x_j$  is the  $j^{th}$  entry in  $x$ , and is thus an element in the set  $\Sigma_j$ . Since we are mostly concerned with the case where  $\Sigma_j$  is a finite set,  $\Phi_j^P(x_{-j})|_{x_j}$  is the entry in the probability mass function corresponding to the element of  $\Sigma_j$  which appears at the  $j^{th}$  index in sequence  $x$ .

We can carry out this calculation for a sequence  $x$  known to be in the population  $P$  as well, which allows us to define the membership degree  $\omega_x^P$ .

**Definition 4** (Membership degree). *Let  $X$  be a random field representing a population  $P$ , ie..  $X = x$  is a randomly drawn sequence from  $P$ . Then the membership degree  $\omega^P$  is a function of the random variable  $X$ :*

$$\omega^P(X) \triangleq \prod_{j=1}^N (\Phi_j^P(X_{-j})|_{X_j}) \quad (7)$$

Note that  $\omega^P$  takes values in the unit interval  $[0, 1]$ , and the probability  $x$  is a member of the population  $P$  is  $\omega^P(X = x)$ , denoted briefly as  $\omega_x^P$  or  $\omega_x$  if  $P$  is clear from context.

Since  $\omega^P(X)$  is a random variable, we can now compute sets of sequences that better represent the population  $P$ , and ones that are on the fringe. We can also evaluate using a pre-specified significance-level if a particular sequence is not from the population  $P$ .

## Theoretical Probability Bounds

The Emergenet framework allows us to rigorously compute bounds on the probability of a spontaneous change of one strain to another, brought about by chance mutations. While any sequence of mutations is equally likely, the “fitness” of the resultant strain, or the probability that it will even result in a viable strain, or not. Thus the necessity of preserving function dictates that not all random changes are viable, and the probability of observing some trajectories through the sequence space are far greater than others. The Emergenet framework allows us to explore this constrained dynamics, as revealed by a sufficiently large set of genomic sequences.

We show in Theorem 1 in the supplementary text that at a significance level  $\alpha$ , with a sequence length  $N$ , the probability of spontaneous jump of sequence  $x$  from population  $P$  to sequence  $y$  in population  $Q$ ,  $\Pr(x \rightarrow y)$ , is bounded by:

$$\omega_y^Q e^{-\frac{\sqrt{8}N^2}{1-\alpha}\theta(x, y)} \geq \Pr(x \rightarrow y) \geq \omega_y^Q e^{-\frac{\sqrt{8}N^2}{1-\alpha}\theta(x, y)} \quad (8)$$

where  $\omega_y^Q$  is the membership probability of strain  $y$  in the target population,  $N$  is the sequence length, and  $\alpha$  is the statistical significance level.

## Problem 1: Predicting Dominant Seasonal Strains

Analyzing the distribution of sequences observed to circulate in the human population at the present time allows us to forecast dominant strain(s) in the next flu season as follows:

Let  $x_*^{t+\delta}$  be a dominant strain in the upcoming flu season at time  $t + \delta$ , where  $H^t$  is the set of observed strains presently in circulation in the human population (at time  $t$ ). We will assume that the Emergenet is constructed using the sequences in the set  $H^t$ , and remains unchanged up to  $t + \delta$ . Since this set is a function of time, the inferred Emergenet also changes with time, and the induced E-distance is denoted as  $\theta^{[t]}(\cdot, \cdot)$ .

From the RHS bound established in Theorem 1 (See Eq. (8) above) in the supplementary text, we have:

$$\ln \frac{Pr(x \rightarrow x^{t+\delta})}{\omega_{x^{t+\delta}}} \geq -\frac{\sqrt{8N^2}}{1-\alpha} \theta^{[t]}(x, x^{t+\delta}) \quad (9)$$

$$\Rightarrow \sum_{x \in H^t} \ln \frac{Pr(x \rightarrow x^{t+\delta})}{\omega_{x^{t+\delta}}} \geq \sum_{x \in H^t} -\frac{\sqrt{8N^2}}{1-\alpha} \theta^{[t]}(x, x^{t+\delta}) \quad (10)$$

$$\Rightarrow \sum_{x \in H^t} \theta^{[t]}(x, x^{t+\delta}) - |H^t| A \ln \omega_{x^{t+\delta}} \geq A \ln \frac{1}{\prod_{x \in H^t} Pr(x \rightarrow x^{t+\delta})} \quad (11)$$

where  $A = \frac{1-\alpha}{\sqrt{8N^2}}$ , where  $N$  is the sequence length considered, and  $\alpha$  is a fixed significance level. Since minimizing the LHS maximizes the lower bound on the probability of the observed strains simultaneously giving rise to  $x^{t+\delta}$ , a dominant strain  $x_*^{t+\delta}$  may be estimated as a solution to the optimization problem:

$$x_*^{t+\delta} = \arg \min_{y \in \cup_{\tau \leq t} H^\tau} \sum_{x \in H^t} \theta^{[t]}(x, y) - |H^t| A \ln \omega_y \quad (12)$$

## Problem 2: Measure of Pandemic Potential

We measure the potential of an animal strain  $x_a^t$  to spillover and become HH capable as a human strain  $x_h^{t+\delta}$ , via the proposed E-risk defined as follows:

$$\rho(x_a^t) \triangleq -\frac{1}{|H^t|} \sum_{x \in H^t} \theta^{[t]}(x_a^t, x) \quad (13)$$

where as before  $H^t$  is the set of human strains observed recently (we take this as strains collected within the past year), and  $\theta^{[t]}$  is the E-distance induced by the Emergenet computed from the sequences in  $H^t$ .

The intuition here is that a lower bound of  $\rho(x_a^t)$  scales as average log-likelihood of the  $x_a^t$  giving rise to a human strains in circulation at time  $t$ . Since the strains in  $H^t$  are already HH capable, a high average likelihood of producing a similar strain has a high potential of being a HH capable novel variant, which is a necessary condition of a pandemic strain. To establish the lower bound, we note that from Theorem 1 (See Eq. (8) above) in the supplementary text, we have:

$$\sum_{y \in H^t} \ln \left| \frac{Pr(x_a^t \rightarrow y)}{\omega_y} \right| \leq -\frac{\sqrt{8N^2}}{1-\alpha} |H^t| \rho(x_a^t) \quad (14)$$

Denoting,  $A = \frac{1-\alpha}{\sqrt{8N^2}}$ ,  $A \ln(\prod_{y \in H^t} \omega_y) = C$ , and  $\langle \cdot \rangle$  as the geometric mean function, we have:

$$\Rightarrow \rho(x_a^t) \geq A \ln \left( \prod_{y \in H^t} Pr(x_a^t \rightarrow y) \right)^{1/|H^t|} + C \quad (15)$$

$$\Rightarrow \rho(x_a^t) \geq A \ln \langle Pr(x_a^t \rightarrow x_h^{t+\delta}) \rangle + C \quad (16)$$

Noting that  $A, C$  are not functions of  $x_a^t$ , we conclude that a lower bound of the proposed risk measure  $\rho(\cdot)$  scales with the average loglikelihood of producing strains close to a circulating human strain at the current time.

## Proof of Probability Bounds

**Theorem 1** (Probability bound). *Given a sequence  $x$  of length  $N$  that transitions to a strain  $y \in Q$ , we have the following bounds at significance level  $\alpha$ .*

$$\omega_y^Q e^{\frac{\sqrt{8N^2}}{1-\alpha} \theta(x, y)} \geq Pr(x \rightarrow y) \geq \omega_y^Q e^{-\frac{\sqrt{8N^2}}{1-\alpha} \theta(x, y)} \quad (17)$$

where  $\omega_y^Q$  is the membership probability of strain  $y$  in the target population  $Q$  (See Def. 3), and  $\theta(x, y)$  is the q-distance between  $x, y$  (See Def. 2).

*Proof.* Using Sanov's theorem<sup>28</sup> on large deviations, we conclude that the probability of spontaneous jump from strain  $x \in P$  to strain  $y \in Q$ , with the possibility  $P \neq Q$ , is given by:

$$Pr(x \rightarrow y) = \prod_{i=1}^N (\Phi_i^P(x_{-i})|_{y_i}) \quad (18)$$

Writing the factors on the right hand side as:

$$\Phi_i^P(x_{-i})|_{y_i} = \Phi_i^Q(y_{-i})|_{y_i} \left( \frac{\Phi_i^P(x_{-i})|_{y_i}}{\Phi_i^Q(y_{-i})|_{y_i}} \right) \quad (19)$$

we note that  $\Phi_i^P(x_{-i}), \Phi_i^Q(y_{-i})$  are distributions on the same index  $i$ , and hence:

$$|\Phi_i^P(x_{-i})|_{y_i} - \Phi_i^Q(y_{-i})|_{y_i}| \leq \sum_{y_i \in \Sigma_i} |\Phi_i^P(x_{-i})|_{y_i} - \Phi_i^Q(y_{-i})|_{y_i}| \quad (20)$$

Using a standard refinement of Pinsker's inequality<sup>46</sup>, and the relationship of Jensen-Shannon divergence with total variation, we get:

$$\theta_i \geq \frac{1}{8} |\Phi_i^P(x_{-i})|_{y_i} - \Phi_i^Q(y_{-i})|_{y_i}|^2 \Rightarrow \left| 1 - \frac{\Phi_i^Q(y_{-i})|_{y_i}}{\Phi_i^P(x_{-i})|_{y_i}} \right| \leq \frac{1}{a_0} \sqrt{8\theta_i} \quad (21)$$

where  $a_0$  is the smallest non-zero probability value of generating the entry at any index. We will see that this parameter is related to statistical significance of our bounds. First, we can formulate a lower bound as follows:

$$\log \left( \prod_{i=1}^N \frac{\Phi_i^P(x_{-i})|_{y_i}}{\Phi_i^Q(y_{-i})|_{y_i}} \right) = \sum_i \log \left( \frac{\Phi_i^P(x_{-i})|_{y_i}}{\Phi_i^Q(y_{-i})|_{y_i}} \right) \geq \sum_i \left( 1 - \frac{\Phi_i^Q(y_{-i})|_{y_i}}{\Phi_i^P(x_{-i})|_{y_i}} \right) \geq \frac{\sqrt{8}}{a_0} \sum_i \theta_i^{1/2} = -\frac{\sqrt{8}N}{a_0} \theta \quad (22)$$

Similarly, the upper bound may be derived as:

$$\log \left( \prod_{i=1}^N \frac{\Phi_i^P(x_{-i})|_{y_i}}{\Phi_i^Q(y_{-i})|_{y_i}} \right) = \sum_i \log \left( \frac{\Phi_i^P(x_{-i})|_{y_i}}{\Phi_i^Q(y_{-i})|_{y_i}} \right) \leq \sum_i \left( \frac{\Phi_i^Q(y_{-i})|_{y_i}}{\Phi_i^P(x_{-i})|_{y_i}} - 1 \right) \leq \frac{\sqrt{8}N}{a_0} \theta \quad (23)$$

Combining Eqs. 22 and 23, we conclude:

$$\omega_y^Q e^{\frac{\sqrt{8}N}{a_0} \theta} \geq Pr(x \rightarrow y) \geq \omega_y^Q e^{-\frac{\sqrt{8}N}{a_0} \theta} \quad (24)$$

Now, interpreting  $a_0$  as the probability of generating an unlikely event below our desired threshold (*i.e.* a "failure"), we note that the probability of generating at least one such event is given by  $1 - (1 - a_0)^N$ . Hence if  $\alpha$  is the pre-specified significance level, we have for  $N \gg 1$ :

$$a_0 \approx (1 - \alpha)/N \quad (25)$$

Hence, we conclude, that at significance level  $\geq \alpha$ , we have the bounds:

$$\omega_y^Q e^{\frac{\sqrt{8}N^2}{1-\alpha} \theta} \geq Pr(x \rightarrow y) \geq \omega_y^Q e^{-\frac{\sqrt{8}N^2}{1-\alpha} \theta} \quad (26)$$

□

**Remark 1.** This bound can be rewritten in terms of the log-likelihood of the spontaneous jump and constants independent of the initial sequence  $x$  as:

$$|\log Pr(x \rightarrow y) - C_0| \leq C_1 \theta \quad (27)$$

where the constants are given by:

$$C_0 = \log \omega_y^Q \quad (28)$$

$$C_1 = \frac{\sqrt{8}N^2}{1 - \alpha} \quad (29)$$

## DATA SHARING

Working software is publicly available at <https://pypi.org/project/emergenet/>. Accession numbers of all sequences used, and acknowledgement documentation for GISAID sequences is available as supplementary information.

## Data Source

In this study, we use sequences for the Hemagglutinin (HA) and Neuraminidase (NA) for Influenza A (for subtypes H1N1 and H3N2), which are key enablers of cellular entry and exit mechanisms respectively<sup>47</sup>. We use two sequences databases: 1) National Center for Biotechnology Information (NCBI) virus<sup>48</sup> and 2) GISAID<sup>49</sup> databases. The former is a community portal for viral sequence data, aiming to increase the usability of data archived in various NCBI repositories. GISAID has a somewhat more restricted user agreement, and use of GISAID data in an analysis requires acknowledgment of the contributions of both the submitting and the originating laboratories (Corresponding acknowledgment tables are included as supplementary information). We collected a total of 98,299 sequences in our analysis, although not all were used due to some being duplicates (see SI-Table ??).

## REFERENCES

- [1] Shao, W., Li, X., Goraya, M. U., Wang, S. & Chen, J.-L. Evolution of influenza a virus by mutation and reassortment. *International journal of molecular sciences* **18**, 1650 (2017).
- [2] Mills, C. E., Robins, J. M. & Lipsitch, M. Transmissibility of 1918 pandemic influenza. *Nature* **432**, 904–906 (2004).
- [3] Reid, A. H. & Taubenberger, J. K. The origin of the 1918 pandemic influenza virus: a continuing enigma. *Journal of general virology* **84**, 2285–2292 (2003).
- [4] Landolt, G. A. & Olsen, C. W. Up to new tricks—a review of cross-species transmission of influenza a viruses. *Animal Health Research Reviews* **8**, 1–21 (2007).
- [5] Dos Santos, G., Neumeier, E. & Bekkati-Berkani, R. Influenza: Can we cope better with the unpredictable? *Human vaccines & immunotherapeutics* **12**, 699–708 (2016).
- [6] Huddleston, J. *et al.* Integrating genotypes and phenotypes improves long-term forecasts of seasonal influenza a/h3n2 evolution. *Elife* **9**, e60067 (2020).

- 
- [7] Boni, M. F. Vaccination and antigenic drift in influenza. *Vaccine* **26**, C8–C14 (2008).
  - [8] Tricco, A. C. *et al.* Comparing influenza vaccine efficacy against mismatched and matched strains: a systematic review and meta-analysis. *BMC medicine* **11**, 153 (2013).
  - [9] Neher, R. A., Russell, C. A. & Shraiman, B. I. Predicting evolution from the shape of genealogical trees. *Elife* **3**, e03568 (2014).
  - [10] Vergara-Alert, J. *et al.* The ns segment of h5n1 avian influenza viruses (aiv) enhances the virulence of an h7n1 aiv in chickens. *Veterinary research* **45**, 1–11 (2014).
  - [11] Hothorn, T., Hornik, K. & Zeileis, A. Unbiased recursive partitioning: A conditional inference framework. *JOURNAL OF COMPUTATIONAL AND GRAPHICAL STATISTICS* **15**, 651–674 (2006).
  - [12] Wille, M., Geoghegan, J. L. & Holmes, E. C. How accurately can we assess zoonotic risk? *PLoS biology* **19**, e3001135 (2021).
  - [13] Pulliam, J. R. & Dushoff, J. Ability to replicate in the cytoplasm predicts zoonotic transmission of livestock viruses. *The Journal of infectious diseases* **199**, 565–568 (2009).
  - [14] Grawelle, R. E. Larger viral genome size facilitates emergence of zoonotic diseases. *bioRxiv* (2020).
  - [15] Grange, Z. L. *et al.* Ranking the risk of animal-to-human spillover for newly discovered viruses. *Proceedings of the National Academy of Sciences* **118**, e2002324118 (2021).
  - [16] CDC. Influenza risk assessment tool (irat) — pandemic influenza (flu) — cdc. <https://www.cdc.gov/flu/pandemic-resources/national-strategy/risk-assessment.htm>. (Accessed on 07/02/2021).
  - [17] Gamblin, S. J. & Skehel, J. J. Influenza hemagglutinin and neuraminidase membrane glycoproteins. *Journal of Biological Chemistry* **285**, 28403–28409 (2010).
  - [18] Posada, D. & Crandall, K. A. Modeltest: testing the model of dna substitution. *Bioinformatics (Oxford, England)* **14**, 817–818 (1998).
  - [19] Eng, C. L., Tong, J. C. & Tan, T. W. Predicting host tropism of influenza a virus proteins using random forest. *BMC medical genomics* **7**, 1–11 (2014).
  - [20] Ahlquist, P. Rna-dependent rna polymerases, viruses, and rna silencing. *Science* **296**, 1270–1273 (2002).
  - [21] Chen, R. & Holmes, E. C. Avian influenza virus exhibits rapid evolutionary dynamics. *Molecular biology and evolution* **23**, 2336–2341 (2006).
  - [22] Woolthuis, R. G., van Dorp, C. H., Keşmir, C., de Boer, R. J. & van Boven, M. Long-term adaptation of the influenza a virus by escaping cytotoxic t-cell recognition. *Scientific reports* **6**, 1–8 (2016).
  - [23] Fan, K. *et al.* Role of itk signalling in the interaction between influenza a virus and t-cells. *Journal of general virology* **93**, 987–997 (2012).
  - [24] van de Sandt, C. E. *et al.* Differential recognition of influenza a viruses by m158–66 epitope-specific cd8+ t cells is determined by extraepitopic amino acid residues. *Journal of virology* **90**, 1009–1022 (2016).
  - [25] Berkhoff, E., Geelhoed-Mieras, M., Fouchier, R., Osterhaus, A. & Rimmelzwaan, G. Assessment of the extent of variation in influenza a virus cytotoxic t-lymphocyte epitopes by using virus-specific cd8+ t-cell clones. *Journal of General Virology* **88**, 530–535 (2007).
  - [26] Van de Sandt, C. E., Kreijtz, J. H. & Rimmelzwaan, G. F. Evasion of influenza a viruses from innate and adaptive immune responses. *Viruses* **4**, 1438–1476 (2012).
  - [27] Wood, J. M. *et al.* Reproducibility of serology assays for pandemic influenza h1n1: collaborative study to evaluate a candidate who international standard. *Vaccine* **30**, 210–217 (2012).
  - [28] Cover, T. M. & Thomas, J. A. *Elements of Information Theory (Wiley Series in Telecommunications and Signal Processing)* (Wiley-Interscience, New York, NY, USA, 2006).
  - [29] Varadhan, S. S. Large deviations. In *Proceedings of the International Congress of Mathematicians 2010 (ICM 2010) (In 4 Volumes) Vol. I: Plenary Lectures and Ceremonies Vols. II–IV: Invited Lectures*, 622–639 (World Scientific, 2010).
  - [30] Agor, J. K. & Özaltın, O. Y. Models for predicting the evolution of influenza to inform vaccine strain selection. *Human vaccines & immunotherapeutics* **14**, 678–683 (2018).
  - [31] Cobey, S. *et al.* Poor immunogenicity, not vaccine strain egg adaptation, may explain the low h3n2 influenza vaccine effectiveness in 2012–2013. *Clinical Infectious Diseases* **67**, 327–333 (2018).
  - [32] Gouma, S., Weirick, M. & Hensley, S. E. Antigenic assessment of the h3n2 component of the 2019-2020 northern hemisphere influenza vaccine. *Nature communications* **11**, 1–5 (2020).
  - [33] Ma, W., Kahn, R. E. & Richt, J. A. The pig as a mixing vessel for influenza viruses: human and veterinary implications. *Journal of molecular and genetic medicine: an international journal of biomedical research* **3**, 158 (2009).
  - [34] Baumann, J., Kouassi, N. M., Foni, E., Klenk, H.-D. & Matrosovich, M. H1N1 Swine Influenza Viruses Differ from Avian Precursors by a Higher pH Optimum of Membrane Fusion .
  - [35] Qi, X. *et al.* Probable person to person transmission of novel avian influenza a (h7n9) virus in eastern china, 2013: epidemiological investigation. *Bmj* **347** (2013).
  - [36] Sun, H. *et al.* Mink is a highly susceptible host species to circulating human and avian influenza viruses. *Emerging microbes & infections* **10**, 472–480 (2021).
  - [37] Goldberger, A. L. & Peng, C.-K. Genomic classification using an information-based similarity index: application to the sars coronavirus. *Journal of Computational Biology* **12**, 1103–1116 (2005).
  - [38] Hulsenbeck, J. P. & Crandall, K. A. Phylogeny estimation and hypothesis testing using maximum likelihood. *Annual Review of Ecology and systematics* **28**, 437–466 (1997).

- 
- [39] van der Meer, F. J. U. M., Orsel, K. & Barkema, H. W. The new influenza A H1N1 virus: balancing on the interface of humans and animals. *The Canadian veterinary journal = La revue veterinaire canadienne* **51**, 56–62 (2010).
  - [40] Smith, G. J. D. *et al.* Origins and evolutionary genomics of the 2009 swine-origin H1N1 influenza A epidemic. *Nature* **459**, 1122–1125 (2009).
  - [41] Mollentze, N., Babayan, S. A. & Streicker, D. G. Identifying and prioritizing potential human-infecting viruses from their genome sequences. *PLoS biology* **19**, e3001390 (2021).
  - [42] Maher, M. C. *et al.* Predicting the mutational drivers of future sars-cov-2 variants of concern. *Science Translational Medicine* **14**, eabk3445 (2022).
  - [43] Hernández-Orozco, S., Kiani, N. A. & Zenil, H. Algorithmically probable mutations reproduce aspects of evolution, such as convergence rate, genetic memory and modularity. *Royal Society open science* **5**, 180399 (2018).
  - [44] Vanmarcke, E. *Random fields: analysis and synthesis* (World scientific, 2010).
  - [45] Manning, C. D., Manning, C. D. & Schütze, H. *Foundations of statistical natural language processing* (MIT press, 1999).
  - [46] Fedotov, A. A., Harremoës, P. & Topsøe, F. Refinements of pinsker's inequality. *IEEE Transactions on Information Theory* **49**, 1491–1498 (2003).
  - [47] McAuley, J., Gilbertson, B., Trifkovic, S., Brown, L. E. & McKimm-Breschkin, J. Influenza virus neuraminidase structure and functions. *Frontiers in microbiology* **10**, 39 (2019).
  - [48] Hatcher, E. L. *et al.* Virus variation resource—improved response to emergent viral outbreaks. *Nucleic acids research* **45**, D482–D490 (2017).
  - [49] Bogner, P., Capua, I., Lipman, D. J. & Cox, N. J. A global initiative on sharing avian flu data. *Nature* **442**, 981–981 (2006).

Extended Data Table 1  
Out-performance of Qnet recommendations over WHO for Influenza A vaccine composition

Subtype	Gene	Hemisphere	Two decades			One decade			2015-2019		
			WHO Error	Qnet Error	% Improvement	WHO Error	Qnet Error	% Improvement	WHO Error	Qnet Error	% Improvement
H1N1	HA	North	12.67	8.76	30.83	4.38	1.19	72.83	2.52	0.33	86.79
H1N1	HA	South	13.57	9.00	33.68	4.67	1.62	65.31	2.52	0.62	75.47
H1N1	HA	Average	13.12	8.88	32.25	4.53	1.40	69.07	2.52	0.48	81.13
H3N2	HA	North	7.65	4.71	38.46	5.00	2.94	41.18	1.82	0.88	51.61
H3N2	HA	South	7.59	4.82	36.43	4.94	3.00	39.29	1.82	0.94	48.39
H3N2	HA	Average	7.62	4.77	37.44	4.97	2.97	40.24	1.82	0.91	50.00
H1N1	NA	North	8.29	6.90	16.67	2.62	1.10	58.18	2.10	0.48	77.27
H1N1	NA	South	9.14	8.38	8.33	3.00	1.43	52.38	2.10	0.76	63.64
H1N1	NA	Average	8.72	7.64	12.50	2.81	1.27	55.28	2.10	0.62	70.46
H3N2	NA	North	4.21	3.63	13.75	2.11	1.79	15.00	1.32	0.32	76.00
H3N2	NA	South	4.68	4.16	11.24	2.58	2.05	20.41	1.32	0.42	68.00
H3N2	NA	Average	4.44	3.90	12.50	2.34	1.92	17.70	1.32	0.37	72.00

Extended Data Table 2

Examples: Qnet induced distance varying for fixed sequence pair when background population changes (rows 1 -5), sequences with small edit distance and large q-distance, and the converse (rows 6-9)

	Edit dist.	Sequence A	Sequence B	Q-dist.	Year A*	Year B*
1	18	A/Singapore/23J/2007	A/Tennessee/UR06-0294/2007	0.0111	2007	2007
2	18	A/Singapore/23J/2007	A/Tennessee/UR06-0294/2007	0.0094	2008	2008
3	18	A/Singapore/23J/2007	A/Tennessee/UR06-0294/2007	0.0027	2009	2009
4	18	A/Singapore/23J/2007	A/Tennessee/UR06-0294/2007	0.0025	2010	2010
5	18	A/Singapore/23J/2007	A/Tennessee/UR06-0294/2007	0.6163	2007	2010
6	11	A/Naypyitaw/M783/2008	A/Singapore/201/2008	0.8852	2008	2008
7	15	A/Cambodia/W0908339/2012	A/Singapore/DMS1233/2012	0.2737	2012	2012
8	126	A/South Dakota/03/2008	A/Singapore/10/2008	0.3034	2008	2008
9	141	A/Jodhpur/3248/2012	A/Cambodia/W0908339/2012	0.2405	2012	2012

\*Year A and year B correspond to the assumed collection years for sequences A and B respectively for the purpose of this example. Sequence A in row 1 is collected in 2007, but is assumed to be from different years in rows 2-4 to demonstrate the change in q-distance from sequence B, arising only from a change in the background population.

Extended Data Table 4  
Number of sequences collected from public databases

Extended Data Table 3  
Correlation between q-distance and edit distance between sequence pairs

Phenotypes	Correlation
Influenza H1N1 HA	0.76
Influenza H1N1 NA	0.74
Influenza H3N2 HA	0.85
Influenza H3N2 NA	0.79

Database	Strain	No. of Sequences
NCBI	Influenza H1N1 HA	17,894
NCBI	Influenza H1N1 NA	16,637
NCBI	Influenza H3N2 HA	18,265
NCBI	Influenza H3N2 NA	14,699
GISAID	Influenza H1N1 HA	1,528
GISAID	Influenza H1N1 NA	1,490
GISAID	Influenza H3N2 HA	13,975
GISAID	Influenza H3N2 NA	13,811
Total		98,299

Extended Data Table 5  
H1N1 HA Northern Hemisphere

Year	WHO Recommendation	Dominant Strain	Qnet Recommendation	WHO Error	Qnet Error
2001-02	A/New Caledonia/20/99	A/Canterbury/41/2001	A/Dunedin/2/2000	4	6
2002-03	A/New Caledonia/20/99	A/Taiwan/567/2002	A/New York/241/2001	3	1
2003-04	A/New Caledonia/20/99	A/Memphis/5/2003	A/New York/291/2002	5	2
2004-05	A/New Caledonia/20/99	A/Thailand/Siriraj-Rama-TT/2004	A/New York/222/2003	7	4
2005-06	A/New Caledonia/20/99	A/Niedersachsen/217/2005	A/Canterbury/106/2004	8	10
2006-07	A/New Caledonia/20/99	A/India/34980/2006	A/Auckland/619/2005	6	1
2007-08	A/Solomon Islands/3/2006	A/Norway/1701/2007	A/New York/8/2006	8	11
2008-09	A/Brisbane/59/2007	A/Pennsylvania/02/2008	A/Kentucky/UR06-0476/2007	2	2
2009-10	A/Brisbane/59/2007	A/Singapore/ON1060/2009	A/Hong Kong/549/2008	119	119
2010-11	A/California/7/2009	A/England/01220740/2010	A/New York/14/2009	5	1
2011-12	A/California/7/2009	A/Punjab/041/2011	A/Kansas/01/2010	7	2
2012-13	A/California/7/2009	A/British Columbia/001/2012	A/Moscow/WRAIR4308T/2011	11	4
2013-14	A/California/7/2009	A/Moscow/CRIE-32/2013	A/Helsinki/1199/2012	10	2
2014-15	A/California/7/2009	A/Thailand/CU-C5169/2014	A/Maryland/02/2013	12	0
2015-16	A/California/7/2009	A/Georgia/15/2015	A/Utah/3691/2014	14	2
2016-17	A/California/7/2009	A/Hawaii/21/2016	A/Adana/08/2015	16	0
2017-18	A/Michigan/45/2015	A/Michigan/291/2017	A/Beijing-Huairou/SWL1335/2016	5	4
2018-19	A/Michigan/45/2015	A/Washington/55/2018	A/India/C1721549/2017	6	1
2019-20	A/Brisbane/02/2018	A/Kentucky/06/2019	A/New Jersey/01/2018	5	1
2020-21	A/Hawaii/70/2019	A/Togo/905/2020	A/Italy/8949/2019	4	8
2021-22	A/Victoria/2570/2019	A/Ireland/20935/2022	A/Togo/45/2021	9	3
2022-23	-1	-1	A/Netherlands/00068/2022	-1	-1

\* Dominant strain is calculated as the one closest to the centroid in the strain space that year in the edit distance metric

Extended Data Table 6  
H1N1 HA Southern Hemisphere

Year	WHO Recommendation	Dominant Strain	Qnet Recommendation	WHO Error	Qnet Error
2001-02	A/New Caledonia/20/99	A/Canterbury/41/2001	A/South Canterbury/50/2000	4	6
2002-03	A/New Caledonia/20/99	A/Taiwan/567/2002	A/Canterbury/41/2001	3	1
2003-04	A/New Caledonia/20/99	A/Memphis/5/2003	A/New York/291/2002	5	2
2004-05	A/New Caledonia/20/99	A/Thailand/Siriraj-Rama-TT/2004	A/Memphis/5/2003	7	4
2005-06	A/New Caledonia/20/99	A/Niedersachsen/217/2005	A/Canterbury/106/2004	8	10
2006-07	A/New Caledonia/20/99	A/India/34980/2006	A/Niedersachsen/217/2005	6	2
2007-08	A/New Caledonia/20/99	A/Norway/1701/2007	A/Thailand/CU68/2006	14	6
2008-09	A/Solomon Islands/3/2006	A/Pennsylvania/02/2008	A/Kentucky/UR06-0476/2007	9	2
2009-10	A/Brisbane/59/2007	A/Singapore/ON1060/2009	A/Belem/241/2008	119	119
2010-11	A/California/7/2009	A/England/01220740/2010	A/Singapore/ON1060/2009	5	1
2011-12	A/California/7/2009	A/Punjab/041/2011	A/England/01220740/2010	7	2
2012-13	A/California/7/2009	A/British Columbia/001/2012	A/Punjab/041/2011	11	4
2013-14	A/California/7/2009	A/Moscow/CRIE-32/2013	A/India/P122045/2012	10	5
2014-15	A/California/7/2009	A/Thailand/CU-C5169/2014	A/Jiangsuhailing/SWL1382/2013	12	4
2015-16	A/California/7/2009	A/Georgia/15/2015	A/Thailand/CU-C5169/2014	14	2
2016-17	A/California/7/2009	A/Hawaii/21/2016	A/Georgia/15/2015	16	2
2017-18	A/Michigan/45/2015	A/Michigan/291/2017	A/Beijing-Huairou/SWL1335/2016	5	4
2018-19	A/Michigan/45/2015	A/Washington/55/2018	A/Michigan/291/2017	6	1
2019-20	A/Michigan/45/2015	A/Kentucky/06/2019	A/Washington/55/2018	7	1
2020-21	A/Brisbane/02/2018	A/Togo/905/2020	A/Italy/8451/2019	10	8
2021-22	A/Victoria/2570/2019	A/Abidjan/457/2021	A/Togo/0298/2021	9	5
2022-23	-1	-1	A/Cote_D'Ivoire/1270/2021	-1	-1

\* Dominant strain is calculated as the one closest to the centroid in the strain space that year in the edit distance metric

Extended Data Table 7  
H3N2 HA Northern Hemisphere



Year	WHO Recommendation	Dominant Strain	Qnet Recommendation	WHO Error	Qnet Error
2005-06	A/California/7/2004	A/Denmark/195/2005	A/Tairawhiti/369/2004	10	2
2006-07	A/Wisconsin/67/2005	A/New York/5/2006	A/South Australia/22/2005	5	4
2007-08	A/Wisconsin/67/2005	A/Tennessee/11/2007	A/Colorado/05/2006	8	5
2008-09	A/Brisbane/10/2007	A/Massachusetts/13/2008	A/Virginia/UR06-0021/2007	3	2
2009-10	A/Brisbane/10/2007	A/Hawaii/14/2009	A/Manhean/03/2008	7	6
2010-11	A/Perth/16/2009	A/Utah/12/2010	A/Philippines/5/2009	8	7
2011-12	A/Perth/16/2009	A/Piaui/14202/2011	A/Singapore/C2010.310/2010	4	4
2012-13	A/Victoria/361/2011	A/Alborz/927/2012	A/Tehran/895/2012	4	3
2013-14	A/Victoria/361/2011	A/Delaware/01/2013	A/Singapore/H2012.934/2012	4	1
2014-15	A/Texas/50/2012	A/Alborz/72205/2014	A/Nebraska/03/2013	10	9
2015-16	A/Switzerland/9715293/2013	A/Parma/471/2015	A/Ontario/01/2014	10	0
2016-17	A/Hong Kong/4801/2014	A/Guangdong/12/2016	A/Oregon/02/2015	0	0
2017-18	A/Hong Kong/4801/2014	A/Maryland/25/2017	A/New York/03/2016	3	1
2018-19	A/Singapore/INFIMH-16-0019/2016	A/Vermont/04/2018	A/Ontario/038/2017	8	5
2019-20	A/Kansas/14/2017	A/Kentucky/27/2019	A/California/7330/2018	16	12
2020-21	A/Hong Kong/2671/2019	A/India/Pun-NIV289524/2021_Jan	A/California/NHRC-OID_FDX100215/2019	16	14
2021-22	A/Cambodia/e0826360/2020	A/Human/New_York/PV60641/2022	A/India/Pun-NIV291000/2021_Jan	14	5
2022-23	-1	-1	A/Ireland/14993/2022	-1	-1

\* Dominant strain is calculated as the one closest to the centroid in the strain space that year in the edit distance metric

Extended Data Table 8  
H3N2 HA Southern Hemisphere

Year	WHO Recommendation	Dominant Strain	Qnet Recommendation	WHO Error	Qnet Error
2005-06	A/Wellington/1/2004	A/Denmark/195/2005	A/Waikato/21/2004	3	3
2006-07	A/California/7/2004	A/New York/5/2006	A/South Australia/22/2005	12	4
2007-08	A/Wisconsin/67/2005	A/Tennessee/11/2007	A/New York/923/2006	8	5
2008-09	A/Brisbane/10/2007	A/Massachusetts/13/2008	A/Tennessee/11/2007	3	2
2009-10	A/Brisbane/10/2007	A/Hawaii/14/2009	A/Manhean/03/2008	7	6
2010-11	A/Perth/16/2009	A/Utah/12/2010	A/Hawaii/14/2009	8	7
2011-12	A/Perth/16/2009	A/Piaui/14202/2011	A/Utah/12/2010	4	4
2012-13	A/Perth/16/2009	A/Alborz/927/2012	A/Piaui/14202/2011	8	4
2013-14	A/Victoria/361/2011	A/Delaware/01/2013	A/Callao/IPE00830/2012	4	7
2014-15	A/Texas/50/2012	A/Alborz/72205/2014	A/Delaware/01/2013	10	7
2015-16	A/Switzerland/9715293/2013	A/Parma/471/2015	A/Alborz/72205/2014	10	0
2016-17	A/Hong Kong/4801/2014	A/Guangdong/12/2016	A/Parma/471/2015	0	0
2017-18	A/Hong Kong/4801/2014	A/Maryland/25/2017	A/Ontario/196/2016	3	4
2018-19	A/Singapore/INFIMH-16-0019/2016	A/Vermont/04/2018	A/Texas/279/2017	8	5
2019-20	A/Switzerland/8060/2017	A/Kentucky/27/2019	A/Santa Catarina/1200/2018	13	12
2020-21	A/South Australia/34/2019	A/India/Pun-NIV289524/2021_Jan	A/Kentucky/27/2019	12	14
2021-22	A/Hong Kong/2671/2019	A/Darwin/9a/2021	A/India/PUN-NIV301718/2021	19	1
2022-23	-1	-1	A/Latvia/04-86261/2022	-1	-1

\* Dominant strain is calculated as the one closest to the centroid in the strain space that year in the edit distance metric



**Extended Data Figure 1. Sequence comparisons.** The observed dominant strain, we note that the correct Qnet deviations tend to be within the RBD, both for H1N1 and H3N2 for HA (panel a shows one example). Additionally, by comparing the type, side chain area, and the accessible side chain area, we note that the changes often have very different properties (panel b-f). Panels g-i show the localization of the deviations in the molecular structure of HA, where we note that the changes are most frequent in the HA1 sub-unit (the globular head), and around residues and structures that have been commonly implicated in receptor binding interactions e.g the ≈ 200 loop, the ≈ 220 loop and the ≈ 180-helix.

**Extended Data Table 9**  
**Riskiest Strains Currently Circulating in Swine**

H1N1 Strain	HA Risk	NA Risk	Overall Risk	Converted IRAT Score
A/swine/Tennessee/A02524414/2022	0.0201	0.0030	0.0077	6.2
A/swine/Missouri/A02750646/2022	0.0201	0.0070	0.0118	6.2
A/swine/Kansas/A02711847/2022	0.0201	0.0098	0.0141	6.2
A/swine/Iowa/A02636572/2022	0.0166	0.0225	0.0193	6.1
A/swine/Iowa/A02636308/2021	0.0143	0.0266	0.0195	6.1
A/swine/Illinois/A02750711/2022	0.0166	0.0233	0.0197	6.1
A/swine/Iowa/A02636616/2022	0.0166	0.0233	0.0197	6.1
A/swine/Oklahoma/A02246915/2022	0.0166	0.0233	0.0197	6.1
A/swine/Colorado/A02636469/2022	0.0166	0.0233	0.0197	6.1
A/swine/Iowa/A02636297/2021	0.0149	0.0267	0.0200	6.1
H3N2 Strain	HA Risk	NA Risk	Overall Risk	Converted IRAT Score
A/swine/Indiana/A02636492/2022	0.0104	0.0113	0.0108	6.2
A/swine/Indiana/A02636512/2022	0.0104	0.0113	0.0108	6.2
A/swine/Iowa/A02750695/2022	0.0110	0.0120	0.0115	6.2
A/swine/Oklahoma/A02711859/2022	0.0122	0.0114	0.0118	6.2
A/swine/Iowa/A02636351/2022	0.0121	0.0119	0.0120	6.2
A/swine/Iowa/A02636476/2022	0.0121	0.0120	0.0121	6.2
A/swine/Texas/A02636569/2022	0.0122	0.0120	0.0121	6.2
A/swine/Iowa/A02750726/2022	0.0123	0.0120	0.0121	6.2
A/swine/Iowa/A02750740/2022	0.0104	0.0156	0.0127	6.2
A/swine/Indiana/A02636521/2022	0.0104	0.0156	0.0127	6.2

\* Converted IRAT Score computed using regression generated from the IRAT vs. Qnet comparison

---

## SUPPLEMENTARY METHODS: NOTES ON Q-DISTANCE & SUPPORTING RESULTS

The *q-distance* is a pseudo-metric since distinct sequences can induce the same distributions over each index, and thus evaluate to have a zero distance. This is actually desirable; we do not want our distance to be sensitive to changes that are not biologically relevant. The intuition is that not all sequence variations brought about by substitutions are equally important or likely. Even with no selection pressure, we might still see random variations at an index if such variations do not affect the replicative fitness. Under that scenario, the corresponding  $\Phi_i$  will predict a flat distribution no matter what the input sequence is, thus contributing nothing to the overall distance. And even if two strains  $x, y$  have the same entry at some index  $i$ , the remaining residues might induce different distributions  $\Phi_i$  based on the remote dependencies, *i.e.*, the entries in  $x_{-i}, y_{-i}$ . Also, it matters if the sequences come from two different background populations  $P, Q$ , *i.e.*, if the induced Qnets  $\Phi^P, \Phi^Q$  are different. Thus, if we construct Qnets for H1N1 Influenza A separately for the collection years 2008 and 2009, then the same exact sequence collected in the respective years might have a non-zero distance between them, reflecting the fact that the background population the sequences arose from are different, inducing possibly different expected mutational tendencies (See SI-Table ??).

Next, we induce q-distance between a sequence and a population and between two populations.

**Definition 5** (Pseudo-metric between populations). *Using the notion of Hausdorff metric between sets:*

$$\forall x \in P, y \in Q, \theta(x, Q) = \min_{y \in Q} \theta(x, y) \quad (30)$$

$$\theta(P, Q) = \max \left\{ \max_{x \in P} \theta(x, Q), \max_{y \in Q} \theta(y, P) \right\} \quad (31)$$

### In-silico Corroboration of Qnet Constraints

We carry out in-silico experiments to corroborate that the constraints represented within an inferred Qnet are indeed reflective of the biology in play. We compare the results of simulated mutational perturbations to sequences from our databases (for which we have already constructed Qnets), and then use NCBI BLAST (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>) to identify if our perturbed sequences match with existing sequences in the databases (See SI-Fig. 1). We find that in contrast to random variations, which rapidly diverge the trajectories, the Qnet constraints tend to produce smaller variance in the trajectories, maintain a high degree of match as we extend our trajectories, and produces matches closer in time to the collection time of the initial sequence — suggesting that the Qnet does indeed capture realistic constraints.

### Multivariate Regression to Identify Factors in Strain Prediction

We investigate the key factors that contribute to our successful prediction of the dominant strain in the next season. We carry out a multivariate regression with data diversity, the complexity of inferred Qnet and the edit distance of the WHO recommendation from the dominant strain as independent variables. Here we define data diversity as the number of clusters we have in the input set of sequences, such that any two sequences five or less mutations apart are in the same cluster. Qnet complexity is measured by the number of decision nodes in the component decision trees of the recursive forest.

We select several plausible structures of the regression equation, and in each case conclude that data diversity has the most important and statistically significant contribution (See SI-Tab. 12).

## REFERENCES

- [1] Shao, W., Li, X., Goraya, M. U., Wang, S. & Chen, J.-L. Evolution of influenza a virus by mutation and reassortment. *International journal of molecular sciences* **18**, 1650 (2017).
- [2] Mills, C. E., Robins, J. M. & Lipsitch, M. Transmissibility of 1918 pandemic influenza. *Nature* **432**, 904–906 (2004).
- [3] Reid, A. H. & Taubenberger, J. K. The origin of the 1918 pandemic influenza virus: a continuing enigma. *Journal of general virology* **84**, 2285–2292 (2003).
- [4] Landolt, G. A. & Olsen, C. W. Up to new tricks—a review of cross-species transmission of influenza a viruses. *Animal Health Research Reviews* **8**, 1–21 (2007).
- [5] Dos Santos, G., Neumeier, E. & Bekkati-Berkani, R. Influenza: Can we cope better with the unpredictable? *Human vaccines & immunotherapeutics* **12**, 699–708 (2016).
- [6] Huddleston, J. *et al.* Integrating genotypes and phenotypes improves long-term forecasts of seasonal influenza a/h3n2 evolution. *Elife* **9**, e60067 (2020).
- [7] Boni, M. F. Vaccination and antigenic drift in influenza. *Vaccine* **26**, C8–C14 (2008).
- [8] Tricco, A. C. *et al.* Comparing influenza vaccine efficacy against mismatched and matched strains: a systematic review and meta-analysis. *BMC medicine* **11**, 153 (2013).
- [9] Neher, R. A., Russell, C. A. & Shraiman, B. I. Predicting evolution from the shape of genealogical trees. *Elife* **3**, e03568 (2014).


- 
- [10] Vergara-Alert, J. *et al.* The ns segment of h5n1 avian influenza viruses (aiv) enhances the virulence of an h7n1 aiv in chickens. *Veterinary research* **45**, 1–11 (2014).
- [11] Hothorn, T., Hornik, K. & Zeileis, A. Unbiased recursive partitioning: A conditional inference framework. *JOURNAL OF COMPUTATIONAL AND GRAPHICAL STATISTICS* **15**, 651–674 (2006).
- [12] Wille, M., Geoghegan, J. L. & Holmes, E. C. How accurately can we assess zoonotic risk? *PLoS biology* **19**, e3001135 (2021).
- [13] Pulliam, J. R. & Dushoff, J. Ability to replicate in the cytoplasm predicts zoonotic transmission of livestock viruses. *The Journal of infectious diseases* **199**, 565–568 (2009).
- [14] Grawelle, R. E. Larger viral genome size facilitates emergence of zoonotic diseases. *bioRxiv* (2020).
- [15] Grange, Z. L. *et al.* Ranking the risk of animal-to-human spillover for newly discovered viruses. *Proceedings of the National Academy of Sciences* **118**, e2002324118 (2021).
- [16] CDC. Influenza risk assessment tool (irat) — pandemic influenza (flu) — cdc. <https://www.cdc.gov/flu/pandemic-resources/national-strategy/risk-assessment.htm>. (Accessed on 07/02/2021).
- [17] Gamblin, S. J. & Skehel, J. J. Influenza hemagglutinin and neuraminidase membrane glycoproteins. *Journal of Biological Chemistry* **285**, 28403–28409 (2010).
- [18] Posada, D. & Crandall, K. A. Modeltest: testing the model of dna substitution. *Bioinformatics (Oxford, England)* **14**, 817–818 (1998).
- [19] Eng, C. L., Tong, J. C. & Tan, T. W. Predicting host tropism of influenza a virus proteins using random forest. *BMC medical genomics* **7**, 1–11 (2014).
- [20] Ahlquist, P. Rna-dependent rna polymerases, viruses, and rna silencing. *Science* **296**, 1270–1273 (2002).
- [21] Chen, R. & Holmes, E. C. Avian influenza virus exhibits rapid evolutionary dynamics. *Molecular biology and evolution* **23**, 2336–2341 (2006).
- [22] Woolthuis, R. G., van Dorp, C. H., Keşmir, C., de Boer, R. J. & van Boven, M. Long-term adaptation of the influenza a virus by escaping cytotoxic t-cell recognition. *Scientific reports* **6**, 1–8 (2016).
- [23] Fan, K. *et al.* Role of itk signalling in the interaction between influenza a virus and t-cells. *Journal of general virology* **93**, 987–997 (2012).
- [24] van de Sandt, C. E. *et al.* Differential recognition of influenza a viruses by m158–66 epitope-specific cd8+ t cells is determined by extraepitopic amino acid residues. *Journal of virology* **90**, 1009–1022 (2016).
- [25] Berkhoff, E., Geelhoed-Mieras, M., Fouchier, R., Osterhaus, A. & Rimmelzwaan, G. Assessment of the extent of variation in influenza a virus cytotoxic t-lymphocyte epitopes by using virus-specific cd8+ t-cell clones. *Journal of General Virology* **88**, 530–535 (2007).
- [26] Van de Sandt, C. E., Kreijtz, J. H. & Rimmelzwaan, G. F. Evasion of influenza a viruses from innate and adaptive immune responses. *Viruses* **4**, 1438–1476 (2012).
- [27] Wood, J. M. *et al.* Reproducibility of serology assays for pandemic influenza h1n1: collaborative study to evaluate a candidate who international standard. *Vaccine* **30**, 210–217 (2012).
- [28] Cover, T. M. & Thomas, J. A. *Elements of Information Theory (Wiley Series in Telecommunications and Signal Processing)* (Wiley-Interscience, New York, NY, USA, 2006).
- [29] Varadhan, S. S. Large deviations. In *Proceedings of the International Congress of Mathematicians 2010 (ICM 2010) (In 4 Volumes) Vol. I: Plenary Lectures and Ceremonies Vols. II–IV: Invited Lectures*, 622–639 (World Scientific, 2010).
- [30] Agor, J. K. & Özaltın, O. Y. Models for predicting the evolution of influenza to inform vaccine strain selection. *Human vaccines & immunotherapeutics* **14**, 678–683 (2018).
- [31] Cobey, S. *et al.* Poor immunogenicity, not vaccine strain egg adaptation, may explain the low h3n2 influenza vaccine effectiveness in 2012–2013. *Clinical Infectious Diseases* **67**, 327–333 (2018).
- [32] Gouma, S., Weirick, M. & Hensley, S. E. Antigenic assessment of the h3n2 component of the 2019-2020 northern hemisphere influenza vaccine. *Nature communications* **11**, 1–5 (2020).
- [33] Ma, W., Kahn, R. E. & Richt, J. A. The pig as a mixing vessel for influenza viruses: human and veterinary implications. *Journal of molecular and genetic medicine: an international journal of biomedical research* **3**, 158 (2009).
- [34] Baumann, J., Kouassi, N. M., Foni, E., Klenk, H.-D. & Matrosovich, M. H1N1 Swine Influenza Viruses Differ from Avian Precursors by a Higher pH Optimum of Membrane Fusion .
- [35] Qi, X. *et al.* Probable person to person transmission of novel avian influenza a (h7n9) virus in eastern china, 2013: epidemiological investigation. *Bmj* **347** (2013).
- [36] Sun, H. *et al.* Mink is a highly susceptible host species to circulating human and avian influenza viruses. *Emerging microbes & infections* **10**, 472–480 (2021).
- [37] Goldberger, A. L. & Peng, C.-K. Genomic classification using an information-based similarity index: application to the sars coronavirus. *Journal of Computational Biology* **12**, 1103–1116 (2005).
- [38] Huelsenbeck, J. P. & Crandall, K. A. Phylogeny estimation and hypothesis testing using maximum likelihood. *Annual Review of Ecology and systematics* **28**, 437–466 (1997).
- [39] van der Meer, F. J. U. M., Orsel, K. & Barkema, H. W. The new influenza A H1N1 virus: balancing on the interface of humans and animals. *The Canadian veterinary journal = La revue veterinaire canadienne* **51**, 56–62 (2010).
- [40] Smith, G. J. D. *et al.* Origins and evolutionary genomics of the 2009 swine-origin H1N1 influenza A epidemic. *Nature* **459**, 1122–1125 (2009).
- [41] Mollentze, N., Babayan, S. A. & Streicker, D. G. Identifying and prioritizing potential human-infecting viruses from

- 
- their genome sequences. *PLoS biology* **19**, e3001390 (2021).
- [42] Maher, M. C. *et al.* Predicting the mutational drivers of future sars-cov-2 variants of concern. *Science Translational Medicine* **14**, eabk3445 (2022).
- [43] Hernández-Orozco, S., Kiani, N. A. & Zenil, H. Algorithmically probable mutations reproduce aspects of evolution, such as convergence rate, genetic memory and modularity. *Royal Society open science* **5**, 180399 (2018).
- [44] Vanmarcke, E. *Random fields: analysis and synthesis* (World scientific, 2010).
- [45] Manning, C. D., Manning, C. D. & Schütze, H. *Foundations of statistical natural language processing* (MIT press, 1999).
- [46] Fedotov, A. A., Harremoës, P. & Topsøe, F. Refinements of pinsker's inequality. *IEEE Transactions on Information Theory* **49**, 1491–1498 (2003).
- [47] McAuley, J., Gilbertson, B., Trifkovic, S., Brown, L. E. & McKimm-Breschkin, J. Influenza virus neuraminidase structure and functions. *Frontiers in microbiology* **10**, 39 (2019).
- [48] Hatcher, E. L. *et al.* Virus variation resource—improved response to emergent viral outbreaks. *Nucleic acids research* **45**, D482–D490 (2017).
- [49] Bogner, P., Capua, I., Lipman, D. J. & Cox, N. J. A global initiative on sharing avian flu data. *Nature* **442**, 981–981 (2006).

SI Fig. 1. **Q-distance validation in-silico using Influenza A sequences from NCBI database.** Panel **a** illustrates that the Qnet induced modeling of evolutionary trajectories initiated from known haemagglutinin (HA) sequences are distinct from random paths in the strain space. In particular, random trajectories have more variance, and more importantly, diverge to different regions of the landscape compared to Qnet predictions. Panels **b-e** show that unconstrained Q-sampling produces sequences maintain a higher degree of similarity to known sequences, as verified by blasting against known HA sequences, have a smaller rate of growth of variance, and produce matches in closer time frames to the initial sequence. Panel **c** shows that this is not due to simply restricting the mutational variations, which increases rapidly in both the Qnet and the classical metric.

## SUPPLEMENTARY FIGURES & TABLES

**a. Distribution around dominant strain**



SI Fig. 2. **No. of mutations from the seasonal dominant strain over the years** The quasispecies that circulates each season for each sub-type is tightly distributed around the dominant strain on average.

SI Tab. 1  
H1N1 NA Northern Hemisphere

Year	WHO Recommendation	Dominant Strain	Qnet Recommendation	WHO Error	Qnet Error
2001-02	A/New Caledonia/20/99	A/New York/447/2001	A/Memphis/15/2000	4	4
2002-03	A/New Caledonia/20/99	A/Paris/0833/2002	A/New York/341/2001	1	5
2003-04	A/New Caledonia/20/99	A/Memphis/5/2003	A/New York/291/2002	3	5
2004-05	A/New Caledonia/20/99	A/Singapore/14/2004	A/New York/223/2003	2	3
2005-06	A/New Caledonia/20/99	A/Taiwan/5524/2005	A/Florida/3e/2004	3	0
2006-07	A/New Caledonia/20/99	A/Massachusetts/08/2006	A/Sofia/361/2005	4	2
2007-08	A/Solomon Islands/3/2006	A/Tennessee/UR06-0106/2007	A/Sofia/490/2006	9	2
2008-09	A/Brisbane/59/2007	A/Sendai/TU66/2008	A/Maryland/04/2007	0	3
2009-10	A/Brisbane/59/2007	A/Thailand/SR08021/2009	A/Paris/910/2008	87	87
2010-11	A/California/7/2009	A/Finland/2460N/2010	A/Rome/709/2009	2	9
2011-12	A/California/7/2009	A/Tula/CRIE-GSYu/2011	A/Oman/SQUH-40/2010	4	2
2012-13	A/California/7/2009	A/Bangalore/697-32/2012	A/Nizhnii Novgorod/CRIE-ZCA/2011	4	0
2013-14	A/California/7/2009	A/Jiangsugusu/SWL1824/2013	A/LongYan/SWL33/2013	5	3
2014-15	A/California/7/2009	A/LongYan/SWL2457/2014	A/Utah/06/2013	9	3
2015-16	A/California/7/2009	A/Michigan/45/2015	A/Maryland/02/2014	14	4
2016-17	A/California/7/2009	A/Mexico/4436/2016	A/India/Pun151245/2015	14	0
2017-18	A/Michigan/45/2015	A/Illinois/37/2017	A/Utah/02/2016	3	3
2018-19	A/Michigan/45/2015	A/Kenya/47/2018	A/Maine/24/2017	4	0
2019-20	A/Brisbane/02/2018	A/Texas/7939/2019	A/Missouri/03/2018	1	0
2020-21	A/Hawaii/70/2019	A/Togo/897/2020	A/Texas/112/2019	0	5
2021-22	A/Victoria/2570/2019	A/Cote_d'Ivoire/3729/2021	A/Togo/0071/2021	1	5
2022-23	-1	-1	A/Lyon/820/2021	-1	-1

\* Dominant strain is calculated as the one closest to the centroid in the strain space that year in the edit distance metric

SI Tab. 2  
H1N1 NA Southern Hemisphere

Year	WHO Recommendation	Dominant Strain	Qnet Recommendation	WHO Error	Qnet Error
2001-02	A/New Caledonia/20/99	A/New York/447/2001	A/Canterbury/37/2000	4	6
2002-03	A/New Caledonia/20/99	A/Paris/0833/2002	A/New York/447/2001	1	5
2003-04	A/New Caledonia/20/99	A/Memphis/5/2003	A/New York/291/2002	3	5
2004-05	A/New Caledonia/20/99	A/Singapore/14/2004	A/Memphis/5/2003	2	3
2005-06	A/New Caledonia/20/99	A/Taiwan/5524/2005	A/Canterbury/106/2004	3	6
2006-07	A/New Caledonia/20/99	A/Massachusetts/08/2006	A/Sofia/361/2005	4	2
2007-08	A/New Caledonia/20/99	A/Tennessee/UR06-0106/2007	A/Thailand/RMSC-UDN-20/2006	4	8
2008-09	A/Solomon Islands/3/2006	A/Sendai/TU66/2008	A/Tennessee/UR06-0151/2007	15	13
2009-10	A/Brisbane/59/2007	A/Thailand/SR08021/2009	A/Nebraska/07/2008	87	87
2010-11	A/California/7/2009	A/Finland/2460N/2010	A/Rome/709/2009	2	9
2011-12	A/California/7/2009	A/Tula/CRIE-GSYu/2011	A/Finland/2460N/2010	4	2
2012-13	A/California/7/2009	A/Bangalore/697-32/2012	A/Tula/CRIE-GSYu/2011	4	0
2013-14	A/California/7/2009	A/Jiangsugusu/SWL1824/2013	A/Oman/SQUH-63/2012	5	4
2014-15	A/California/7/2009	A/LongYan/SWL2457/2014	A/NanPing/SWL1640/2013	9	6
2015-16	A/California/7/2009	A/Michigan/45/2015	A/LongYan/SWL2457/2014	14	5
2016-17	A/California/7/2009	A/Mexico/4436/2016	A/Michigan/45/2015	14	0
2017-18	A/Michigan/45/2015	A/Illinois/37/2017	A/Mexico/4436/2016	3	3
2018-19	A/Michigan/45/2015	A/Kenya/47/2018	A/Kentucky/26/2017	4	2
2019-20	A/Michigan/45/2015	A/Texas/7939/2019	A/Kenya/47/2018	4	0
2020-21	A/Brisbane/02/2018	A/Togo/897/2020	A/Texas/7939/2019	6	5
2021-22	A/Victoria/2570/2019	A/Cote_D'Ivoire/1496/2021	A/NAGASAKI/8/2020	1	6
2022-23	-1	-1	A/Dakar/35/2021	-1	-1

\* Dominant strain is calculated as the one closest to the centroid in the strain space that year in the edit distance metric

SI Fig. 3. IRAT vs. Q-distance relationship for H1- and H3- sub-types, using past year data vs. using all data. On the result when computing average q-distance between the target strain and the circulating human strains from the past year, and on the right is the result when using all available human strains of that sub-type. Evidently, the former has a much higher correlation, since a strain being "close" to humans at some point does not necessarily mean being close now.

SI Tab. 3  
H3N2 NA Northern Hemisphere

Year	WHO Recommendation	Dominant Strain	Qnet Recommendation	WHO Error	Qnet Error
2003-04	A/Moscow/10/99	A/Denmark/107/2003	A/New York/100/2002	13	3
2004-05	A/Fujian/411/2002	A/Hyogo/36/2004	A/New York/20/2003	3	16
2005-06	A/California/7/2004	A/Denmark/203/2005	A/Hong Kong/HKU20/2004	4	0
2006-07	A/Wisconsin/67/2005	A/Berlin/32/2006	A/Mexico/InDRE2227/2005	1	1
2007-08	A/Wisconsin/67/2005	A/Brazil/80/2007	A/Baden-Wuerttemberg/17/2006	8	7
2008-09	A/Brisbane/10/2007	A/Missouri/05/2008	A/Washington/01/2007	3	2
2009-10	A/Brisbane/10/2007	A/Oklahoma/09/2009	A/Wisconsin/24/2008	3	1
2010-11	A/Perth/16/2009	A/California/17/2010	A/New York/70/2009	2	3
2011-12	A/Perth/16/2009	A/Texas/14/2011	A/California/14/2010	3	2
2012-13	A/Victoria/361/2011	A/New York/02/2012	A/Singapore/C2011.493/2011	4	1
2013-14	A/Victoria/361/2011	A/Michigan/02/2013	A/New York/01/2012	3	1
2014-15	A/Texas/50/2012	A/Tehran/69634/2014	A/Boston/DOA2-176/2013	3	1
2015-16	A/Switzerland/9715293/2013	A/Parma/471/2015	A/Thailand/CU-B10520/2014	3	0
2016-17	A/Hong Kong/4801/2014	A/North Carolina/62/2016	A/Delaware/02/2015	7	2
2017-18	A/Hong Kong/4801/2014	A/Texas/277/2017	A/New York/03/2016	8	0
2018-19	A/Singapore/INFIMH-16-0019/2016	A/Japan/NHRC_FDX70352/2018	A/Colorado/11/2017	4	3
2019-20	A/Kansas/14/2017	A/Washington/9757/2019	A/Guangxi-Fangcheng/54/2019	3	11
2020-21	A/Hong Kong/2671/2019	A/Bangladesh/1004005/2020	A/Maryland/02/2019	3	13
2021-22	A/Cambodia/e0826360/2020	A/Stockholm/10/2022	A/Bangladesh/1916/2020	2	2
2022-23	-1	-1	A/Iowa/20/2022	-1	-1

\* Dominant strain is calculated as the one closest to the centroid in the strain space that year in the edit distance metric

SI Tab. 4  
H3N2 NA Southern Hemisphere

Year	WHO Recommendation	Dominant Strain	Qnet Recommendation	WHO Error	Qnet Error
2003-04	A/Moscow/10/99	A/Denmark/107/2003	A/New York/101/2002	13	3
2004-05	A/Fujian/411/2002	A/Hyogo/36/2004	A/New York/20/2003	3	16
2005-06	A/Wellington/1/2004	A/Denmark/203/2005	A/Wellington/1/2004	2	2
2006-07	A/California/7/2004	A/Berlin/32/2006	A/Mexico/InDRE2227/2005	3	1
2007-08	A/Wisconsin/67/2005	A/Brazil/80/2007	A/Ohio/06/2006	8	10
2008-09	A/Brisbane/10/2007	A/Missouri/05/2008	A/Brazil/80/2007	3	2
2009-10	A/Brisbane/10/2007	A/Oklahoma/09/2009	A/Wisconsin/24/2008	3	1
2010-11	A/Perth/16/2009	A/California/17/2010	A/New York/70/2009	2	3
2011-12	A/Perth/16/2009	A/Texas/14/2011	A/Virginia/05/2010	3	2
2012-13	A/Perth/16/2009	A/New York/02/2012	A/Texas/14/2011	4	1
2013-14	A/Victoria/361/2011	A/Michigan/02/2013	A/New York/02/2012	3	3
2014-15	A/Texas/50/2012	A/Tehran/69634/2014	A/Michigan/02/2013	3	1
2015-16	A/Switzerland/9715293/2013	A/Parma/471/2015	A/Tehran/69634/2014	3	2
2016-17	A/Hong Kong/4801/2014	A/North Carolina/62/2016	A/Parma/471/2015	7	2
2017-18	A/Hong Kong/4801/2014	A/Texas/277/2017	A/Guangdong/264/2016	8	0
2018-19	A/Singapore/INFIMH-16-0019/2016	A/Japan/NHRC_FDX70352/2018	A/Texas/277/2017	4	3
2019-20	A/Switzerland/8060/2017	A/Washington/9757/2019	A/Pennsylvania/317/2018	10	10
2020-21	A/South Australia/34/2019	A/Bangladesh/1004005/2020	A/Washington/9757/2019	1	13
2021-22	A/Hong Kong/2671/2019	A/India/PUN-NIV301718/2021	A/India/PUN-NIV301132/2021	6	4
2022-23	-1	-1	A/Michigan/UOM10045036720/2022	-1	-1

\* Dominant strain is calculated as the one closest to the centroid in the strain space that year in the edit distance metric

SI Tab. 5  
H1N1 NA Northern Hemisphere (Multi-cluster)

Year	WHO Recommendation	WHO Error	Qnet Error 1	Qnet Error 2	Qnet Recommendation 1	Qnet Recommendation 2
2001-02	A/New Caledonia/20/99	4	1	6	A/New South Wales/26/2000	A/Canterbury/37/2000
2002-03	A/New Caledonia/20/99	1	0	5	A/Wellington/1/2001	A/New York/447/2001
2003-04	A/New Caledonia/20/99	3	2	8	A/Paris/0833/2002	A/Taiwan/141/2002
2004-05	A/New Caledonia/20/99	2	3	4	A/Memphis/5/2003	A/Hanoi/1004/2003
2005-06	A/New Caledonia/20/99	3	0	1	A/Denmark/130/2004	A/Paris/650/2004
2006-07	A/New Caledonia/20/99	4	2	8	A/Sofia/361/2005	A/Wellington/11/2005
2007-08	A/Solomon Islands/3/2006	9	4	8	A/Sofia/246/2006	A/New York/8/2006
2008-09	A/Brisbane/59/2007	0	13	19	A/Tennessee/UR06-0151/2007	A/Ohio/UR06-0178/2007
2009-10	A/Brisbane/59/2007	87	88	90	A/Sendai/TU66/2008	A/Japan/618/2008
2010-11	A/California/7/2009	2	1	6	A/South Carolina/WRAIR1645P/2009	A/Wisconsin/629-D00809/2009
2011-12	A/California/7/2009	4	1	3	A/England/21680633/2010	A/Hangzhou/178/2010
2012-13	A/California/7/2009	4	1	22	A/Joshkar-Ola/CRIE-BLP/2011	A/Rio Grande do Sul/578/2011
2013-14	A/California/7/2009	5	4	13	A/Thailand/MR10580/2012	A/Mexico/INMEGEN-INER 15/2012
2014-15	A/California/7/2009	9	3	7	A/Minnesota/02/2013	A/Helsinki/430/2013
2015-16	A/California/7/2009	14	4	7	A/Helsinki/808M/2014	A/Virginia/NHRC430739/2014
2016-17	A/California/7/2009	14	0	3	A/Michigan/45/2015	A/Colorado/30/2015
2017-18	A/Michigan/45/2015	3	3	8	A/Mexico/4436/2016	A/Arizona/03/2016
2018-19	A/Michigan/45/2015	4	0	4	A/California/NHRC_QV11073/2017	A/Minnesota/35/2017
2019-20	A/Brisbane/02/2018	1	0	2	A/Kenya/47/2018	A/Colorado/7682/2018
2020-21	A/Hawaii/70/2019	0	3	8	A/California/NHRC-OID_BOX-ILI-0012/2019	A/Indiana/30/2019
2021-22	A/Victoria/2570/2019	1	5	51	A/Togo/0071/2021	A/Yunnan-Mengzi/1462/2020
2022-23	-1	-1	-1	-1	A/Netherlands/10646/2022	A/Sydney/234/2022

\* Dominant strain is calculated as the one closest to the centroid in the strain space that year in the edit distance metric

SI Tab. 6  
H1N1 NA Southern Hemisphere (Multi-cluster)

Year	WHO Recommendation	WHO Error	Qnet Error 1	Qnet Error 2	Qnet Recommendation 1	Qnet Recommendation 2
2001-02	A/New Caledonia/20/99	4	1	6	A/New South Wales/26/2000	A/Canterbury/37/2000
2002-03	A/New Caledonia/20/99	1	0	5	A/Wellington/1/2001	A/New York/447/2001
2003-04	A/New Caledonia/20/99	3	2	8	A/Paris/0833/2002	A/Taiwan/141/2002
2004-05	A/New Caledonia/20/99	2	3	4	A/Memphis/5/2003	A/Hanoi/1004/2003
2005-06	A/New Caledonia/20/99	3	0	1	A/Denmark/130/2004	A/Paris/650/2004
2006-07	A/New Caledonia/20/99	4	2	8	A/Sofia/361/2005	A/Wellington/11/2005
2007-08	A/New Caledonia/20/99	4	4	8	A/Sofia/246/2006	A/New York/8/2006
2008-09	A/Solomon Islands/3/2006	15	13	19	A/Tennessee/UR06-0151/2007	A/Ohio/UR06-0178/2007
2009-10	A/Brisbane/59/2007	87	88	90	A/Sendai/TU66/2008	A/Japan/618/2008
2010-11	A/California/7/2009	2	1	6	A/South Carolina/WRAIR1645P/2009	A/Wisconsin/629-D00809/2009
2011-12	A/California/7/2009	4	1	3	A/England/21680633/2010	A/Hangzhou/178/2010
2012-13	A/California/7/2009	4	1	22	A/Joshkar-Ola/CRIE-BLP/2011	A/Rio Grande do Sul/578/2011
2013-14	A/California/7/2009	5	4	13	A/Thailand/MR10580/2012	A/Mexico/INMEGEN-INER 15/2012
2014-15	A/California/7/2009	9	3	7	A/Minnesota/02/2013	A/Helsinki/430/2013
2015-16	A/California/7/2009	14	4	7	A/Helsinki/808M/2014	A/Virginia/NHRC430739/2014
2016-17	A/California/7/2009	14	0	3	A/Michigan/45/2015	A/Colorado/30/2015
2017-18	A/Michigan/45/2015	3	3	8	A/Mexico/4436/2016	A/Arizona/03/2016
2018-19	A/Michigan/45/2015	4	0	4	A/California/NHRC_QV11073/2017	A/Minnesota/35/2017
2019-20	A/Michigan/45/2015	4	0	2	A/Kenya/47/2018	A/Colorado/7682/2018
2020-21	A/Brisbane/02/2018	5	2	7	A/California/NHRC-OID_BOX-ILI-0012/2019	A/Indiana/30/2019
2021-22	A/Victoria/2570/2019	1	7	58	A/Togo/0155/2021	A/Shandong/00204/2021
2022-23	-1	-1	-1	-1	A/Switzerland/86136/2022	A/Wisconsin/04/2021

\* Dominant strain is calculated as the one closest to the centroid in the strain space that year in the edit distance metric

SI Tab. 7  
H3N2 NA Northern Hemisphere (Multi-cluster)

Year	WHO Recommendation	WHO Error	Qnet Error 1	Qnet Error 2	Qnet Recommendation 1	Qnet Recommendation 2
2003-04	A/Moscow/10/99	13	4	5	A/Auckland/612/2002	A/New York/87/2002
2004-05	A/Fujian/411/2002	3	16	18	A/New York/20/2003	A/New York/12/2003
2005-06	A/California/7/2004	4	1	7	A/New York/358/2004	A/Singapore/36/2004
2006-07	A/Wisconsin/67/2005	1	3	8	A/Macau/557/2005	A/Hong Kong/HKU53/2005
2007-08	A/Wisconsin/67/2005	8	0	10	A/Wisconsin/42/2006	A/Wisconsin/44/2006
2008-09	A/Brisbane/10/2007	3	4	10	A/Missouri/06/2007	A/Japan/72/2007
2009-10	A/Brisbane/10/2007	3	1	7	A/Wisconsin/24/2008	A/Mississippi/UR07-0042/2008
2010-11	A/Perth/16/2009	2	3	8	A/New York/70/2009	A/Japan/883/2009
2011-12	A/Perth/16/2009	3	2	2	A/California/19/2010	A/Virginia/05/2010
2012-13	A/Victoria/361/2011	4	1	12	A/Texas/14/2011	A/Singapore/GP1684/2011
2013-14	A/Victoria/361/2011	3	1	5	A/Idaho/38/2012	A/Pavia/135/2012
2014-15	A/Texas/50/2012	3	1	1	A/Nevada/05/2013	A/Michigan/02/2013
2015-16	A/Switzerland/9715293/2013	3	0	4	A/Nicaragua/6866_14/2014	A/Iran/91244/2014
2016-17	A/Hong Kong/4801/2014	7	1	25	A/New Jersey/13/2015	A/California/NHRC_BRD41056N/2015
2017-18	A/Hong Kong/4801/2014	9	1	4	A/Guangdong/264/2016	A/Victoria/668/2016
2018-19	A/Singapore/INFIMH-16-0019/2016	3	2	4	A/Netherlands/3530/2017	A/Washington/17/2017
2019-20	A/Kansas/14/2017	3	4	10	A/England/538/2018	A/California/BRD12490N/2018
2020-21	A/Hong Kong/2671/2019	3	1	13	A/England/9738/2019	A/Washington/9757/2019
2021-22	A/Cambodia/e0826360/2020	2	3	7	A/Laos/527/2021	A/Michigan/UOM10045655748/2020
2022-23	-1	-1	-1	-1	A/Maine/02/2022	A/Michigan/UOM10042819294/2021

\* Dominant strain is calculated as the one closest to the centroid in the strain space that year in the edit distance metric

SI Tab. 8  
H3N2 NA Southern Hemisphere (Multi-cluster)

Year	WHO Recommendation	WHO Error	Qnet Error 1	Qnet Error 2	Qnet Recommendation 1	Qnet Recommendation 2
2003-04	A/Moscow/10/99	13	4	5	A/Auckland/612/2002	A/New York/87/2002
2004-05	A/Fujian/411/2002	3	16	18	A/New York/20/2003	A/New York/12/2003
2005-06	A/Wellington/1/2004	2	1	7	A/New York/358/2004	A/Singapore/36/2004
2006-07	A/California/7/2004	3	3	8	A/Macau/557/2005	A/Hong Kong/HKU53/2005
2007-08	A/Wisconsin/67/2005	8	0	10	A/Wisconsin/42/2006	A/Wisconsin/44/2006
2008-09	A/Brisbane/10/2007	3	4	10	A/Missouri/06/2007	A/Japan/72/2007
2009-10	A/Brisbane/10/2007	3	1	7	A/Wisconsin/24/2008	A/Mississippi/UR07-0042/2008
2010-11	A/Perth/16/2009	2	3	8	A/New York/70/2009	A/Japan/883/2009
2011-12	A/Perth/16/2009	3	2	2	A/California/19/2010	A/Virginia/05/2010
2012-13	A/Perth/16/2009	4	1	12	A/Texas/14/2011	A/Singapore/GP1684/2011
2013-14	A/Victoria/361/2011	3	1	5	A/Idaho/38/2012	A/Pavia/135/2012
2014-15	A/Texas/50/2012	3	1	1	A/Nevada/05/2013	A/Michigan/02/2013
2015-16	A/Switzerland/9715293/2013	3	0	4	A/Nicaragua/6866_14/2014	A/Iran/91244/2014
2016-17	A/Hong Kong/4801/2014	7	1	25	A/New Jersey/13/2015	A/California/NHRC_BRD41056N/2015
2017-18	A/Hong Kong/4801/2014	9	1	4	A/Guangdong/264/2016	A/Victoria/668/2016
2018-19	A/Singapore/INFIMH-16-0019/2016	3	2	4	A/Netherlands/3530/2017	A/Washington/17/2017
2019-20	A/Switzerland/8060/2017	10	4	10	A/England/538/2018	A/California/BRD12490N/2018
2020-21	A/South Australia/34/2019	1	1	13	A/England/9738/2019	A/Washington/9757/2019
2021-22	A/Hong Kong/2671/2019	6	1	49	A/Darwin/11/2021	A/Hawaii/28/2020
2022-23	-1	-1	-1	-1	A/Congo/313/2021	A/Texas/12723/2022

\* Dominant strain is calculated as the one closest to the centroid in the strain space that year in the edit distance metric

SI Tab. 9  
Influenza A Strains Evaluated by IRAT and Corresponding Qnet Computed Risk Scores

Influenza Virus	Subtype	IRAT Date	IRAT Emergence Score	IRAT Impact Score	HA Sample	NA Sample	HA Avg. E-distance	NA Avg. E-distance	Geom. Mean	Emergent Emergence Score	Emergent Impact Score
A/swine/Shandong/1207/2016	H1N1	Jul 2020	7.5	6.9	1000	1000	0.0941	0.0205	0.0440	6.0	6.2
A/Ohio/13/2017	H3N2	Jul 2019	6.6	5.8	1000	1000	0.0184	0.0306	0.0238	6.3	6.2
A/Hong Kong/125/2017	H7N9	May 2017	6.5	7.5	437	437	0.0296	0.0058	0.0131	6.6	6.5
A/Shanghai/02/2013	H7N9	Apr 2016	6.4	7.2	178	178	0.0055	0.0036	0.0044	6.7	6.6
A/Anhui-Lujiang/39/2018	H9N2	Jul 2019	6.2	5.9	31	30	0.0290	0.1681	0.0698	5.2	5.0
A/Indiana/08/2011	H3N2	Dec 2012	6.0	4.5	1000	1000	0.0523	0.0091	0.0218	6.4	6.5
A/California/62/2018	H1N2	Jul 2019	5.8	5.7	55	55	0.1089	0.0610	0.0815	5.4	5.5
A/Bangladesh/0994/2011***	H9N2	Feb 2014	5.6	5.4			0.2078	0.1823	0.1947	4.3	4.9
A/Sichuan/06681/2021	H5N6	Oct 2021	5.3	6.3	45	45	0.3616	0.0518	0.1369	5.2	6.4
A/Vietnam/1203/2004	H5N1	Nov 2011	5.2	6.6	258	246	0.1673	0.0111	0.0430	6.2	6.7
A/Yunnan/14564/2015**	H5N6	Apr 2016	5.0	6.6	344	331	0.3482	0.2987	0.3225	4.9	6.5
A/Astrakhan/3212/2020**	H5N8	Mar 2021	4.6	5.2	381	365	0.1603	0.3472	0.2359	3.9	4.4
A/Netherlands/219/2003	H7N7	Jun 2012	4.6	5.8	46	46	0.2757	0.3521	0.3115	4.6	5.8
A/American wigeon/South Carolina/AH0195145/2021	H5N1	Mar 2022	4.4	5.1	335	323	0.1722	0.5114	0.2967	4.0	4.7
A/Jiangxi-Donghu/346/2013***	H10N8	Feb 2014	4.3	6.0			0.2088	0.2101	0.2094	4.3	4.8
A/gyr falcon/Washington/41088/2014**	H5N8	Mar 2015	4.2	4.6	341	328	0.1532	0.3424	0.2290	3.9	4.3
A/Northern pintail/Washington/40964/2014**	H5N2	Mar 2015	3.8	4.1	341	328	0.1529	0.3799	0.2410	3.9	4.3
A/canine/Illinois/12191/2015	H3N2	Jun 2016	3.7	3.7	1000	1000	0.0607	0.1509	0.0957	4.9	4.8
A/American green-winged teal/Washington/1957050/2014	H5N1	Mar 2015	3.6	4.1	326	314	0.1911	0.4482	0.2927	4.1	4.9
A/turkey/Indiana/1573-2/2016**	H7N8	Jul 2017	3.4	3.9	495	494	0.1130	0.7738	0.2957	3.4	3.9
A/chicken/Tennessee/17-007431-3/2017	H7N9	Oct 2017	3.1	3.5	496	495	0.1027	0.2569	0.1624	4.1	4.2
A/chicken/Tennessee/17-007147-2/2017	H7N9	Oct 2017	2.8	3.5	496	495	0.2095	0.2541	0.2307	4.2	4.8

\* HA strain is not available for A/duck/New York/1996, so this strain is omitted.

\*\* Could not construct a Qnet of human sequence data available for that virus sub-type (less than 30 strains), so we constructed a Qnet using all human strains that match the HA sub-type, i.e. H5NX for H5N6.

\*\*\* These strains did not have enough human sequence data to generate a Qnet, even when only considering the HA sub-type. Thus, we estimated the risk score using every Qnet from the other IRAT strains, and took the average among NA and HA. Finally, we took the geometric mean of the resulting NA and HA averages.

SI Tab. 10  
Influenza A Strains Evaluated by IRAT and Corresponding Qnet Computed Current Risk Scores

Influenza Virus	Subype	IRAT Date	IRAT Emer-gence Score	IRAT Impact Score	HA Qnet Sample	NA Qnet Sample	HA Avg. Q-dist.	NA Avg. Q-dist.	Geom. Mean	Qnet Emer-gence Score	Qnet Impact Score
A/swine/Shandong/1207/2016	H1N1	Jul 2020	7.5	6.9	1000	1000	0.0599	0.0417	0.0500	5.8	5.8
A/Ohio/13/2017	H3N2	Jul 2019	6.6	5.8	1000	1000	0.0091	0.0692	0.0251	6.2	6.0
A/Hong Kong/125/2017	H7N9	May 2017	6.5	7.5	1000	1000	0.0092	0.0046	0.0065	6.7	6.6
A/Shanghai/02/2013	H7N9	Apr 2016	6.4	7.2	1000	1000	0.0031	0.0044	0.0037	6.8	6.6
A/Anhui-Lujiang/39/2018	H9N2	Jul 2019	6.2	5.9	58	58	0.0157	0.0467	0.0271	6.2	6.0
A/Indiana/08/2011	H3N2	Dec 2012	6.0	4.5	1000	1000	0.0176	0.0184	0.0180	6.4	6.3
A/California/62/2018	H1N2	Jul 2019	5.8	5.7	37	37	0.2038	0.0477	0.0986	5.3	5.9
A/Bangladesh/0994/2011	H9N2	Feb 2014	5.6	5.4	58	58	0.0473	0.4654	0.1484	3.8	3.6
A/Sichuan/06681/2021	H5N6	Oct 2021	5.3	6.3	46	46	0.3443	0.0600	0.1437	5.1	6.2
A/Vietnam/1203/2004	H5N1	Nov 2011	5.2	6.6	48	45	0.1323	0.0411	0.0738	5.6	5.8
A/Yunnan/14564/2015	H5N6	Apr 2016	5.0	6.6	46	46	0.2187	0.0415	0.0953	5.4	6.0
A/Astrakhan/3212/2020	H5N8	Mar 2021	4.6	5.2	95	92	0.2366	0.5451	0.3591	4.8	6.1
A/Netherlands/219/2003	H7N7	Jun 2012	4.6	5.8	1000	1000	0.1658	0.4596	0.2760	3.9	4.5
A/American wigeon/South Carolina/AH0195145/2021	H5N1	Mar 2022	4.4	5.1	48	45	0.2355	0.3135	0.2717	4.3	5.2
A/Jiangxi-Donghu/346/2013**	H10N8	Feb 2014	4.3	6.0	-1	-1	0.2097	0.2299	0.2196	4.2	4.8
A/gyrfalcon/Washington/41088/2014	H5N8	Mar 2015	4.2	4.6	95	92	0.2387	0.5438	0.3603	4.8	6.1
A/Northern pintail/Washington/40964/2014	H5N2	Mar 2015	3.8	4.1	95	92	0.2327	0.5099	0.3445	4.6	5.8
A/canine/Illinois/12191/2015	H3N2	Jun 2016	3.7	3.7	1000	1000	0.0179	0.0374	0.0259	6.2	6.1
A/American green-winged teal/Washington/1957050/2014	H5N1	Mar 2015	3.6	4.1	48	45	0.2352	0.3067	0.2686	4.3	5.1
A/turkey/Indiana/1573-2/2016	H7N8	Jul 2017	3.4	3.9	1000	1000	0.0438	0.4165	0.1351	4.0	3.8
A/chicken/Tennessee/17-007431-3/2017	H7N9	Oct 2017	3.1	3.5	1000	1000	0.0335	0.5127	0.1310	3.8	3.6
A/chicken/Tennessee/17-007147-2/2017	H7N9	Oct 2017	2.8	3.5	1000	1000	0.0839	0.5127	0.2075	3.5	3.6
A/duck/New York/1996*	H1N1	Nov 2011	2.3	2.4	1000	1000	-1	-1	-1	-1	-1

\*This table contains Qnet scores for IRAT computed using current sequence data, thereby computing the current risk of these strains. -1 indicates missing data, either from lack of human sequence data available for that virus sub-type (less than 30 strains) or missing IRAT sequence data (in the case of A/duck/New York/1996)

SI Tab. 11  
General linear model for evaluating effect of data diversity on Qnet performance

Variable Name	Description
qnet_complexity	Cumulative number of nodes in all predictors in the corresponding Qnet
data_diversity	Number of clusters in set of input sequence where each sequence in a specific cluster is separated by at least 5 mutations from sequences not in the cluster
ldistance_WHO	Deviation of WHO predicted strain from the dominant strain

```
model:dev ~ qnet_complexity + data_diversity + qnet_complexity * data_diversity + ldistance_WHO
Generalized Linear Model Regression Results
=====
Dep. Variable: dev No. Observations: 235
Model: GLM Df Residuals: 230
Model Family: Gaussian Df Model: 4
Link Function: identity Scale: 23.214
Method: IRLS Log-Likelihood: -700.43
Date: Thu, 11 Jun 2020 Deviance: 5339.2
Time: 16:45:46 Pearson chi2: 5.34e+03
No. Iterations: 3 Covariance Type: nonrobust
=====
      coef    std err     z   P>|z|    [0.025    0.975]
-----
Intercept      -0.1116    1.090   -0.102    0.918    -2.248    2.025
qnet_complexity 0.0005    0.000    1.075    0.282    -0.000    0.001
data_diversity  0.3197    0.126    2.531    0.011    0.072    0.567
qnet_complexity:data_diversity -6.932e-05  5.01e-05   -1.383    0.167    -0.000    2.89e-05
ldistance_WHO   -0.0348    0.035   -1.007    0.314    -0.102    0.033
=====
```

```
model:dev ~ qnet_complexity + data_diversity + ldistance_WHO
Generalized Linear Model Regression Results
=====
Dep. Variable: dev No. Observations: 235
Model: GLM Df Residuals: 231
Model Family: Gaussian Df Model: 3
Link Function: identity Scale: 23.306
Method: IRLS Log-Likelihood: -701.41
Date: Thu, 11 Jun 2020 Deviance: 5383.6
Time: 16:45:47 Pearson chi2: 5.38e+03
No. Iterations: 3 Covariance Type: nonrobust
=====
      coef    std err     z   P>|z|    [0.025    0.975]
-----
Intercept      1.0841    0.665    1.630    0.103    -0.219    2.387
qnet_complexity -4.12e-05  0.000   -0.156    0.876    -0.001    0.000
data_diversity  0.1788    0.075    2.392    0.017    0.032    0.325
ldistance_WHO   -0.0695    0.024   -2.930    0.003    -0.116    -0.023
=====
```

SI Tab. 12  
General linear model evaluating Qnet emergence risk predictions against IRAT estimates

---

Model: IRAT_Emergence_Score ~ Geometric_Mean						
<hr/>						
Dep. Variable:	IRAT_Emergence_Score	No. Observations:	22			
Model:	GLM	Df Residuals:	20			
Model Family:	Gaussian	Df Model:	1			
Link Function:	identity	Scale:	0.86853			
Method:	IRLS	Log-Likelihood:	-28.618			
Date:	Tue, 25 Oct 2022	Deviance:	17.371			
Time:	00:58:27	Pearson chi2:	17.4			
No. Iterations:	3	Pseudo R-squ. (CS):	0.5919			
Covariance Type:	nonrobust					
<hr/>						
	coef	std err	z	P> z	[0.025	0.975]
Intercept	6.2467	0.356	17.529	0.000	5.548	6.945
Geometric_Mean	-8.1063	1.830	-4.429	0.000	-11.693	-4.519
<hr/>						
Model: IRAT_Emergence_Score ~ Geometric_Mean + HA_Avg_Qdist*NA_Avg_Qdist						
<hr/>						
Dep. Variable:	IRAT_Emergence_Score	No. Observations:	22			
Model:	GLM	Df Residuals:	17			
Model Family:	Gaussian	Df Model:	4			
Link Function:	identity	Scale:	0.69369			
Method:	IRLS	Log-Likelihood:	-24.357			
Date:	Tue, 25 Oct 2022	Deviance:	11.793			
Time:	00:58:59	Pearson chi2:	11.8			
No. Iterations:	3	Pseudo R-squ. (CS):	0.7797			
Covariance Type:	nonrobust					
<hr/>						
	coef	std err	z	P> z	[0.025	0.975]
Intercept	6.8403	0.442	15.459	0.000	5.973	7.708
Geometric_Mean	-23.7466	9.674	-2.455	0.014	-42.707	-4.786
HA_Avg_Qdist	1.9097	3.979	0.480	0.631	-5.889	9.708
NA_Avg_Qdist	-1.8133	2.826	-0.642	0.521	-7.353	3.726
HA_Avg_Qdist:NA_Avg_Qdist	54.2280	21.474	2.525	0.012	12.139	96.317
<hr/>						

SI Tab. 13  
Numbering Conversion to pdm09 and H3 Schemes

Query	H1N1pdm	H3	Query	H1N1pdm	H3	Query	H1N1pdm	H3	Query	H1N1pdm	H3
1	-	-	77	60	69	157	140	143	-	-	-
2	-	-	78	61	70	158	141	144	-	-	-
3	-	-	79	62	71	159	142	145	-	-	-
4	-	-	80	63	72	160	143	146	238	221	224
5	-	-	81	64	73	161	144	147	239	222	225
6	-	-	82	65	74	162	145	148	240	223	226
7	-	-	83	66	75	163	146	149	241	224	227
8	-	-	84	67	76	164	147	150	242	225	228
9	-	-	85	68	77	165	148	151	243	226	229
10	-	-	86	69	78	166	149	152	244	227	230
11	-	-	87	70	79	167	150	153	245	228	231
12	-	-	88	71	80	168	151	154	246	229	232
13	-	-	89	72	81	169	152	155	247	230	233
14	-	-	90	73	82	170	153	156	248	231	234
15	-	-	91	74	-	171	154	157	249	232	235
16	-	-	92	75	83	172	155	158	250	233	236
17	-	-	93	76	84	-	-	-	251	234	237
-	-	1	94	77	85	-	-	-	252	235	238
-	-	2	95	78	86	-	-	-	253	236	239
-	-	3	96	79	87	-	-	-	254	237	240
-	-	4	97	80	88	173	156	159	255	238	241
-	-	5	98	81	89	174	157	160	256	239	242
-	-	6	99	82	90	175	158	161	257	240	243
-	-	7	100	83	91	176	159	162	258	241	244
-	-	8	101	84	92	177	160	163	259	242	245
-	-	9	102	85	-	178	161	164	260	243	246
-	-	10	103	86	93	179	162	165	261	244	247
18	1	11	104	87	94	180	163	166	262	245	248
19	2	12	105	88	95	181	164	167	263	246	249
20	3	13	106	89	96	182	165	168	264	247	250
21	4	14	107	90	97	183	166	169	265	248	251
22	5	15	108	91	98	184	167	170	266	249	252
23	6	16	109	92	99	-	-	-	267	250	253
24	7	17	110	93	100	185	168	171	268	251	254
25	8	18	111	94	101	186	169	172	269	252	255
26	9	19	112	95	102	187	170	173	270	253	256
27	10	20	-	-	-	-	-	-	271	254	257
28	11	21	-	-	-	188	171	174	272	255	258
29	12	22	113	96	103	189	172	175	273	256	259
30	13	23	114	97	104	190	173	176	274	257	260
31	14	24	115	98	105	191	174	177	275	258	261
32	15	25	116	99	106	192	175	178	276	259	262
33	16	26	117	100	107	193	176	179	-	-	-
34	17	27	118	101	108	194	177	180	-	-	-
35	18	28	119	102	109	195	178	181	-	-	-
36	19	29	120	103	110	196	179	182	-	-	-
37	20	30	121	104	111	197	180	183	-	-	-
38	21	31	122	105	112	198	181	184	-	-	-
39	22	32	123	106	113	199	182	185	-	-	-
40	23	33	124	107	114	200	183	186	-	-	-
41	24	34	125	108	115	201	184	187	-	-	-
42	25	35	126	109	116	202	185	188	277	260	-
43	26	36	127	110	117	203	186	189	278	261	263
44	27	37	128	111	118	204	187	190	279	262	264
45	28	38	129	112	119	205	188	191	280	263	265
46	29	39	130	113	120	206	189	192	281	264	266
47	30	40	131	114	121	207	190	193	282	265	267
48	31	41	132	115	122	208	191	194	283	266	268
49	32	42	133	116	123	209	192	195	284	267	269
50	33	43	-	-	-	210	193	196	285	268	270
51	34	44	-	-	-	211	194	197	286	269	271
52	35	45	134	117	124	212	195	198	287	270	272
53	36	46	135	118	125	213	196	199	288	271	273
54	37	47	136	119	-	-	-	-	289	272	274
55	38	48	137	120	-	214	197	200	290	273	275
56	39	49	138	121	-	215	198	201	291	274	276
57	40	50	139	122	126	216	199	202	292	275	277
58	41	51	140	123	127	217	200	203	293	276	278
59	42	52	141	124	128	218	201	204	294	277	279
60	43	53	-	-	-	219	202	205	295	278	280
61	44	54	-	-	-	220	203	206	296	279	281
62	45	-	-	-	-	221	204	207	297	280	282
63	46	55	-	-	-	222	205	208	298	281	283
64	47	56	-	-	-	223	206	209	299	282	284
65	48	57	142	125	129	224	207	210	300	283	285
66	49	58	143	126	130	225	208	211	-	-	-
67	50	59	144	127	131	226	209	212	301	284	286
68	51	60	145	128	132	227	210	213	302	285	287
-	-	-	146	129	133	228	211	214	303	286	288
-	-	-	147	130	-	229	212	215	304	287	289
-	-	-	148	131	134	230	213	216	305	288	290
-	-	-	149	132	135	231	214	217	306	289	291
-	-	-	150	133	136	232	215	218	307	290	292
69	52	61	151	134	137	233	216	219	308	291	293
70	53	62	152	135	138	234	217	220	309	292	294
71	54	63	153	136	139	235	218	221	310	293	295
72	55	64	154	137	140	236	219	222	311	294	296
73	56	65	155	138	141	237	220	223	-	-	-
74	57	66	-	-	-	-	-	-	312	295	297
75	58	67	156	139	142	-	-	-	313	296	298