

# opioid hysteria and the social prejudice against the chronically ill - Jarif Wasif.pdf

*by Sanaul Haque*

---

**Submission date:** 13-Oct-2025 11:08PM (UTC+0700)

**Submission ID:** 2779950057

**File name:** opioid\_hysteria\_and\_the\_social\_prejudice\_against\_the\_chronically\_ill\_-\_Jarif\_Wasif.pdf (276.95K)

**Word count:** 1895

**Character count:** 10848

## **IARCO RESEARCH PROPOSAL**

**Full Legal Name:** Jarif Wasif

**Institution:** Methodist English Medium School

**Category:** Junior

**Class/Grade/Year:** A-Level

**Country:** Bangladesh

**Major (if senior):** N/A

**Submission Date:** September 30, 2025

**Registered Email Address:** jarif3408@gmail.com

**Research Topic:** Opioid hysteria and the <sup>10</sup> social prejudice against the chronically ill

# OPIOID HYSTERIA AND THE SOCIAL PREJUDICE AGAINST THE CHRONICALLY ILL

## Research Problem

In 2023, 24.3% of adults in the USA had chronic pain, 8.5% of whom had high-impact chronic pain that limits daily life or work [1]. Diseases such as fibromyalgia, Ehlers-Danlos syndrome, and Complex regional pain syndrome are incurable conditions that coincide with unbearable pain and fatigue, which severely impair an individual's life. For this reason, the people in chronic pain states have a significant prevalence of anxiety and depression (A/D) symptoms, with 39.5% of those with high-impact chronic pain having A/D symptoms along with a high risk for suicide death. [2] [3]. Thus, many chronically ill patients rely on opioid related pain meds as a tool to ease their pain and allow them to perform basic tasks. However, the ongoing opioid crisis, marked by a sharp rise in opioid overdose deaths, has resulted in the portrayal of these substances as addictive drugs that are exclusively taken as a measure to escape into a deep sense of euphoria and thus inducing a stigma against chronically ill patients who are dependent on these medications. Misinterpretation of this crisis has prompted the DEA to issue increasingly strict and often vague guidelines on opioid prescribing. While intended to curb misuse, these regulations carry serious consequences: even well-meaning practitioners risk investigation or loss of their medical license if their prescriptions are deemed illegitimate. As a result, many healthcare providers have grown fearful and reluctant to prescribe opioids, even when medically justified, leaving patients with chronic or severe pain caught in the crossfire. [4][5]

## Existing Literature

Studies have brought attention to the unfair treatment of the chronically ill, highlighting both the psychological and physiological toll of discrimination. An example is a study published in SSM Population Health, which notes that 18.4% of adults aged 50 and older in the United States with chronic illnesses like diabetes, heart disease, or lung disease reported experiencing discrimination among healthcare facilities, where their interactions with these places are defined by a lack of respect and poor service [6]. These adverse experiences had strong correlations with lower provider trust, lower satisfaction with care, and poor self-reported health. Another study, published in Psychoneuroendocrinology, pointed out that individuals exposed to repetitive unfair treatment in a medical context or everyday life scored much higher on allostatic load, a physiological stress biomarker [7]. This increased biological load denotes the physical consequences of decades-long discrimination and stress that can accelerate the progression of chronic illnesses and weaken overall well-being. Notwithstanding such a challenge and the role that opioids have played in pain management among many chronically ill persons in the United States, there still exists a significant gap in research comparing the dangers and disadvantages associated with restricting prescription opioids and whether the guidelines were able to overcome the opioid crisis. Moreover, there lies a severe scarcity in the literature on chronic pain patients' experiences due to opioids being depicted as psychoactive drugs that are used by addicts to get into a psychedelic state. Understanding the impact of this stigmatization and the risk-reward of prescription opioids is thus crucial for helping chronically ill patients alleviate their pain, mentally and physically.

## Research Question

→ Does the fearmongering of opioids promoted by societal stigmatization correlate with declining mental health amongst chronically ill patients?

→ Does suppressing prescriptions result in a lower overdose?

## Methodology

This study is based upon various complementary datasets based on the US spanning from 1999-2023. The research utilizes a mixed-methods approach; combining quantitative trend analysis and comparative assessments on the relevant secondary data to determine the risk-reward balance between prescribing necessary opioids versus the danger of non-prescribed patients seeking illicit alternatives, considering factors such as overdose rates, long-term mental health decline, and suicide mortality.

### Datasets:

1. Prescription opioid overdose death rate in the US (1999-2023) [8]
  - Variables: Annual death rates per 100,000 population, temporal trends
  - Source: Statista database on US prescription opioid overdose mortality
2. Total Opioid Overdose deaths [9]
  - Variables: Total overdose deaths, specific causes
  - Source: CDC National Center for Health Statistics
3. Mental Health Outcomes following Opioid discontinuation [10] [11]
  - Variables: Suicide rates among chronic pain patients, depression and anxiety measures
  - Source: PubMed studies
4. Non-prescription Opioid use patterns [12]
  - Variables: Substance abuse patterns, illicit drugs procurement
  - Source: PLOS ONE article

## Analytical Method

### Quantitative Analysis:

1. **Trend Analysis-** A time-series examination of prescription opioid overdose rates between 1999-2023 is conducted initially, and then the annual percentage changes are calculated and inflection points are identified. Correlation analysis done between restrictions and overdose trends.
2. **Comparative Assessment-** At first, a comparison is derived between overdose rates between prescription and non-prescription opioids and risk-ratio calculations between different patient populations is calculated. Demographics are considered.
3. **Outcome examination-** The outcomes measured are mental health decline rates following opioid discontinuation, suicide mortality rates amongst chronic pain patients and illicit drug use prevalence of patients who were denied necessary opioids for treatment.

**Project Practicalities:**

Secondary data will be acquired and cleaned within the first week. A comprehensive statistical analysis will be conducted in the subsequent 2 weeks and the results will be interpreted and arranged in the week after. The final manuscript will be prepared and submitted in the 4th week of the study's duration. The total estimated time to complete the research is 4 weeks.

The study population includes vulnerable populations: chronic illness and substance use disorder. Hence, analysis will be done sensitively to avoid stigmatization of such demographics. The results will be objective and rational without any perpetuating emotional biases. The limitations of secondary data analysis will be acknowledged clearly. Findings will be contextualized solely to present evidence on the hypothesis, and harmful narratives will be avoided strictly. Moreover, the results are intended to advocate for improved patient care rather than reinforcing negative stereotypes amongst these populations.

**6. Roadblocks and Potential Limitations:**

This research relies entirely on secondary data in the U.S., since time and resource constraints prevent primary data collection. The latest data sets might not be readily available, and some secondary sources might be biased. Additionally, variables such as mental health deterioration or suicide may not be affected only by the denial of prescriptions, as some other factors might overlap, making causality harder to establish.

**7. Expected outcomes:**


An examination of data regarding the deaths involving both prescribed and non-prescribed opioids can help us determine whether legitimate prescription of opioids can play a significant role in the general opioid overdose epidemic. The difference within is critical to the distinction between the effects of illicit drug use and medically controlled pain management. Moreover, questioning the necessity of limiting access to opioid medication for patients with chronic conditions, as a result worsening the quality of their lives, increasing the possible transition to illegal sources, we will be able to judge whether all this is worth limiting access to drugs or whether this measure is harming rather than helping. The current analysis can assist in greater opioid policy-making, one that safeguards the health of the population at large and does not miss the population struggling with chronic pain.

**8. Conclusion:**

The methodology described seeks to test the hypothesis that reducing prescription rates for patients who require opioids for pain management not only fails to achieve the intended safety outcomes but also tends to exacerbate the risks these patients might face. If the study supports the hypothesis, this research will present evidence that will compel fewer restrictions on opioid based treatments for documented medical conditions. Moreover, the findings will help humanize chronically ill patients and remove the stigma that comes from taking narcotics for legitimate symptoms. The study will help recognize the importance of opioid therapy for chronic pain conditions whilst tackling the harm that current stringent regulatory practices bring forth.

**References:**

- [1] J. W. Lucas and I. Sohi, "Chronic Pain and High-Impact Chronic Pain in U.S. Adults, 2023," *NCHS Data Brief*, no. 518, Oct. 2024. [Online]. Available: <https://doi.org/10.15620/cdc/169630>
- [2] J. S. De La Rosa, B. R. Brady, M. M. Ibrahim, K. E. Herder, J. S. Wallace, A. R. Padilla, and T. W. Vanderah, "Co-occurrence of chronic pain and anxiety/depression symptoms in U.S. adults: Prevalence, functional impacts, and opportunities," *Pain*, vol. 165, no. 3, pp. 666–673, Mar. 2024. [Online]. Available: <https://doi.org/10.1097/j.pain.0000000000003056>
- [3] A. L. Hassett, J. K. Aquino, and M. A. Ilgen, "The risk of suicide mortality in chronic pain patients," *Current Pain and Headache Reports*, vol. 18, p. 436, 2014. [Online]. Available: <https://doi.org/10.1007/s11916-014-0436-1>
- [4] Y. Heung, M. Clark, J. Tschanz, and E. Bruera, "Opioid prescription denials by community pharmacies for cancer-related pain: A case series," *Journal of Pain and Symptom Management*, vol. 66, no. 3, pp. e431–e435, 2023. [Online]. Available: <https://doi.org/10.1016/j.jpainsymman.2023.06.027>
- [5] K. Corredor, "Reports of refusals to fill buprenorphine prescriptions hits new high, but overall reports down," *The ASAM Weekly*, Oct. 31, 2024. [Online]. Available: <https://www.asam.org/publications-resources/the-asam-weekly/detail/2024/10/31/reports-of-refusals-to-fill-buprenorphine-prescriptions-hits-new-high-but-overall-reports-down>
- [6] K. A. Ajillo, "Opiophobia and the tragedy of needless pain: A call for education and balance," *Pharmacology, Biochemistry and Behavior*, vol. 230, p. 173616, 2023. [Online]. Available: <https://doi.org/10.1016/j.pbb.2023.173616>
- [7] J. J. Bettinger, W. Amarquaye, J. Fudin, and M. E. Schatman, "Misinterpretation of the 'overdose crisis' continues to fuel misunderstanding of the role of the prescription opioids," *Journal of Pain Research*, vol. 15, pp. 949–958, Apr. 2022. [Online]. Available: <https://doi.org/10.2147/JPR.S367753>
- [8] Statista Research Department, "Death rate from prescription opioid overdose in the U.S. from 1999 to 2023," *Statista*, Jul. 9, 2025. [Online]. Available: <https://www.statista.com/statistics/638335/death-rate-prescription-opioid-overdose-us/>
- [9] Centers for Disease Control and Prevention, "Provisional drug overdose death counts," *National Center for Health Statistics*, May 15, 2024. [Online]. Available: [https://www.cdc.gov/nchs/pressroom/nchs\\_press\\_releases/2024/20240515.htm](https://www.cdc.gov/nchs/pressroom/nchs_press_releases/2024/20240515.htm)
- [10] T. L. Mark and C. Alexander, "Discontinuation of long-term opioid therapy in patients with chronic pain: A comparative effectiveness study," *Journal of General Internal Medicine*, vol. 35, no. 10, pp. 2920–2927, 2020. [Online]. Available: <https://doi.org/10.1007/s11606-020-05761-w>
- [11] M. I. Demidenko, K. L. Dobscha, and S. K. Dobscha, "Suicidal ideation and suicide attempt following discontinuation of long-term opioid therapy for pain," *Pain*, vol. 162, no. 8, pp. 2186–2194, 2021. [Online]. Available: <https://doi.org/10.1097/j.pain.0000000000002201>

[12] H. Guarino, P. Mateu-Gelabert, L. Elliott, A. S. Bennett, and R. rtis, "Nonmedical opioid use and pathways of risk among formerly prescribed opioid users," *PLOS ONE*, vol. 15, no. 3, p. e0230408, 2020. [Online]. Available: <https://doi.org/10.1371/journal.pone.0230408>

# opioid hysteria and the social prejudice against the chronically ill - Jarif Wasif.pdf

## ORIGINALITY REPORT

15%

SIMILARITY INDEX

12%

INTERNET SOURCES

12%

PUBLICATIONS

0%

STUDENT PAPERS

## PRIMARY SOURCES

- |   |   |    |
|---|---|----|
| 1 | Karlyn A. Edwards, Michele J. Buonora, Jessica S. Merlin, Jane M. Liebschutz. "Recent advances in the treatment of chronic pain and substance use disorders", Current Opinion in Psychology, 2025<br>Publication  | 1% |
| 2 | <a href="http://www.statista.com">www.statista.com</a><br>Internet Source   | 1% |
| 3 | <a href="http://www.ncbi.nlm.nih.gov">www.ncbi.nlm.nih.gov</a><br>Internet Source   | 1% |
| 4 | <a href="http://www.uvmcora.org">www.uvmcora.org</a><br>Internet Source   | 1% |
| 5 | <a href="http://researchinformation.amsterdamumc.org">researchinformation.amsterdamumc.org</a><br>Internet Source   | 1% |
| 6 | Christiane Bertachini Lombello, Nasser Ali Daghashtanli, Hermann Windisch Neto, Fernando Luiz Affonso Fonseca et al. "Characterization of DELUX: Ultraviolet light sterilization device for PFF2 / N95 masks against COVID-19", Ingenius, 2021<br>Publication | 1% |
| 7 | Grunwell, Daniel, Baptista, Paulo, Campos, Sergio, Sahama, Tony. "Managing and sharing health data through Information Accountability protocols", 'Institute of   | 1% |



# Electrical and Electronics Engineers (IEEE)', 2015

Internet Source

8

[www.researchprotocols.org](http://www.researchprotocols.org)

Internet Source

1 %

9

Campbell, Gabrielle, Raimondo Bruno, Shane Darke, Fiona Shand, Wayne Hall, Michael Farrell, and Louisa Degenhardt. "Prevalence and Correlates of Suicidal Thoughts and Suicide Attempts in People Prescribed Pharmaceutical Opioids for Chronic Pain :", Clinical Journal of Pain, 2015.

Publication

1 %

10

Jung-won Lim. Health and Quality of Life Outcomes, 2004

Publication

1 %

11

Patty Wilson, Tracey Day. "Benzodiazepine Withdrawal in the Context of Benzodiazepine-Contaminated Opioids: Practice Implications", The Journal for Nurse Practitioners, 2024

Publication

1 %

12

[ijcem.in](http://ijcem.in)

Internet Source

1 %

13

[ouci.dntb.gov.ua](http://ouci.dntb.gov.ua)

Internet Source

1 %

14

Federico Mento, Antonio Di Sabatino, Anna Fiengo, Umberto Sabatini et al. "Automatically Scoring Lung Ultrasound Videos of COVID-19 and post-COVID-19 Patients", 2022 IEEE International Ultrasonics Symposium (IUS), 2022

Publication

1 %

[www.medrxiv.org](http://www.medrxiv.org)

15

Internet Source

1 %

16

Elaine L. Miller, Robert V. Miller. "Does Music Have a Role in Pain Management?", Pain Management Nursing, 2025

Publication

&lt;1 %

17

Patrick Coyne, Sarah Lowry, Carol Mulvenon, Judith A. Paice. "American Society for Pain Management Nursing and Hospice and Palliative Nurses Association Position Statement: Pain Management at the End of Life", Pain Management Nursing, 2024

Publication

&lt;1 %

18

[www.mdpi.com](http://www.mdpi.com)

Internet Source

&lt;1 %

19

Jeffrey J Bettinger, William Amarquaye, Jeffrey Fudin, Michael E Schatman. "Misinterpretation of the "Overdose Crisis" Continues to Fuel Misunderstanding of the Role of Prescription Opioids", Journal of Pain Research, 2022

Publication

&lt;1 %

20

[www.chcs.org](http://www.chcs.org)

Internet Source

&lt;1 %

21

John V. Prunskis, Tadas Masys, Stephen T. Pyles, Alaa Abd-Elseyed et al. "The Application of Artificial Intelligence to Enhance Spinal Cord Stimulation Efficacy for Chronic Pain Management: Current Evidence and Future Directions", Current Pain and Headache Reports, 2025

Publication

&lt;1 %

22 Lauren A McCormack, Mark J Edlund, Sonia M Thomas, Li-Tzy Wu et al. "Effectiveness of motivational interviewing plus cognitive behavioral therapy versus shared decision making for voluntary opioid tapering in patients with chronic pain: The INSPIRE randomized pragmatic trial", Pain Medicine, 2025

Publication

<1 %

23 Keith A. Trujillo. "Opiophobia and the tragedy of needless pain: A call for education and balance", Pharmacology Biochemistry and Behavior, 2023

Publication

<1 %

Exclude quotes On Exclude matches Off

Exclude bibliography Off