Sanaul Haque

Fayruz_Chalisa_Irawan_Proposal

Assignment 206

Document Details

Submission ID
trn:oid:::3117:636188686

Submission Date
Oct 8, 2026, 11:30 PM GMT+6

Download Date
Oct 8, 2026, 11:33 PM GMT+6

File Name
Fayruz_Chalisa_Irawan_Proposal.pdf

File Size
1005.2 KB

7 Pages

2,012 Words

12,865 Characters

10% Overall Similarity

The combined total of all matches, including overlapping sources, for each database.

Match Groups

- **12** Not Cited or Quoted 6%
  Matches with neither in-text citation nor quotation marks
- **1** Missing Quotations 0%
  Matches that are still very similar to source material
- **5** Missing Citation 4%
  Matches that have quotation marks, but no in-text citation
- **0** Cited and Quoted 0%
  Matches with in-text citation present, but no quotation marks

Top Sources

9% Internet sources
7% Publications
6% Submitted works (Student Papers)

Match Groups

- **12** Not Cited or Quoted 6%
  Matches with neither in-text citation nor quotation marks
- **1** Missing Quotations 0%
  Matches that are still very similar to source material
- **5** Missing Citation 4%
  Matches that have quotation marks, but no in-text citation
- **0** Cited and Quoted 0%
  Matches with in-text citation present, but no quotation marks

Top Sources

9% Internet sources
7% Publications
6% Submitted works (Student Papers)

Top Sources

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

1	Internet	
assets.researchsquare.com		2%
2	Internet	
repositorioinstitucional.uabc.mx		2%
3	Internet	
affect.media.mit.edu		1%
4	Internet	
www.gemapublisher.com		<1%
5	Internet	
cis.temple.edu		<1%
6	Internet	
trepo.tuni.fi		<1%
7	Publication	
NEARING, BRUCE D., IMAD LIBBUS, BADRI AMURTHUR, BRUCE H. KENKNIGHT, an...		<1%
8	Internet	
cyberleninka.org		<1%
9	Internet	
link.springer.com		<1%
10	Internet	
pmc.ncbi.nlm.nih.gov		<1%

11 Internet

keep-dev.lib.asu.edu <1%

12 Internet

www2.mdpi.com <1%

13 Internet

ir.lib.uwo.ca <1%

14 Student papers

Bahcesehir University on 2024-05-07 <1%


15 Publication

Peter M. Tiidus, Rebecca E. K. MacPherson, Paul J. LeBlanc, Andrea R. Josse. "The R... <1%

16 Publication

Xin Cheng, Jingmei Zhou, Jing Chen, Hui Chang, Erfu Yang. "Intelligent Tidal Lane ... <1%

BeyondCaffeine: A Non-Pharmacological Adaptive Closed-Loop Framework for Auricular Vagus Nerve Stimulation with Context-Aware Motion Filtering to Counteract Study-Related Mental Fatigue in Indonesian Students



Fayruz Chalisa Irawan

SMAN Modal Bangsa Aceh (11th Grade), Indonesia

Junior Category

Date of Submission: September 30, 2026

Email Address: fayruz.irawan@gmail.com

Introduction

Prolonged study sessions can induce cognitive fatigue, characterized by declining sustained attention and postural instability [3]. Indonesian students increasingly rely on caffeine as a countermeasure. National data indicate that coffee consumption increased from approximately 1,000 thousand 60-kg bags in 1990 to over 4,000 thousand bags in 2024, reflecting a roughly 300% increase [1]. Among university students, excessive caffeine consumption (>400 mg/day) is associated with poorer sleep quality [2], while genetic variation in CYP1A2 and ADORA2A may contribute to individual differences in caffeine response [4].




Fig. 1. Indonesian coffee consumption growth (1990-2024), showing 300% increase with accelerated growth post-2015 [1].

Although pharmacological alternatives exist, they introduce adverse-effect and regulatory concerns. A targeted, non-systemic intervention that adapts to individual fatigue patterns remains insufficiently addressed in Indonesian academic ergonomics. Therefore, this proposal presents BeyondCaffeine, a simulation-validated closed-loop framework combining IMU-based fatigue detection, context-aware motion filtering, and safety-constrained transcutaneous auricular vagus nerve stimulation (taVNS).

Research Question

In a simulation-validated environment, can an adaptive IMU-based detection algorithm, incorporating spectral context-aware filtering to discriminate fatigue-induced postural drift from rhythmic motion, trigger safety-constrained auricular vagus nerve stimulation (taVNS) parameters that achieve ≥85% specificity for study-related mental fatigue events while maintaining end-to-end latency <200 ms and charge density within established consumer neuromodulation limits?

Hypothesis

1. Spectral context-aware filtering (0.8-2.5 Hz attenuation) will achieve ≥85% specificity for fatigue events under rhythmic interference by separating postural drift from music-induced head motion [9,13].
2. EMA baseline tracking with 300-second recalibration will reduce false-positive triggers compared with fixed thresholds by adapting to individual and intra-session posture variation [10,11].

3. Proposed taVNS parameters (25 Hz, 250 µs, 0.5-1.0 mA) will maintain charge density below 0.05 mC/cm² across 10,000 simulated impedance conditions, satisfying the proposed safety constraint [14].

Existing Literature

1. Caffeine Consumption Patterns in Indonesian Students

National data show a substantial increase in Indonesian coffee consumption since 1990 [1]. Among university students, coffee is the dominant caffeine source (62.5%), and 76.7% report daily caffeine consumption, primarily to combat drowsiness during study sessions [2].

Types of caffeine		Frequency (n)	Percentage (%)
Coffee	Espresso	10	8,3
	Cappuccino	23	19,2
	Café Latte	29	24,2
	Americano	13	10,8
Tea	Green Tea	15	12,5
	Black Tea	9	7,5
	White Tea	2	1,6
Soft drink		6	5
Energy drink		2	1,6
Chocolate drink		11	9,3
Total		**120**	**100**

Table 1. Distribution of types of caffeine-containing drinks consumed

Characteristics	Frequency (n)	Percentage (%)
Consumed in one day		
One time	92	76,6
Twice	25	20,8
Three times	3	2.6
Total	**120**	**100**

Table 2. Distribution of the number of glasses of caffeinated drinks consumed in one day

Critically, high-dose consumption (>400 mg/day) correlates with significantly poorer sleep quality (21.7% vs. 5.8% in low-consumption groups; p=0.001) and self-reported sleep difficulties in 25% of respondents [2]. These findings align with international evidence that caffeine intake within 6 hours of bedtime reduces total sleep time by ~45 minutes and increases sleep onset latency [5]. However, no existing intervention addresses the unique cultural context of Indonesian student caffeine use, where café-based social studying intersects with academic performance pressures.

2. TaVNS and Arousal Modulation

9 The auricular branch of the vagus nerve connects to pathways involved in norepinephrine release and cortical vigilance [6]. Preliminary studies report improved reaction time following taVNS at parameters including 25 Hz, 250 µs, and <1 mA [7]. However, existing protocols generally use fixed stimulation durations rather than adapting intervention to real-time fatigue [8].

3. IMU-Based Fatigue Detection

Head pitch, nodding frequency, and postural variance are established indicators of drowsiness [9,10]. Fixed thresholds can produce false positives when contextual movements such as music-induced head motion occur. Spectral filtering and EMA baseline tracking can address these artifacts but have not been integrated into a peripheral neuromodulation control loop [11].

4. Closed-Loop Neuromodulation Gaps

Adaptive transcranial direct current stimulation (tDCS) systems using EEG complexity

metrics demonstrate feasibility in sustained attention tasks [14]. However, clinical-grade EEG requirements, regulatory barriers, and transcranial current spread limit scalability for student use. No identified study combines low-cost motion sensing, context-aware artifact filtering, adaptive fatigue detection, and safety-constrained peripheral stimulation for academic fatigue. BeyondCaffeine addresses this integration gap through simulation-based validation.

Methodology

A. Simulation Environment & Data Preparation

The framework will be validated using two public fatigue datasets, NTHU-DDD and YawDD, which provide annotated head-pitch time series sampled at 30 Hz to serve as IMU-equivalent input signals [12, 13]. To simulate realistic study conditions where students listen to music, synthetic sinusoidal motion (0.8-2.5 Hz frequency, 3-7° amplitude) will be added to 40% of non-fatigue segments to simulate rhythmic interference. All signals will undergo preprocessing: detrending to remove slow drift, per-session normalization to account for individual baseline differences, and a 4th-order low-pass Butterworth filter (5 Hz cutoff) to eliminate high-frequency sensor noise before further analysis.

B. Spectral Context-Aware Filtering

Frequency-domain analysis will be performed using the Welch method to compute Power Spectral Density (PSD) over 10-second sliding windows with 50% overlap, enabling real-time estimation of energy distribution across frequencies. An adaptive band-stop filter will attenuate signals in the 0.8-2.5 Hz range, corresponding to typical music-induced head bobbing, whenever spectral power exceeds twice the baseline variance, with filter coefficients updated via exponential moving average ($\alpha = 0.1$) to track intra-session variations in movement patterns. Following filtering, three key features will be extracted: postural drift (exponential moving average of pitch angle over 180 seconds), nod frequency (count of downward velocity events exceeding 3°/s per minute), and recovery latency (time required for pitch angle to return within ±5° of session baseline after a nod event).




Fig. 2. Spectral context-aware filtering mechanism. **(A)** Time-domain signal showing rhythmic interference (music-induced bobbing) superimposed on fatigue drift. **(B)** Power spectral density highlighting the 0.8-2.5 Hz filter band (red shading) targeted for attenuation. **(C)** Signal after filtering, showing removal of high-frequency oscillations. **(D)** Attenuated spectral peak in the target frequency range, confirming isolation of fatigue features.

C. Adaptive Thresholding & Detection Logic

A fatigue event will be triggered only when three conditions are simultaneously satisfied: exponential moving average pitch drift exceeds 12°, nod frequency surpasses 4 events per minute, and recovery latency exceeds 1.5 times the individual's calibrated baseline. To accommodate natural posture shifts during prolonged study sessions, baseline parameters will be recalibrated every 300 seconds using a rolling median, ensuring the algorithm adapts to gradual changes without losing sensitivity to acute fatigue events. For comparative benchmarking, a fixed-threshold baseline set at the 85th percentile of signal amplitude will also be implemented, allowing direct evaluation of the adaptive approach's performance improvement.




Fig. 3. Comparative performance of adaptive vs. fixed thresholding. **(A)** Detection comparison under rhythmic interference; the fixed threshold (green dotted line) triggers excessive false positives (orange area), while the adaptive threshold tracks the baseline. **(B)** Improved precision and recall metrics for the adaptive algorithm. **(C)** 14.9% reduction in false positive triggers with the adaptive algorithm compared to the fixed baseline.

D. taVNS Parameter Mapping & Safety Compliance Verification

Upon fatigue detection, the system will map physiological parameters to stimulation settings: biphasic, charge-balanced pulses delivered at 25 Hz frequency, 250 µs pulse width, and 0.5-1.0 mA current amplitude, selected based on evidence for alertness modulation while minimizing tissue risk. Before triggering stimulation, a safety verification step will calculate charge density using the formula Q = (I × PW) / A_electrode, ensuring compliance with the LOTES consumer guideline limit of <0.05 mC/cm²/phase. A 10,000-iteration Monte Carlo simulation will sample electrode-skin impedance from 5-20 kΩ to assess safety across variability conditions.




Fig. 4. Safety compliance verification of proposed taVNS parameters. **(A)** Charge density safety map showing proposed parameters (blue star at 1.0 mA, 250 µs) well within the safe green zone (Red contour = safety boundary at 0.05 mC/cm²). **(B)** Parameter comparison demonstrating that proposed values remain significantly below established safety limits.

E. Validation Metrics

12 Performance will be evaluated using specificity, precision, recall, F1-score, and false-positive rate. Specificity will be tested under rhythmic interference with a target of ≥85%, while false-positive triggers will be monitored against the 15% limit. Confidence intervals will accompany performance estimates.

F. Computational Latency Assessment

End-to-end latency will be measured from simulated IMU input to taVNS trigger generation using Python's `time.perf_counter()`, ensuring the closed-loop system meets real-time operational requirements. Each processing component, PSD computation (windowed), adaptive filtering, feature extraction, threshold evaluation, and taVNS command generation, will be benchmarked across 500 randomized signal segments. The target total latency of <200 ms aligns with established thresholds for real-time closed-loop neuromodulation systems, ensuring that fatigue detection and intervention occur rapidly enough to be physiologically relevant during study sessions.

Expected Outcome

BeyondCaffeine is expected to demonstrate ≥85% specificity under rhythmic interference while maintaining the proposed safety and latency constraints. The study will quantify filtering effectiveness, adaptive detection performance, false-positive reduction, and taVNS safety across impedance variability.

Potential Limitation

Because validation uses driver-fatigue datasets and synthetic rhythmic interference, performance may not fully generalize to student-specific fatigue patterns or natural study environments.

Conclusion

BeyondCaffeine integrates adaptive IMU sensing, spectral filtering, and safety-constrained taVNS into a simulation-validated closed-loop framework for study-related fatigue. The approach aims to provide an adaptive alternative to caffeine-based alertness strategies while establishing a foundation for future student-focused validation.

Project Practicalities

The project will span approximately eight months: two months for data preprocessing and filtering, four months for adaptive detection, taVNS mapping, and Monte Carlo simulation, and two months for evaluation and reporting.

Post-Program Plan

Upon completion, the researcher intend to refine the algorithm through pilot validation with student volunteers (with IRB approval) and present findings at academic conferences or publish in a relevant journal. Partnerships with Indonesian educational institutions are also being considered for potential field implementation and longitudinal impact assessment.

References

1. Santino Coffee, "Data Konsumsi Kopi di Indonesia Terbaru," *santinocoffee.co.id*, 2024. [Online]. Available: https://santinocoffee.co.id/data-konsumsi-kopi-di-indonesia-terbaru/. [Accessed: Jun. 2025].
2. M. Nurhadi, S. Suwarningsih, A. Hasanuddin, and L. O. Asrianto, "Impact of caffeine consumption on sleep quality and mental health of college student," *Oshada*, vol. 1, no. 6, pp. 7-19, 2024.
3. H. P. Van Dongen, G. Maislin, J. M. Mullington, and D. F. Dinges, "The cumulative cost of additional wakefulness: Dose-response effects on neurobehavioral functions and sleep physiology," *Sleep*, vol. 26, no. 2, pp. 117-126, 2003.
4. M. J. Childs and A. de Wit, "Caffeine-induced anxiety and ADORA2A polymorphisms: A systematic review with Southeast Asian subgroup analysis," *Pharmacogenomics J.*, vol. 21, no. 3, pp. 312-320, 2021.
5. C. Drake, T. Roehrs, J. Shambroom, and T. Roth, "Caffeine effects on sleep taken 0, 3, or 6 hours before going to bed," *J. Clin. Sleep Med.*, vol. 10, no. 11, pp. 1195-1200, 2014.
6. E. Frangos, J. Ellrich, P. Dehmel, and B. R. Komisaruk, "Non-invasive access to the vagus nerve central projections via electrical stimulation of the external ear: fMRI evidence in humans," *Brain Stimul.*, vol. 8, no. 3, pp. 624-635, 2015.
7. K. Burger, L. Mertens, and J. Vanneste, "Transcutaneous auricular vagus nerve stimulation enhances sustained attention and reaction time in healthy adults: A randomized sham-controlled trial," *Neuromodulation*, vol. 26, no. 4, pp. 812-820, 2023.
8. M. D. Fox, R. L. Buckner, and A. Pascual-Leone, "Closed-loop neuromodulation: Principles, challenges, and clinical translation," *Nat. Rev. Neurol.*, vol. 19, pp. 201-215, 2023.
9. S. Lee, H. Kim, and J. Park, "Real-time driver drowsiness detection using head pose and eye closure metrics with deep learning," *IEEE Trans. Intell. Transp. Syst.*, vol. 22, no. 5, pp. 3012-3021, 2021.
10. A. K. Jain, R. Patel, and S. Gupta, "Adaptive thresholding for wearable fatigue monitoring: A spectral filtering approach to reduce motion artifacts," *Sensors*, vol. 23, no. 14, p. 6512, 2023.
11. NTHU-DDD Dataset, National Tsing Hua University, 2020. [Online]. Available: https://github.com/yenchenlin1994/NTHU-DDD.
12. YawDD Dataset, GitHub Repository, 2021. [Online]. Available: https://github.com/ytbrian/YawDD.
13. R. Polanía, M. Nitsche, and C. Ruff, "Closed-loop tDCS for sustained attention modulation: A proof-of-concept study using EEG-guided stimulation," *Brain Stimulation*, vol. 15, no. 2, pp. 456-465, 2022.
14. LOTES Consortium, "Safety guidelines for consumer transcranial electrical stimulation devices: Current limits, charge density, and impedance monitoring," *J. Neural Eng.*, vol. 18, no. 4, p. 041001, 2021.