


Replica Exchange Simulations of Protein-Protein Binding and Multi- protein Complex Formation

Youngchan Kim and Gerhard Hummer



Laboratory of Chemical Physics

National Institute of Diabetes and Digestive and Kidney Diseases


National Institutes of Health

February 3, 2009

NIH Biowulf Symposium

Background

- Many biological functions are carried out by large, multi-protein assemblies
 - DNA transcriptional regulation
 - Signal transduction
 - Nuclear pore complex
 - Membrane-protein trafficking
 - Viral entry and release



ESCRT machinery (Membrane trafficking)

Motivation

Understanding structure and dynamics of multi-protein assemblies

- Many multi-protein assemblies form only transiently
 - Held together by relatively weak pairwise interactions ($K_d > 1\mu\text{M}$)
 - Multi-protein assemblies contain unstructured regions
 - Flexible polymeric linkers connecting structured domains
- ⇒ Challenges for traditional structural approaches
- X-ray crystallography: Difficult to crystallize weak complexes with unstructured regions
 - NMR spectroscopy: Size limits
 - Electron microscopy: Trapping of functional assemblies
- ⇒ New opportunities for modeling, simulation, and theory!
- Complement experiments
 - Provide predictions, insights, and new directions


Outline

1. Model and Method:

- Validation: structures and binding affinities


2. Structure and dynamics of multi-protein assemblies

- Vps27/Hse1: ESCRT protein sorting machinery
- Collaboration with *James H. Hurley*, NIDDK





3. Transient encounter complexes in protein-protein complex formation

- Paramagnetic relaxation enhancement NMR of protein-protein complexes
- Collaboration with *G. Marius Clore*, NIDDK



Coarse-grained model for multi-protein assemblies

- **Residue-level (C_α only) coarse-graining**
 - Rigid body for structured domains
- **Transferable energy function**
 - Long-range Debye-Hückel electrostatic interactions
 - Residue-dependent short-range interactions (Miyazawa-Jernigan statistical contact potentials)
 - Experimental inputs: Lysozyme *osmotic protein second-virial coefficient* and Ub-CUE *protein binding affinity*
- **Flexible linkers: polymer model**
 - Harmonic stretching potential
 - Bending potential
 - Torsion angle potential
- **Membrane interactions**
 - Planar membrane
 - Short-range interactions between residues and membrane
 - Electrostatic interactions



Simulation method

- Replica exchange Monte Carlo
 - Twenty replicas at different temperatures
 - Enhances equilibrium sampling
 - Implemented in the parallel architecture of Biowulf cluster




Validation: complex structure

Ub/CUE
Ub/UIM1
Ub/UBA
Ub/GAT
Ub/DUIM
Ub/UIM2
UbL/UBA
UbL/UIM
UbL/UIM2




Distance to native structure ↑

Energy →




$$\text{DRMS} = \frac{1}{N} \sum_{i,j} \left| d_{ij}^{\text{exp}} - d_{ij}^{\text{model}} \right|$$



Native-like ↓



Native-like ↓




Native-like ↓

(YCK, Hummer; *J. Mol. Biol.* **375**, 1416, 2008)

Validation: binding affinities

Ub/CUE
Ub/UIIM1
Ub/UBA
Ub/GAT
Ub/DUIM
Ub/UIIM2
UbL/UBA
UbL/UIIM
UbL/UIIM2


Experiments



Application I: multi-vesicular body (MVB) protein sorting machinery


- The ESCRT machinery targets ubiquitinated transmembrane proteins for degradation in the lysosome or yeast vacuole
- ESCRTs are required for HIV budding at the plasma membrane

Vps27/Hse1(yeast)
Hrs/STAM(human)




Structure of the assembled Vps27/Hse1 complex



Vps27/Hse1 complex





Vps27/Hse1 complex is dynamic and open



Hrs/STAM(human) complex also shows open structures



Positive cooperativity enhances Vps27 binding to ubiquitin



FYVE domain is tethered to the membrane


Proteins can move freely parallel to the membrane

Summary I: Simulations of Vps27/Hse1

- Dynamic and open structure
 - Important for targeting a variety of ubiquitinated cargos
- Cooperative binding of ubiquitin via nonspecific interactions
 - Essential for function at low biological concentrations
- Are nonspecific interactions detectable?

Application II: Transient encounter complexes probed by simulation and NMR

- Paramagnetic relaxation enhancement (PRE) probes the presence of low-population (<10%) transient encounter complexes




$$\text{PRE} \sim 1/r^6$$

Iwahara, Clore,
Nature **440**,
1227, 2006

NMR Paramagnetic Relaxation Enhancement (encounter complexes of HPr-IIA^{Mannose})

- PRE of backbone amide protons on IIA^{Mannose}





Can we simulate encounter complexes?

(Tang, Iwahara, Clore, *Nature* **444**, 383, 2006)

Replica-exchange simulations of HPr-IIA^{Man} complex


- Coarse-grained simulation model (YCK, Hummer; *J. Mol. Biol.* **375**, 1416, 2008)



(YCK, Tang, Clore, Hummer, *PNAS USA* **105**, 12855, 2008)

PRE profiles of HPr-IIA^{Man} complex

- PRE of backbone amide protons on IIA^{Mannose}




Reweighting of simulation structures

- Simulation model should not be expected to produce accurate populations
- 2 kT binding free energy difference
→ 10-fold difference in population


⇒ Cluster the structures of the specific and non-specific complexes

⇒ Re-weight the populations of the clusters to match PRE profiles




PRE profiles of HPr-IIA^{Man} complex

- PRE of backbone amide protons on IIA^{Mannose}



Energy landscape of protein complex formation

- Funnel-like energy landscape



Summary II: Biology of transient encounter complexes

- **Accelerated on-rate** (barnase: Schreiber, Fersht, *Nat Struct Biol* **3**, 427, 1996)
- **Strengthening of weak specific interactions in multi-protein assemblies (Vps27)**
- **Alternative binding modes** (mannose transport: Hu *et al.* *J Biol Chem* **283**, 11024, 2008)
- **Evolutionary remnants of earlier specific interactions?**

Conclusion

- Coarse-grained model and transferable energy function provide valuable and complementary information regarding structures and dynamics of *multi-protein assemblies* and *transient encounter complexes*

Acknowledgments

- Gerhard Hummer (NIDDK, NIH)

ESCRT complex

- James Hurley (NIDDK, NIH)

PRE of encounter complexes

- G. Marius Clore (NIDDK, NIH)

- Chun Tang (U. Missouri)

Computational resources

- NIH Biowulf
- Helix Systems Staff