



# CALCULUS

## EARLY TRANSCENDENTAL FUNCTIONS

5th EDITION

**ROBERT T. SMITH, ROLAND B. MINTON, ZIAD A. T. RAFHI**

**Mc  
Graw  
Hill**  
Education

© 2017 by McGraw-Hill Education. Permission required for reproduction or display.

## **Methods of Integration**

### **1- Integration by Parts**

### **2- Integration by substitution**

i) **Algebraic substitutions**

ii) **Trigonometric substitutions**

### **3- Integration by partial fractions**

$$\int \frac{p(x)}{q(x)} dx ,$$

*p(x) and q(x) are polynomials*

**The degree of the numerator is less than the degree of the denominator**

*a) The denominator consists of linear factors*

*b) The denominator has repeated linear factors*

*c) A factor of second degree that does not decompose*

**Example** 
$$\int \frac{2x^2 - 5x + 2}{x^3 + x} dx$$

$$\frac{2x^2 - 5x + 2}{x^3 + x} =$$

$$= \frac{2x^2 - 5x + 2}{x(x^2 + 1)} = \frac{A}{x} + \frac{Bx + C}{x^2 + 1}$$

$$\rightarrow 2x^2 - 5x + 2 = A(x^2 + 1) + (Bx + C)x$$

$$\rightarrow 2x^2 - 5x + 2 = Ax^2 + A + Bx^2 + Cx$$

$$\rightarrow 2x^2 - 5x + 2 = (A + B)x^2 + Cx + A$$

$$\rightarrow A = 2$$

$$\rightarrow C = -5$$

$$\rightarrow A + B = 2$$

$$\rightarrow B = 2 - A = 2 - 2 \rightarrow B = 0$$

$$\rightarrow \frac{2x^2 - 5x + 2}{x(x^2 + 1)} = \frac{2}{x} + \frac{-5}{x^2 + 1}$$

$$\rightarrow \int \frac{2x^2 - 5x + 2}{x(x^2 + 1)} dx =$$

$$= \int \frac{2}{x} dx + \int \frac{-5}{1 + x^2} dx$$

$$= 2 \int \frac{1}{x} dx - 5 \int \frac{1}{1 + x^2} dx$$

$$= 2 \ln|x| - 5 \tan^{-1}x + c$$

**The degree of the numerator is more than or equal to the degree of the denominator**

**Example**  $\int \frac{2x^3 - 4x^2 - 15x + 5}{x^2 - 2x - 8} dx$

$$\begin{array}{r} 2x \\ x^2 - 2x - 8 \overline{) 2x^3 - 4x^2 - 15x + 5} \\ \underline{2x^3 - 4x^2 - 16x} \phantom{+ 5} \\ x + 5 \end{array}$$

$$\frac{2x^3 - 4x^2 - 15x + 5}{x^2 - 2x - 8} = 2x + \frac{x + 5}{x^2 - 2x - 8}$$

$$\frac{x + 5}{x^2 - 2x - 8} =$$

$$= \frac{x + 5}{(x + 2)(x - 4)} = \frac{A}{x + 2} + \frac{B}{x - 4}$$

$$\rightarrow x + 5 = A(x - 4) + B(x + 2)$$

$$\text{at } x = -2 \rightarrow 3 = -6A \rightarrow A = -\frac{1}{2}$$

$$\text{at } x = 4 \rightarrow 9 = 6B \rightarrow B = \frac{3}{2}$$

$$\rightarrow \frac{x+5}{x^2-2x-8} = \frac{-\frac{1}{2}}{x+2} + \frac{\frac{3}{2}}{x-4}$$

$$\rightarrow \frac{2x^3 - 4x^2 - 15x + 5}{x^2 - 2x - 8} =$$

$$= 2x + \frac{-\frac{1}{2}}{x+2} + \frac{\frac{3}{2}}{x-4}$$

$$\rightarrow \int \frac{2x^3 - 4x^2 - 15x + 5}{x^2 - 2x - 8} dx =$$

$$= 2 \int x dx - \frac{1}{2} \int \frac{1}{x+2} dx + \frac{3}{2} \int \frac{1}{x-4} dx$$

$$= x^2 - \frac{1}{2} \ln|x+2| + \frac{3}{2} \ln|x-4| + c$$