

Particle Physics II
Lecture 12: The Higgs boson

Lesya Shchutska

June 1, 2023

Fermion masses

- Higgs mechanism gives masses not only to W and Z bosons but also to fermions
- due to different transformation of LH and RH chiral states the mass term of the Dirac lagrangian

$$-m\bar{\psi}\psi = -m(\bar{\psi}_R\psi_L + \bar{\psi}_L\psi_R)$$

is not symmetric under $SU(2)_L \times U(1)_Y$ gauge symmetry

\implies cannot be in the SM lagrangian

- therefore the fermion fields are introduced to the SM as massless states and they acquire their mass thanks to the interaction with Higgs field

Transformations of left doublet

- in the SM, LH chiral fermions are placed in SU(2) doublets (L)
- RH chiral fermions – in SU(2) singlets (R)
- since two complex scalar Higgs fields are in an SU(2) doublet $\phi(x)$, a local gauge transformation affects it as:

$$\phi \rightarrow \phi' = (I + ig_W \vec{\epsilon}(x) \cdot \vec{T})\phi$$

- the same local gauge transformation applies to the LH fermion doublet L
- for $\bar{L} \equiv L^\dagger \gamma^0$:

$$\bar{L} \rightarrow \bar{L}' = \bar{L}(I - ig_W \vec{\epsilon}(x) \cdot \vec{T})$$

- $\implies \bar{L}\phi$ is invariant under SU(2)_L gauge transformations

Invariant form

- combined with a RH singlet: $\bar{L}\phi R \implies$ invariant under $SU(2)_L$ and $U(1)_Y$ gauge transformations
- its Hermitian conjugate is invariant too:

$$(\bar{L}\phi R)^\dagger = \bar{R}\phi^\dagger L$$

- \implies such term satisfies the $SU(2)_L \times U(1)_Y$ gauge symmetry:

$$-g_f (\bar{L}\phi R + \bar{R}\phi^\dagger L)$$

Electron case

- lagrangian part for electrons would look like:

$$\mathcal{L}_e = -g_e \left[(\bar{\nu}_e \quad \bar{e})_L \begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix} e_R + \bar{e}_R (\phi^{+*} \quad \phi^{0*}) \begin{pmatrix} \nu_e \\ e \end{pmatrix}_L \right]$$

- g_e is a constant – Yukawa coupling of the electron to the Higgs field
- after spontaneous symmetry breaking the Higgs doublet in the unitary gauge is:

$$\phi(x) = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v + h(x) \end{pmatrix}$$

- electron term in the lagrangian becomes:

$$\mathcal{L}_e = -\frac{g_e}{\sqrt{2}} v (\bar{e}_L e_R + \bar{e}_R e_L) - \frac{g_e}{\sqrt{2}} h (\bar{e}_L e_R + \bar{e}_R e_L)$$

- first term there has exactly the form needed for the fermion masses

Electron case


- the Yukawa coupling g_e is not predicted by the Higgs mechanism but can be chosen to be consistent with the observed electron mass:

$$g_e = \sqrt{2} \frac{m_e}{v}$$

- then electron mass lagrangian becomes:

$$\mathcal{L}_e = -m_e \bar{e}e - \frac{m_e}{v} \bar{e}e h$$

- first term (giving the mass to electron) – is a coupling of electron to the Higgs field through its non-zero v
- second term – is a coupling between the electron and the Higgs boson



Up-type fermions

- term $\bar{L}\phi R + \bar{R}\phi^\dagger L$ can generate masses only for down-type fermions:
 - this happens since non-zero vacuum expectation value v exists only in the lower (neutral) component of the Higgs doublet
- to have masses for up-type fermions, use the conjugate doublet ϕ_c formed as:

$$\phi_c = -i\sigma_2\phi^* = \begin{pmatrix} -\phi^{0*} \\ \phi^- \end{pmatrix} = \frac{1}{\sqrt{2}} \begin{pmatrix} -\phi_3 + i\phi_4 \\ \phi_1 - i\phi_2 \end{pmatrix}$$

- the conjugate doublet ϕ_c transforms in the same way as ϕ under SU(2)
- this is similar to the representation of up- and down-quarks and anti-up and anti-down quarks in SU(2) isospin symmetry

Up-type fermions

- a gauge-invariant mass term for up-type quarks comes from the term:

$$\bar{L}\phi_c R + \bar{R}\phi_c^\dagger L$$

- as an example, for u quark:

$$\mathcal{L}_u = g_u (\bar{u} \quad \bar{d})_L \begin{pmatrix} -\phi^{0*} \\ \phi^- \end{pmatrix} u_R + H.C.$$

- after symmetry breaking it becomes:

$$\mathcal{L}_u = -\frac{g_u}{\sqrt{2}}v(\bar{u}_L u_R + \bar{u}_R u_L) - \frac{g_u}{\sqrt{2}}h(\bar{u}_L u_R + \bar{u}_R u_L)$$

- with Yukawa coupling $g_u = \sqrt{2}m_u/v$ leading to:

$$\mathcal{L}_u = -m_u \bar{u}u - \frac{m_u}{v} \bar{u}u h$$

Fermion masses

- so for all Dirac fermions gauge invariant mass terms can be constructed from:

$$\mathcal{L} = -g_f \left[\bar{L}\phi R + (\bar{L}\phi R)^\dagger \right] \text{ or } \mathcal{L} = g_f \left[\bar{L}\phi_c R + (\bar{L}\phi_c R)^\dagger \right]$$

- these terms produce masses of the fermions and the interactions between the Higgs boson and the fermion
- the Yukawa couplings of the fermions to the Higgs field are given by:

$$g_f = \sqrt{2} \frac{m_f}{v},$$

where $v = 246 \text{ GeV}$




- e.g. for the top quark with $m_t \approx 173.1 \pm 0.9 \text{ GeV}$, the Yukawa coupling is almost exactly 1, resonating with a notion that maybe Yukawa couplings of the fermions should be $\mathcal{O}(1)$
- a special case is neutrinos where Yukawa couplings are $< 10^{-12}$: other mechanisms are suggested for neutrino masses

Properties of the Higgs boson

- the SM Higgs boson H is a neutral scalar particle
- its mass is a free parameter of the SM $m_H = 2\lambda v^2$
- H couples to all fermions with a coupling strength proportional to the fermion mass, vertex factor is:

$$-i\frac{m_f}{v} \equiv -i\frac{m_f}{2m_W}g_W$$

- H decays as $H \rightarrow f\bar{f}$ to all kinematically allowed decay modes $m_H > 2m_f$
- for vector bosons, coupling is also proportional to boson mass:





Higgs boson decays

- for a discovered H boson with $m = 125$ GeV the largest BF is to $b\bar{b}$:

$$\mathcal{B}(H \rightarrow b\bar{b}) = 57.8\%$$

- we can compute partial decay width:
 - vertex is known
 - H is scalar \implies no polarization 4-vector is needed
 - matrix element is simply:

$$\mathcal{M} = \frac{m_b}{v} \bar{u}(p_2) v(p_3) = \frac{m_b}{v} u^\dagger \gamma^0 v$$



Higgs boson decays

- can take b -quark momentum along the z -axis
- $m_H \gg m_b$: can neglect b mass and take

$$p_2 \approx (E, 0, 0, E) \text{ and } p_3 \approx (E, 0, 0, -E), \text{ where } E = m_H/2$$

- in the ultra-relativistic limit the spinors for two possible helicity state for each of the b ($\theta = 0, \phi = 0$) and the \bar{b} ($\theta = \phi, \phi = \pi$) are:

$$u_{\uparrow}(p_2) = \sqrt{E} \begin{pmatrix} 1 \\ 0 \\ 1 \\ 0 \end{pmatrix}, u_{\downarrow}(p_2) = \sqrt{E} \begin{pmatrix} 0 \\ 1 \\ 0 \\ -1 \end{pmatrix},$$

$$v_{\uparrow}(p_3) = \sqrt{E} \begin{pmatrix} 1 \\ 0 \\ -1 \\ 0 \end{pmatrix}, v_{\downarrow}(p_3) = \sqrt{E} \begin{pmatrix} 0 \\ -1 \\ 0 \\ -1 \end{pmatrix}$$

Higgs boson decays

- due to $u^\dagger \gamma^0 v$ form of the ME only two of the possible four helicity combinations give non-0 matrix element:

$$\mathcal{M}_{\uparrow\uparrow} = -\mathcal{M}_{\downarrow\downarrow} = \frac{m_b}{v} 2E$$

- these correspond to spin configurations where $b\bar{b}$ are produced in a spin-0 state
- since H is a spin-0 scalar, it decays isotropically and ME does not have angular dependence
- and since there is only one spin state for H, the spin-averaged ME is:

$$\langle |\mathcal{M}|^2 \rangle = |\mathcal{M}_{\uparrow\uparrow}|^2 + |\mathcal{M}_{\downarrow\downarrow}|^2 = \frac{m_b^2}{v^2} 8E^2 = \frac{2m_b^2 m_H^2}{v^2}$$

- the partial decay width:

$$\Gamma(H \rightarrow b\bar{b}) = 3 \times \frac{m_b^2 m_H}{8\pi v^2},$$

where factor 3 accounts for three colors of $b\bar{b}$ pair

- for SM H: $\Gamma(H \rightarrow b\bar{b}) \approx 2 \text{ MeV}$




Higgs boson decays

- partial H decay rate to fermions is $\propto m_f^2$:

$$\Gamma(H \rightarrow b\bar{b}) : \Gamma(H \rightarrow c\bar{c}) : \Gamma(H \rightarrow \tau^+\tau^-) \approx 3m_b^2 : 3m_c^2 : m_\tau^2$$

- m_q run with q^2 similarly to α_S : masses should be taken at $q^2 = m_H^2$?
- there $m_c(m_H^2) \approx 0.6$ GeV and $m_b(m_H^2) \approx 3.0$ GeV
- decay to massless particles γ and g happen via loops:

Decay mode	Branching ratio
$H \rightarrow b\bar{b}$	57.8%
$H \rightarrow WW^*$	21.6%
$H \rightarrow \tau^+\tau^-$	6.4%
$H \rightarrow gg$	8.6%
$H \rightarrow c\bar{c}$	2.9%
$H \rightarrow ZZ^*$	2.7%
$H \rightarrow \gamma\gamma$	0.2%



Higgs boson decays measurements

- example of measurements by the ATLAS experiment, testing H couplings to all particles:

