

THE CHINESE UNIVERSITY OF HONG KONG  
Department of Mathematics  
MATH3280 Introductory Probability 2022-2023 Term 1  
Suggested Solutions of Homework Assignment 3

**Q1**

(a). The image of  $X$  is  $\{4, 2, 1, 0, -1, -2\}$ .

$$P(X = 4) = \frac{\binom{4}{2}}{\binom{14}{2}} = \frac{6}{91}$$

$$P(X = 2) = \frac{\binom{4}{1} \binom{2}{1}}{\binom{14}{2}} = \frac{8}{91}$$

$$P(X = 1) = \frac{\binom{4}{1} \binom{8}{1}}{\binom{14}{2}} = \frac{32}{91}$$

$$P(X = 0) = \frac{\binom{2}{2}}{\binom{14}{2}} = \frac{1}{91}$$

$$P(X = -1) = \frac{\binom{8}{1} \binom{2}{1}}{\binom{14}{2}} = \frac{16}{91}$$

$$P(X = -2) = \frac{\binom{8}{2}}{\binom{14}{2}} = \frac{4}{13}$$

(b).

$$E(X) = \sum_{x:p(x)>0} x \cdot p(x) = 4 \cdot \frac{6}{91} + 2 \cdot \frac{8}{91} + 1 \cdot \frac{32}{91} + 0 \cdot \frac{1}{91} - 1 \cdot \frac{16}{91} - 2 \cdot \frac{4}{13} = 0$$

The expected value of the money we are going to get for playing 100 games is  $(0 - 2) \times 100 = -200$  dollars.

(c). It is not fair. The game is biased against us.

## Q2

The possible values of  $X$  are 1, 2, 3, 4, 5, 6 and the probabilities that  $X$  takes on each of these values are

$$P(X = 1) = \frac{5 \cdot 9!}{10!} = \frac{1}{2}$$

$$P(X = 2) = \frac{5 \cdot 5 \cdot 8!}{10!} = \frac{5}{18}$$

$$P(X = 3) = \frac{5 \cdot 4 \cdot 5 \cdot 7!}{10!} = \frac{5}{36}$$

$$P(X = 4) = \frac{5 \cdot 4 \cdot 3 \cdot 5 \cdot 6!}{10!} = \frac{5}{84}$$

$$P(X = 5) = \frac{5 \cdot 4 \cdot 3 \cdot 2 \cdot 5 \cdot 5!}{10!} = \frac{5}{252}$$

$$P(X = 6) = \frac{5!5!}{10!} = \frac{1}{252}$$

$$P(X = 7) = P(X = 8) = P(X = 9) = P(X = 10) = 0$$

### Q3

First, find the mass probability functions of  $X$  and  $Y$ . Let  $\Omega = \{40, 33, 25, 50\}$ .

$$P(X = 40) = \frac{40}{148}$$

$$P(X = 33) = \frac{33}{148}$$

$$P(X = 25) = \frac{25}{148}$$

$$P(X = 50) = \frac{50}{148}$$

$$P(Y = i) = \frac{1}{4}, \quad i \in \Omega$$

Then we are going to calculate the expectation and variance of  $X$  and  $Y$ .

$$E(X) = \sum_{k \in \Omega} kP(X = k) = 40 \cdot \frac{40}{148} + 33 \cdot \frac{33}{148} + 25 \cdot \frac{25}{148} + 50 \cdot \frac{50}{148} \approx 39.28$$

$$E(Y) = \sum_{k \in \Omega} kP(Y = k) = 40 \cdot \frac{1}{4} + 33 \cdot \frac{1}{4} + 25 \cdot \frac{1}{4} + 50 \cdot \frac{1}{4} = 37$$

$$E(X^2) = \sum_{k \in \Omega} k^2P(X = k) = 40^2 \cdot \frac{40}{148} + 33^2 \cdot \frac{33}{148} + 25^2 \cdot \frac{25}{148} + 50^2 \cdot \frac{50}{148}$$

$$\text{Var}(X) = E(X^2) - E(X)^2 \approx 82.20$$

$$E(Y^2) = \sum_{k \in \Omega} k^2P(Y = k) = 40^2 \cdot \frac{1}{4} + 33^2 \cdot \frac{1}{4} + 25^2 \cdot \frac{1}{4} + 50^2 \cdot \frac{1}{4}$$

$$\text{Var}(Y) = E(Y^2) - E(Y)^2 = 84.5$$

## Q4

(a). By simple observation, we can see the event  $\{X = 1\}$  is equal to  $\{(H, T), (T, H)\}$ .

$$P(X = 1) = P\{(H, T), (T, H)\} = 0.6 \times (1 - 0.7) + (1 - 0.6) \times 0.7 = 0.46$$

(b). Similarly,  $\{X = 2\} = \{(H, H)\}$ . Then

$$\begin{aligned} E(X) &= \sum_{k=0}^2 kP(X = k) \\ &= 0 \cdot P(X = 0) + 1 \cdot P(X = 1) + 2 \cdot P(X = 2) \\ &= 1 \cdot 0.46 + 2 \cdot (0.6 \times 0.7) \\ &= 1.3 \end{aligned}$$

## Q5

(a). Denote the event “first coin is flipped” by  $C_1$ , with  $C_2$  defined similarly. Let  $X$  be the number of heads out of 10 tosses.

$$\begin{aligned} P(X = 7) &= P(X = 7 | C_1) P(C_1) + P(X = 7 | C_2) P(C_2) \\ &= \left[ \binom{10}{7} \cdot 4^7 \cdot 6^3 \right] \frac{1}{2} + \left[ \binom{10}{7} \cdot 7^7 \cdot 3^3 \right] \frac{1}{2} \\ &= [.0425] \cdot 5 + [.2668] \cdot 5 \\ &= .1547 \end{aligned}$$

(b). By conditioning on the outcome of the first flip, we update the probability (now evenly split between coins 1 and 2) that coin 1 is being flipped. Let  $H_1$  denote the event “the first flip is heads”.

$$\begin{aligned} P(C_1 | H_1) &= \frac{P(H_1 | C_1) P(C_1)}{P(H_1 | C_1) P(C_1) + P(H_1 | C_2) P(C_2)} \\ &= \frac{.4 \times .5}{.4 \times .5 + .7 \times .5} \\ &= \frac{4}{11} \end{aligned}$$

Our updated probabilities are now:  $P(C_1 | H_1) = \frac{4}{11}$  and  $P(C_2 | H_1) = \frac{7}{11}$ .  
Then

$$\begin{aligned}
& P(X = 7 | H_1) \\
&= \frac{P(\{X = 7\} \cap H_1)}{P(H_1)} \\
&= \frac{P(\{X = 7\} \cap C_1 \cap H_1) + P(\{X = 7\} \cap C_2 \cap H_1)}{P(H_1)} \\
&= \frac{P(\{X = 7\} \cap C_1 \cap H_1)}{P(C_1 \cap H_1)} \cdot \frac{P(C_1 \cap H_1)}{P(H_1)} + \frac{P(\{X = 7\} \cap C_2 \cap H_1)}{P(C_2 \cap H_1)} \cdot \frac{P(C_2 \cap H_1)}{P(H_1)} \\
&= P(X = 7 | C_1 \cap H_1) \cdot P(C_1 | H_1) + P(X = 7 | C_2 \cap H_1) \cdot P(C_2 | H_1) \\
&= \left[ \binom{9}{6} \cdot 4^6 \cdot 6^3 \right] \frac{4}{11} + \left[ \binom{9}{6} \cdot 7^6 \cdot 3^3 \right] \frac{7}{11} \\
&= [.0743].3636 + [.2668].6364 \\
&= .1968
\end{aligned}$$

## Q6

The number of potential interviewees who consent to the interview  $X$  is a binomial random variable, with  $n = 5$  and  $p = \frac{2}{3}$ . Q: What is the probability that each of the 5 people consents to the interview?

(a).  $P(X = 5) = \binom{5}{5} \left(\frac{2}{3}\right)^5 \left(\frac{1}{3}\right)^0 = \left(\frac{2}{3}\right)^5 = \frac{32}{243} = .1317$

(b). Now  $X \sim \text{Binom}\left(8, \frac{2}{3}\right)$

$$\begin{aligned} P(X \geq 5) &= \sum_{k=5}^8 \binom{8}{k} \left(\frac{2}{3}\right)^k \left(\frac{1}{3}\right)^{8-k} \\ &= .7414 \end{aligned}$$

(c). We are asked for the probability that the 6th potential interviewee will be the 5th to consent.

$$\begin{aligned} P(X = 6) &= \binom{5}{4} \left(\frac{2}{3}\right)^5 \frac{1}{3} \\ &= \frac{160}{729} = .2195 \end{aligned}$$

(d).

$$\begin{aligned} P(X = 7) &= \binom{6}{4} \left(\frac{2}{3}\right)^5 \left(\frac{1}{3}\right)^2 \\ &= \frac{160}{729} = .2195 \end{aligned}$$

## Q7

As the expected value of a function of a discrete random variable  $X$  equals  $\sum_i f(x_i) P\{X = x_i\}$ ,  $c^X$  has as an expected value of

$$\begin{aligned} E[c^X] &= c^1 P\{X = 1\} + c^{-1} P\{X = -1\} \\ E[c^X] &= c(p) + \left(\frac{1}{c}\right)(1-p) \\ E[c^X] &= cp + \frac{1-p}{c} \end{aligned}$$

Setting this equal to 1,

$$\begin{aligned}cp + \frac{1-p}{c} &= 1 \\pc^2 + 1 - p &= c \\pc^2 - c + 1 - p &= 0\end{aligned}$$

If  $p = 0$ , then  $c = 1$  (disregarded). If  $p \neq 0$ , we solve this quadratic equation and get

$$\begin{aligned}c &= \frac{1 \pm \sqrt{(-1)^2 - 4p(1-p)}}{2p} \\c &= \frac{1 \pm \sqrt{(2p-1)^2}}{2p} \\c &= \frac{1 \pm (2p-1)}{2p}\end{aligned}$$

Solving the equation above, we get  $c = \frac{1}{p} - 1$  or  $c = 1$  (disregarded).

**Q8**

$$\begin{aligned}E\left(\frac{1}{X+1}\right) &= \sum_{k=0}^n \frac{1}{k+1} P(X=k) \\&= \sum_{k=0}^n \frac{1}{k+1} \frac{n!}{k!(n-k)!} p^k (1-p)^{n-k} \\&= \frac{1}{(n+1)p} \sum_{k=0}^n \frac{(n+1)!}{(k+1)!(n-k)!} p^{k+1} (1-p)^{n-k} \\&= \frac{1}{(n+1)p} \left( \sum_{i=0}^{n+1} \frac{(n+1)!}{i!(n+1-i)!} p^i (1-p)^{n+1-i} - (1-p)^{n+1} \right) \\&= \frac{1}{(n+1)p} ((p + (1-p))^{n+1} - (1-p)^{n+1}) \\&= \frac{1 - (1-p)^{n+1}}{(n+1)p}\end{aligned}$$

## Q9

Note that a Poisson r.v. has the parameter  $\lambda > 0$  and  $P(X = k) = \frac{\lambda^k e^{-\lambda}}{k!}$  for  $k = 0, 1, 2, \dots$ . Fix a  $k \in \{0, 1, 2, \dots\}$  and let  $f_k(\lambda) = \frac{\lambda^k e^{-\lambda}}{k!}$ . If  $k = 0$ , note that  $f_0(\lambda) = e^{-\lambda}$ ,  $\lambda > 0$ , so no maximum is attained for  $k = 0$ . If  $k > 0$ , then

$$f'_k(\lambda) = \frac{e^{-\lambda}}{k!} (k\lambda^{k-1} - \lambda^k)$$
$$\begin{cases} > 0 & \text{if } \lambda < k \\ = 0 & \text{if } \lambda = k \\ < 0 & \text{if } \lambda > k \end{cases}$$

Hence  $\lambda = k$  maximizes  $P(X = k)$ .

## Q10

First, according to the expectation of a function of random variables, we have

$$\begin{aligned} E(X^n) &= \sum_{k=1}^{\infty} k^n \frac{\lambda^k e^{-\lambda}}{k!} \\ &= \lambda \sum_{k=1}^{\infty} k^{n-1} \frac{\lambda^{k-1} e^{-\lambda}}{(k-1)!} \\ &= \lambda \sum_{i=0}^{\infty} (i+1)^{n-1} \frac{\lambda^i e^{-\lambda}}{i!} \\ &= \lambda E((X+1)^{n-1}). \end{aligned}$$

We have known  $E(X) = \lambda$ . Then by the formula proved above,

$$\begin{aligned} E(X^2) &= \lambda E(X+1) = \lambda(E(X) + 1) = \lambda^2 + \lambda \\ E(X^3) &= \lambda E((X+1)^2) \\ &= \lambda(E(X^2 + 2X + 1)) \\ &= \lambda(E(X^2) + 2E(X) + 1) \\ &= \lambda(\lambda^2 + \lambda + 2\lambda + 1) \\ &= \lambda^3 + 3\lambda^2 + \lambda. \end{aligned}$$