

MY ARTICLE

ZHOU FENG

1. INTRODUCTION

A good introduction to fractal geometry is Falconer [3]. There is `smallmatrix` environment (e.g,  $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$ ). It is recommended to use  $\dots, \cdots, \cdots, \cdots, \dots$  instead of  $\dots$  and  $\cdots$ . Then we test the `\nobreakdash`:  $p$ -adic, page 1–9,  $n$ -dim, 1-dim,  $\sigma$ -algebra. What about a text-mode fractional:  $\frac{\log_k H}{1212}$ .

Then for the `\xleftarrow`:

(1.1)

$$A \xleftarrow{n+\mu-1} B \xrightarrow[T]{n\pm i-1\mathrm{bla}, \mathrm{bla}, \mathrm{bla}} C \bigcap_{i\geq 1} A_i \bigcup_{k=1}^{100} \mathfrak{r}_k$$

$$\Leftrightarrow$$

Compare the `\choose` and `\binom`:  $\binom{n}{k}\binom{n}{k}$ .  $|z|, \|v\|, \|v\|_\infty$ .

About the user-defined math operators:

$$\mathrm{ex}(\mathrm{conv}(A_i))\mathrm{abc}\lim_{x\rightarrow 0\atop n\rightarrow\infty}$$

Then the `\mod`:  $\gcd(n, m \bmod n); \quad x \equiv y \pmod b, x \equiv y \bmod c, x \equiv y \pmod d$ .

See the following default math environments:

(1.2)

$$\vec{F} = m\vec{a}$$

$$\vec{F} = G\frac{m_1m_2}{r^2}$$

(1.3a)

$$\nabla \cdot \vec{E} = \varepsilon_0 \rho$$

(1.3b)

$$\nabla \cdot \vec{B} = 0$$

(1.3c)

$$\nabla \times \vec{E} = -\frac{\partial \vec{B}}{\partial t}$$

(1.3d)

$$\nabla \times \vec{B} = \mu_0 \varepsilon_0 \vec{J} + \frac{\partial \vec{E}}{\partial t}$$

(1.4)

$$E = \gamma mc^2$$
$$\mathcal{R}_{\mu\nu} - \frac{\mathcal{R}}{2} g_{\mu\nu} + \Lambda g_{\mu\nu} = \frac{8\pi G}{c^4} T_{\mu\nu}$$

• `\substack{}` and `\begin{subarray}`

(1.5)

$$\mathop{\mathrm{Lim}}_{\substack{0\leq i\leq m\\0\leq j\leq n}}P(i,j)$$

(1.6)

$$\sum_{\substack{i\in\Lambda\\0\leq j\leq n}}P(i,j)$$

♡ `\sideset{text}{right}{symbol}`

$$a_0+\frac{1}{a_1+\frac{1}{a_2+\frac{1}{a_3+\cdots}}}$$

(1.7)

$$*\prod_{n=1}^{\infty}*\left[*\prod_{n=1}^{*\infty}\sum_{\text{ex}}\frac{\prod_{n=1}^{\infty}}{\lim_{n\rightarrow\infty}}\right]\Bigg\}\lim_{n\rightarrow\infty}\text{Quantum Computing}$$

The `\mathbf{f}` command is commonly used to obtain bold Latin letters in math, but for most other kinds of math symbols it has no effect.

`\mid` and `\mathbin{}` : $P(A \mid B)P(A \mid B)P(A|B)$

- (a) `f : X\to Y` vs. `f\colon X\to Y`:  $f : X \rightarrow Y$  vs.  $f\colon X \rightarrow Y$ .
- (b) `:=` vs. `\coloneqq` :  $:=$  vs.  $\coloneqq$ .
- (c)  $\{z : z \in \mathbb{Z}\}$  vs.  $\{z\colon z \in \mathbb{Z}\}$ .
- (d)  $v_1, v_2, \ldots, v_n$  vs.  $v_1, \ldots, v_n$ .
- (e)  $f(n) = O(n)$  vs.  $f(n)$  is  $O(n)$  or  $f(n) \in O(n)$ .
- (f)  $A \setminus B$  vs.  $A\backslash B$  vs.  $A - B$ .

- (g) There is a `\`, spacing between integrand and measure

$$\int_a^b x^2 dx$$

- (h) Use `Serre et al.``\ proved`: Serre et al. proved.  
`Serre et al.` proved: Serre et al. proved.

Here is some practical suggestions for mathematical writting.

- (1) The structures for conditional sentences: `If ... , then...`; `When...`, ...; `For ... , ....` No `Let....` `Then...`!
- (2) Avoid using `as` and `for` to introduce reasons after some conclusion.
- (3) `Hence`, `Thus`, and `Therefore`, .
- (4) , `so` is informal and should be used when the conclusion is short.
- (5) A statement that is **assumed** is an axiom, and throughout to be true. Something **supposed** is a hypothesis and more appropriate to introduce a case or an argument by contradiction. For example, `Suppose to the contrary that` and `Toward a contradiction, suppose that`.
- (6) No `v's` or `ai's`.
- (7) No *nested* proof environments.
- (8) `We induct on n` vs. `We use induction on n`.
- (9) Prefer `pairwise` to `mutually`.
- (10) No contractions like `can't`, `won't`, etc.
- (11) Use `\begingroup\allowdisplaybreaks ... \endgroup` to allow the large chunk of math display environments to be broken into pages.
- (12) Replace `$$ ... $$` with `\[...\]` in `sed`:  
`sed '/\$\$/{:x;N;/.*\$\$ *$/!bx;s/\$\$(.*)\$\$ */\[1\]/}'`

## 2. COMMUTATIVE DIAGRAMS

Arrows `@>>>` `@<<<` `@VVV` `@AAA`. Double lines: `@=`. Null arrows: `@`

$$(2.1) \quad \begin{array}{ccc} S^{\mathcal{W}_\Lambda \otimes T} & \xrightarrow{j} & T \\ \parallel & & \downarrow_{\text{End } P} \\ (S \otimes T)/I & \equiv & (Z \otimes T)/J \end{array}$$

`tikzcd` is the ultimate answer to a commutative diagram in  $\text{T}_\text{E}\text{X}$ .

$$\begin{array}{ccc} A & \xrightarrow{\phi} & B \\ & \searrow & \\ & & C \end{array}$$

### 3. REFERENCE & CITATION

Choose a `natbib` compatible `\bibliographystyle`, e.g. `abbrvnat`, `plainnat`.

- `\cite{}`: [1]
- `\citet{}`: Akiyama et al. [1]
- `\citet*{}`: Akiyama, Feng, Kempton, and Persson [1]
- `\citep{}`: [1]
- `\citep*{}`: [1]
- `\citealt*{}`: Akiyama, Feng, Kempton, and Persson 1
- `\citeyear{}`: 2020
- `\citeauthor{}`: Akiyama et al.
- `\citeauthor*{}`: Akiyama, Feng, Kempton, and Persson
- `\cite[text]{keylist}` [1, Theorem 1]
- `\cite[prefix][suffix]{keylist}`: [see e.g. 1, p. 123]
- `\citenum{}`: 1
- `\citeyearpar{}`: [2020]
- `\citefullauthor{}`: Akiyama, Feng, Kempton, and Persson

See also a book Parry [6] and an arXiv preprint [5]. More multi-authors citation like Benoist and Quint [2] and Fan, Lau, and Rao [4].

*Remark 3.1.* For the use of `natbib` and format of arXiv preprint, it is recommended to use the `.bst` files `*nat.bst` or `*natDOI.bst` at

<https://github.com/zfengg/toolkit/tree/master/tex/bst>.

Otherwise, all the other default `bst` styles suffices.

### REFERENCES

- [1] S. Akiyama, D.-J. Feng, T. Kempton, and T. Persson. On the Hausdorff dimension of Bernoulli convolutions. *Int. Math. Res. Not. IMRN*, (19):6569–6595, 2020. 4
- [2] Y. Benoist and J.-F. Quint. *Random walks on reductive groups*, volume 62 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern*

- Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer, Cham, 2016. [4](#)
- [3] K. Falconer. *Fractal geometry*. John Wiley & Sons, Inc., Hoboken, NJ, second edition, 2003. Mathematical foundations and applications. [1](#)
- [4] A.-H. Fan, K.-S. Lau, and H. Rao. Relationships between different dimensions of a measure. *Monatsh. Math.*, 135(3):191–201, 2002. [4](#)
- [5] D.-J. Feng. Dimension of invariant measures for affine iterated function systems. *arXiv preprint arXiv:1901.01691*, 2020. [4](#)
- [6] W. Parry. *Topics in ergodic theory*, volume 75 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge-New York, 1981. [4](#)

DEPARTMENT OF MATHEMATICS, THE CHINESE UNIVERSITY OF HONG KONG, SHATIN, HONG KONG

*Email address:* `zfeng@math.cuhk.edu.hk`