

*Leçon
sur
la
théorie
géné-
rale
des
sur-
faces
Lessons
on
the
gen-
eral
the-
ory
of
sur-
faces*
?¹
2

*in-
ter
alia*

The
num-
bers
be-
tween
the
square
brack-
ets
re-
fer
to
en-
tries
in
the
Bib-
li-
og-
ra-
phy
In-
dex
at
the
end
of
the
book.
Let
us
enun-
ci-
ate
three

$$\begin{array}{c} \mathbf{T} \\ \mathbf{A} \\ 1,2, \vec{\mathbf{I}}_3 \\ \mathbf{A} \\ \mathbf{T} \\ \mathbf{M} \\ x,y,z \\ \mathbf{AM} \end{array}$$

$$\frac{dM}{dt} = \frac{dA}{dt} + x \frac{d_1}{dt} + y \frac{d_2}{dt} + z \frac{d_3}{dt}.$$

$$\omega_i = \xi_i(t)dt, \omega_{ji} = p_{ji}(t)dt, (i = 1, 2, 3; j < i).$$

\mathbf{T}_3
 \mathbf{T}_3
 \mathbf{T}_1
 $t =$
 \mathbf{T}_2
 $t =$
 \mathbf{T}_3^*
 \mathbf{T}_3^*
 ω_3^*
 ω_3^*
 \mathbf{T}_3
 \mathbf{T}_1^*
 \mathbf{T}_1^*
 \mathbf{T}_3^*
 \mathbf{T}_3^*
 \mathbf{T}_2^*
 \mathbf{T}_2^*
 \mathbf{T}_2^*
 \mathbf{T}^*
 \mathbf{T}^*
 \mathbf{T}^*

Con-
 sider
 bi-
 jec-
 tive
 re-
 spon-
 dance
 es-
 tab-
 lished
 be-
 tween
 two
 pa-
 ram-
 eter
 fam-
 ilies
 of
 tri-
 he-
 drals
 and
 For
 a
 sin-
 gle
 dis-
 place-
 ment
 to
 suf-
 fice
 to
 per-
 pose
 all
 the
 re-
 spond-
 ing
 tri-
 he-
 drals,
 it
 is
 nec-
 es-
 sary
 and
 suf-
 fi-
 cient
 that
 the
 cor-
 spon-
 dance
 in
 ques-
 tion
 makes
 the
 com-
 po-
 nents
 of
 in-
 finites-
 i-

$$10$$

$$x(t)$$

$$y(t)$$

$$z(t)$$

$$^2 +$$

$$y'^2 +$$

$$\tilde{0}, \tilde{z}'^2 =$$

$$x'x'' +$$

$$y'y'' +$$

$$\tilde{0}, \tilde{z}'z'' =$$

\mathbf{O}_{xyz}
 \mathbf{O}
 $\prime\prime\prime$
 \mathbf{O}

\mathbf{O}_{xyz}
 \mathbf{O}_{xyz}

\mathbf{A}_{123}

$$(16) \quad (\)_1^2 \times_{12} \times_{21} \times_{32} \times_{12}^2 \times_{33} \times_{13} \times_{23}^2 is (0) 01010100.$$

$$(17) \quad {}_1\wedge_2=i_{1,\,2}\wedge_3=i_{3,\,3}\wedge_1=i_2$$

$$(18) \quad ({}_1\wedge_2)\times_3=i.$$

the-
rem

Every
tri-
he-
dral
 \mathbf{A}_{123}
sat-
is-
fy-
ing
eq:2.3
and
eq:2.5
is
a
right
cyclic
tri-
he-
dral.

\mathbf{O}_3
 $x\mathbf{O}_y$
 \mathbf{O}
 $\mathbf{O}_{\underline{z}}$
 $\mathbf{O}_{\underline{z}}$

$$\omega_1$$

$$\begin{array}{c} C \\ C \\ T \\ C \\ T \\ C \\ T \\ T \\ T \\ T \\ T_0 \\ T_0 \end{array}$$

$$\omega_1=\omega_1,\omega_{11}=\omega_{11},\omega_{21}=\omega_{21},$$

$$\begin{array}{c} T_0 \\ T_0 \\ C \\ C \\ fun- \\ da- \\ men- \\ tal \\ con- \\ di- \\ tions \\ of \\ equal- \\ ity \\ ?? \end{array}$$

$$\begin{array}{c} C \\ C \end{array}$$

$$\begin{array}{l} \omega_{12}= \\ 0 \\ d_1= \\ \omega_{111} \\ 14 \end{array}$$

$$\begin{array}{c} k \\ d\sigma \\ ?? \end{array}$$

$$\begin{array}{c} de- \\ ter- \\ min- \\ ing \\ the \\ dis- \\ place- \\ ments \\ which \\ per- \\ mitted \\ two \\ given \\ ori- \\ ented \\ min- \\ i- \\ mal \\ curves \\ C \\ and \\ C \\ C \\ C \\ C \\ C \\ C \\ C \end{array}$$

$$(28) \quad k(\sigma)=k(\sigma),d\sigma=d\sigma.$$

$$\begin{array}{c} C \\ C \end{array}$$

$$(29) \quad k(\sigma)=k(\sigma),d\sigma=-d\sigma.$$

$$\begin{array}{c} C \\ C \\ spe- \\ cial \\ case \\ C \\ C \\ C \\ C \end{array}$$

$$\begin{matrix} 15 \\ P \\ (\geq \\ 0) \\ P \\ P \\ P \end{matrix}$$

16
 G

$\overset{G}{\mathbf{A}}_{123}$
 $\overset{1}{G}$
 G

Comparison
of
two
re-
orth
or-
der
tri-
he-
drals
 \mathbf{A}_{123}
and
 \mathbf{B}_{123} .

$\{\mathbf{A} = \mathbf{B} + \rho_1$ (*this signifies*

$\overrightarrow{\mathbf{BA}}\rho_1$
 $\overset{1}{\underset{2}{\overset{1}{=}}}$
 $\overset{2}{\underset{3}{\overset{1}{=}}}$
 $\overset{2}{\underset{3}{\overset{1}{=}}}\cos\theta +_3$
 $\overset{2}{\underset{3}{\overset{1}{=}}}\sin\theta,$
 $\overset{2}{\underset{3}{\overset{1}{=}}}\sin\theta +_3$
 $\overset{2}{\underset{3}{\overset{1}{=}}}\cos\theta,$
 $(38)\rho$
 θ

$\overset{1}{\underset{2}{\overset{1}{=}}}$
 $\overset{1}{\underset{2}{\overset{1}{=}}}$

six
 ω_i
 ω_{ji}
 \mathbf{A}_{123}
two

$$\omega_2 = \bar{\omega}_2 + (\rho + \lambda k) \bar{\omega}_{12},$$

$$\begin{array}{c}
20 \\
\mathbf{A}_{123} \\
\mathbf{A} \\
\uparrow \\
\mathbf{k} \\
k
\end{array}$$

$$\begin{array}{l} 21 \\ \omega_{12} \\ \omega_1 + \\ k\omega_{21} \end{array}$$

$$(48) \qquad \omega_{12} = \omega_1 + k\omega_{21}.$$

$$\begin{array}{c} 1 \\ 3 \\ -3 \end{array}$$

$$\begin{array}{l} \textit{pri-} \\ \textit{ta-} \\ \textit{tion} \end{array}$$

$$\begin{array}{l} k \\ \textit{The} \\ \textit{sec-} \\ \textit{ond} \\ \textit{der-} \\ \textit{tact} \\ \textit{prob-} \\ \textit{lem} \end{array}$$

\mathbf{P}^α
 \mathbf{P}_{123}

$k =$

\mathbf{A}
 \mathbf{P}

k

$??$

$$\omega_{12} = dk, \omega_1 + k\omega_{21} = 0.$$

(52)

sec-
ond

der

tri-

he-
dral

SECOND

OR-

DER-

CON-

TACT

PROB-

LEM.

Con-

sider

two

gen-

er-

a-

tri-

cs

G_0

and

G_0

of

two

sur-

faces

and

a

dis-

place-

ment

is-

ing

a

con-

tact

of

or-

der

\geq

Γ

along

these

two

gen-

er-

a-

tri-

ces:

is

per-

un-

poses

one

of

the

first

or-

der

tri-

he-

dral

\mathbf{T}_0

at-

tached

at

G_0

to

a

first

or-

der

tri-

he-

dral

\mathbf{T}_0

at-

tached

at

G_0 .

For

the

of

der

of

con-

tact

to

be

greater

than

it

is

rec-

sary

and

23

$$\begin{array}{l} \beta_0\alpha_0,\beta_0\gamma_0,\beta_0\delta_0,\beta_0\mathbf{P}),\\ X=(\alpha_0\beta_0,\alpha_0\gamma_0,\alpha_0\delta_0,\alpha_0\mathbf{P}),\\ Z=\\ \mathbf{I}\\ \mathbf{M}\\ X=(\beta\alpha,\beta\gamma,\beta\delta,\beta\mathbf{M});Y=(\alpha\beta,\alpha\gamma,\alpha\delta,\alpha\mathbf{M});Z=1. \end{array}$$

$$(59) \quad \begin{array}{l} n\\ \{x'=ax+by+cz,y'=a'x+b'y+c'z,z'=a''x+b''y+c''z,\; abca'b'c'a''b''c''=0. \end{array}$$

$$\begin{array}{l} a\\ b\\ c\\ c'.\\ (x,y,z)\\ g\\ m-\\ g\\ lytic\\ point\\ ge-\\ b-\\ met-\\ r-\\ cal\\ point\\ (x,y,z)\\ T\\ \mathbf{M}'(x',y',z')\\ T(T'\mathbf{M})=\\ (TT')\mathbf{M}\\ p\\ \mathbf{M}_1(x_1,y_1,z_1),\mathbf{M}_2(x_2,y_2,z_2),\ldots, \end{array}$$

$$\begin{array}{l} {}_1x_1+\\ {}_2x_2+\\ \dot{a}_p\dot{x}_p^+=\\ 0,\\ {}_1y_1+\\ {}_2y_2+\\ \dot{a}_p\dot{y}_p^+=\\ 0,\\ {}_1z_1+\\ {}_2z_2+\\ \dot{a}_p\dot{z}_p^+=\\ 0,\\ {}_1\mathbf{M}_1+{}_2\mathbf{M}_2+\ldots+{}_p\mathbf{M}_p=0. \end{array}$$

$$\begin{array}{l} \mathbf{M}'_1\\ \dot{\mathbf{M}}'_p\\ {}_1\mathbf{M}'_1+{}_2\mathbf{M}'_2+\ldots+{}_p\mathbf{M}'_p=0. \end{array}$$

$$[\mathbf{M}_1,\mathbf{M}_2,\mathbf{M}_3]$$

$$x_1y_1z_1x_2y_2z_2x_3y_3z_3$$

$$[\mathbf{M}'_1,\mathbf{M}'_2,\mathbf{M}'_3]=[\mathbf{M}_1,\mathbf{M}_2,\mathbf{M}_3].$$

$$\mathbf{A}^0(0,0,1),\mathbf{A}^0_1(1,0,0),\mathbf{A}^0_2(0,1,0).$$

$$\begin{array}{l} \mathbf{R}_0\\ \mathbf{R}_0\\ \mathbf{R}\\ \mathbf{A}(c,c',c''),\mathbf{A}_1(a,a',a''),\mathbf{A}_2(b,b',b''). \end{array}$$

$$\begin{array}{l} \mathbf{A}\\ \mathbf{A}_1\\ \mathbf{A}_2\\ [\mathbf{A},\mathbf{A}_1,\mathbf{A}_2]=1 \end{array}$$

$$\begin{array}{l} \mathbf{R}\\ \mathbf{X}\\ \mathbf{Y}\\ \mathbf{Z}\\ \mathbf{R}\\ \mathbf{M}\\ \mathbf{R}\\ \mathbf{R}_0\\ \mathbf{M}\\ \mathbf{P}\\ \mathbf{X}\\ \mathbf{Y}\\ \mathbf{Z} \end{array}$$

$$\mathbf{P}=X\mathbf{A}_1^0+Y\mathbf{A}_2^0+Z\mathbf{A}_3^0.$$

$$\mathbf{M} = X\mathbf{A}_1 + Y\mathbf{A}_2 + Z\mathbf{A}_3.$$

$$(60) \quad \begin{array}{l} T_\epsilon\\ \xi_T\\ T_\xi\\ x_i' = x_i + \epsilon \eta_i(x_1,\ldots,x_n) + \cdots \\ \xi_{\hat{x}_i} \end{array}$$

rel-
a-
tive
com-
po-
nents
ab-
so-
lute
com-
po-
nents
with
a
sign
change
pa-
ram-
e-
ters

Example
 $x' =$
 $\alpha x +$
 $b,$
 $x' =$
 $\alpha x +$
 β

$$\alpha = \frac{1}{\alpha}, \beta = -\frac{b}{a}.$$

??

$$\omega_1(a,b,da,db) = \frac{da}{a}, \omega_2(a,b,da,db) = \frac{db}{a}.$$

$$\begin{aligned} 1(\alpha, \beta, d\alpha, d\beta) &= \\ -\frac{da}{a} &= \\ \frac{d\alpha}{\alpha}, & \\ 2(\alpha, \beta, d\alpha, d\beta) &= \\ -\frac{db}{a} &= \\ \alpha d\left(\frac{\beta}{\alpha}\right) &= \\ d\beta - & \\ \frac{\beta}{\alpha} d\alpha. & \end{aligned}$$

There-
fore
every
group
is
holoedri-
cally
isomor-
phic
to
its
pa-
ram-
eter
group.

Θ_a
 ξ
 ξ'
the
pa-
ram-
eter
group
is
sim-
ply
tran-
si-
tive.

ξ
 Θ_a
 $\xi' =$
 $\Theta_a \xi$
??

$$\omega_p(\xi', d\xi') = \omega_p(\xi, d\xi).$$

the
pa-
ram-
eter
group
leaves
the
Pfaff-
ian
forms
 $\omega_p(\xi, d\xi)$
in-
vari-
ant

$\Omega(\xi, d\xi)$
 $\omega_p(\xi, d\xi)$
??
 $\tilde{\Omega}(\xi, d\xi)$

$$\Omega(\xi, d\xi) = a_1(\xi)\omega_1(\xi, d\xi) + \dots + a_r(\xi)\omega_r(\xi, d\xi).$$

Θ_a
 ξ
 ξ'

$$a_1(\xi)\omega_1(\xi, d\xi) + \dots + a_r(\xi)\omega_r(\xi, d\xi) = a_1(\xi')\omega_1(\xi', d\xi') + \dots + a_r(\xi')\omega_r(\xi', d\xi').$$

$$\omega_p(\xi, d\xi) = \omega_p(\xi', d\xi')$$

$$[a_1(\xi) - a_1(\xi')]\omega_1(\xi, d\xi) + \dots + [a_r(\xi) - a_r(\xi')]\omega_r(\xi, d\xi) = 0.$$

$$\begin{matrix} 31 \\ S \end{matrix}.$$

$$1, S_1, \ldots, S_n, S,$$

$$S_1, S_1^{-1}S_2, S_2^{-1}S_3, \ldots, S_{n-1}^{-1}S_n, S_n^{-1}S$$

$$?$$

36
 \mathbf{R}_1
 \mathbf{R}_1
 g_1
 G_1
 \mathcal{G}_1
 \mathcal{G}_1
 \mathcal{G}_1
 $Sg_1\mathbf{R}_1$
 \mathcal{G}
 \mathcal{G}

Example.
practically

Generalities
of the operation called orientation.
 G
 g_1
 g_1
 g_1
 g_1
 g_1
 ω_1
 $g_1\mathbf{R}_1$
 ω_1
 G

38

δ
 \mathbf{A}_1

r
 $n-$

r
 x_r^1

x_r^{+1}
 x_1^1

x_n
 q

a
 class
 for
 which
 G
 op-
 er-
 ates
 tran-
 si-
 tively
 is
 equiv-
 a-
 lent
 to
 a
 body
 c_1
 de-
 fined
 by
 the
 sub-
 group
 g_1
 leav-
 ing
 at
 ob-
 ject
 ω_1
 ar-
 bi-
 trar-
 ily
 cho-
 sen
 from
 the
 class
 ω
 in-
 vari-
 ant.

$$\begin{array}{l} \mathbf{R} = \\ \mathbf{S} \mathbf{R}_0 \\ \mathbf{S} \mathbf{A} \\ \mathbf{S} \\ \mathbf{R} \end{array}$$

Every
sim-
ply
tran-
si-
tive
group
is
sim-
ilar
to
its
pa-
ram-
eter
group.

$$\begin{array}{l} 44 \\ u_1 \\ u_2 \\ u_{r-n} \end{array}$$

$$\begin{array}{l} a_1 \\ u_r \\ (x_1, \dots, x_n) \\ (x'_1, \dots, x'_n) \end{array}$$

$$\begin{array}{l} x'_i = \phi_i(x_1, \dots, x_n; a_1, \dots, a_r), \\ (97) \quad (x_1, \dots, x_n; u_1, \dots, u_{r-n}) \\ (x'_1, \dots, x'_n) \\ x'_1 \\ x_p \\ u_1 \\ u_{r-n} \end{array}$$

$$(98) \quad \{x'_i = \phi_i(x_1, \dots, x_n; a_1, \dots, a_r), (i = 1, \dots, n), u'_j = \psi_j(u_1, \dots, u_{r-n}; x_1, \dots, x_n; a_1, \dots, a_r), (j = 1, \dots, r - n).$$

$$(106) \quad \begin{matrix} \{\xi'_1 = \psi_1(\xi_1, \xi_2, \dots; a_1, \dots, a_r), \dots \\ G(\lambda) \end{matrix}$$

$$\begin{matrix} \mu \\ \lambda \\ \mu \\ G(\mu) \\ G \\ G(\lambda) \\ \mu \\ \xi_1 \\ \xi_2. \\ \lambda \\ \zeta_1 \\ \zeta_2. \\ \delta \\ \mu \end{matrix}$$

$$G(\mu)$$

\mathcal{B}

$$(114) \quad x'_i = \phi_i(x_1, \dots, x_n; a_1, \dots, a_r), (i = 1, 2, \dots, n).$$

$$\omega_i(X, dX) = \omega_i(x, dx).$$

$$(118) \quad \begin{aligned} & \begin{pmatrix} x'_1, x'_2, \dots, x'_n \\ x'_1 + \\ dx'_1, \dots, x'_n + \\ dx'_n \\ (X_i + \\ dX_i) \\ (X_i) \\ (x'_i) \end{pmatrix} \\ & dX_i = \alpha_{i1}(X_1, \dots, X_n; x'_1, \dots, x'_n) dx'_1 + \dots + \alpha_{in}(X_1, \dots, X_n; x'_1, \dots, x'_n) dx'_n. \\ & \begin{pmatrix} \alpha_{i1}(X_1, \dots, X_n; x_1, \dots, x_n) dx_1 + \\ \dots + \\ \underline{\alpha_{in}}(X_1, \dots, X_n; x_1, \dots, x_n) dx_n \\ \alpha_{i1}(X_1, \dots, X_n; x'_1, \dots, x'_n) dx'_1 + \\ \dots + \\ \alpha_{in}(X_1, \dots, X_n; x'_1, \dots, x'_n) dx'_n. \\ (X_i) \\ r \end{pmatrix} \\ & \alpha_{i1}(X_1, \dots, X_n; x_1, \dots, x_n) dx_1 + \dots + \alpha_{in}(X_1, \dots, X_n; x_1, \dots, x_n) dx_n \end{aligned}$$

$$\begin{pmatrix} x_i \\ x'_i \end{pmatrix}$$

Remark.

$$\begin{pmatrix} ?? \\ \mathbf{R}_0 \\ \mathbf{A} \end{pmatrix}$$

$$(X_i)$$

$$\omega_i(x, dx)$$

$$(x_i +$$

$$dx_i)$$

$$(x_i)$$

$$a_i$$

$$S_a$$

$$(x_i)$$

$$\mathbf{R}_0 \equiv$$

$$\mathbf{A}$$

$$(X_i +$$

$$dX_i)$$

$$(x_i +$$

$$dx_i)$$

$$S_a$$

$$dX_i = \omega_i(x, dx),$$

$$i =$$

$$\phi_i(x_1, \dots, x_n; a_1, \dots, a_n),$$

$$dX_i =$$

$$\frac{\phi_i(x_1, \dots, x_n; a_1, \dots, a_n)}{x_1} dx_1 +$$

$$\dots +$$

$$\frac{\phi_i(x_1, \dots, x_n; a_1, \dots, a_n)}{x_n} dx_n.$$

$$\omega_i(x, dx)$$

$$a_i$$

Example:

$$a$$

$$dX =$$

$$X dx$$

$$X =$$

$$1$$

$$dx$$

$$dx'x' =$$

$$dx$$

$$x =$$

$$x_1$$

$$x_2$$

$$x_n$$

$$u_1$$

$$u_{r-n}$$

$$r$$

$$\omega_p(x, u, dx, du).$$

$$(119) \quad \left\{ \sum_{p=1}^r \omega_p(a, da) X_p f = \sum_{j=1}^n \frac{f}{x_j} \delta x_j; \sum_{p=1}^r \frac{\phi_i(x_1, \dots, x_n; a_1, \dots, a_r)}{a_p} da_p = \sum_{j=1}^n \frac{\phi_i(x_1, \dots, x_n; a_1, \dots, a_r)}{x_j} \delta x_j (i = 1, \dots, n). \right.$$

Group

of

pla-

nar

dis-

place-

50
of
the
system
of
partial
differential
equations
is
equal
to
the
number
of
classes
51
of
the
variables
 u_i ,
 v_i ,
 w_m
introduced
in
paragraph
1
2
3

??

G
 p
 p
 p
 x_1
 x_p
 G
 r

in-
transi-
tive
group
 g
 p
characterised
leaving
in-
variant

p
functions
 y_1 ,
 y_p
 x_1 ,
 x_n

$$\begin{array}{l}53\\G(V)\end{array}$$

$$\begin{array}{l}G\\G(V)\\(\xi_1,\ldots,\xi_r)\\(x_1,\ldots,x_n;u_1,\ldots,u_{r-n})\\V\\x_1=\\x_n=\\ \omega_1^1\\ \omega_n^1\\V\\du_j\\n\end{array}$$

$$\begin{array}{l}A\\first\\case\\which\\can\\arise\\u_j^n\end{array}$$

$$\begin{array}{l}V^0(x_i^0)\\V^1(x_i^1)\\G\\(x_i^0,u_k^0)\\V^0\\(x_i^1,u_k^1)\\V^1\\G\\V^0\\V^1\\r^-\\n\\G(V)\\n\\ \omega_i(x,dx)\\V^0\\V^1\\??\\G(V)\\G\end{array}$$

$$\begin{array}{l}If\\the-\\pre-\\ced-\\ing\\case\\does\\not\\arise\\ \omega_1^1\\ \omega_n^1\\u_{r-n}\end{array}$$

$$\omega_i=\sum_{k=1}^n\alpha_{ik}(x,u)dx_k,(i=1,\ldots,n).$$

$$\begin{array}{l}u_j\\ \alpha_{ik}\\u_{p+1}\\u_{r-n}\end{array}$$

con-
tract
el-
ments
 λ
 V_λ
 D_λ
the-
ret-
i-
cally
pos-
si-
ble
 $\gamma\gamma$
prac-
ti-
cally

\mathbf{A}
 μ_0
 $0 \dots$
 μ_{P-1}
 $P \dots$

The
~~con-~~
~~ta~~~~ct~~
~~el-~~
~~ment~~
~~of~~
~~or-~~
~~der~~
~~P~~
~~of~~
~~a~~
~~point~~
~~A~~
~~of~~
 ~~V_λ~~
~~will~~
~~be~~
~~dis-~~
~~tin-~~
~~guished~~
~~by~~
~~the~~
~~fam-~~
~~ily~~
~~of~~
~~frames~~
~~of~~
~~or-~~
~~der~~
~~P~~
~~of~~
~~this~~
~~point~~
~~and~~
~~by~~
~~the~~
~~set~~
~~of~~
~~in-~~
~~vari-~~
~~ants~~
~~of~~
~~or-~~
~~der~~
 $< P$
~~of~~
~~the~~
~~point.~~

$P+$
 $\frac{1}{P}$

given
any
two
points
of
 V_λ
there
always
ex-
ists
an
op-
er-
a-
tion
of
 G
trans-
form-
ing
 V_λ
into
it-
self
and
trans-
form-
ing
the
two
points
into
each
other?

$$(2) \quad {}_1\wedge_2=1.$$

$$\begin{array}{l} \mathbf{R}_{da} \\ \mathbf{R}_0 \\ \omega_1(0,da) \\ \omega_2(0,da) \\ \mathbf{A} \\ 1+ \\ \omega_{11}(0,da) \\ \omega_{12}(0,da) \\ {}^1 \\ \omega_{21}(0,da) \\ 1+ \\ \omega_{22}(0,da) \\ 2 \end{array}$$

$$\omega_{11}(0,da)+\omega_{22}(0,da)=0.$$

$$\omega_1(0,da),\omega_2(0,da),\omega_{11}(0,da),\omega_{12}(0,da),\omega_{21}(0,da)$$

$$\begin{array}{l} \mathbf{R}_0 \\ \mathbf{R}_{da} \end{array}$$

$$\begin{array}{l} ?? \\ rel- \\ a- \\ tive \\ com- \\ po- \\ nents \\ \mathbf{A}_{12} \\ \omega \end{array}$$

$$(3) \quad \left\{ \overrightarrow{d\mathbf{A}} = \omega_1 \ (a,da)_1 + \omega_2 \ (a,da)_2, d_1 = \omega_{11}(a,da)_1 + \omega_{12}(a,da)_2, d_2 = \omega_{21}(a,da)_1 - \omega_{11}(a,da)_2. \right.$$

$$\begin{array}{l} 0 \\ 1 \\ 0 \\ frames \\ of \\ or- \\ der \\ 0 \\ \mathbf{A} \\ \mathbf{A}_{12} \\ \mathbf{A} \\ frames \\ of \\ or- \\ der \\ 1 \\ \mathbf{A} \\ \mathbf{A}_{12} \\ 1 \\ 0 \\ 1 \end{array}$$

$$\begin{array}{l} G= \\ 5 \\ 0 \\ v_0= \\ 3 \\ v_1= \\ 2 \\ 0 \\ r_0^- = \\ 2 \\ 1 \\ v_0- \end{array}$$

$$\begin{array}{l} 73 \\ \omega_{00} \\ \omega_{01} \\ \omega_{02} \\ \omega_{10} \\ \omega_{11} \\ \omega_{12} \\ \omega_{20} \\ \omega_{21} \\ \{d\mathbf{A} = \omega_{00}\mathbf{A} + \omega_{01}\mathbf{A}_1 + \omega_{02}\mathbf{A}_2, d\mathbf{A}_1 = \omega_{10}\mathbf{A} + \omega_{11}\mathbf{A}_1 + \omega_{12}\mathbf{A}_2, d\mathbf{A}_2 = \omega_{20}\mathbf{A} + \omega_{21}\mathbf{A}_1 + \omega_{22}\mathbf{A}_2, \\ ??\omega_{00}+ \\ \omega_{11} \vdash \\ \omega_{22} = \\ 0. \end{array}$$

$$\begin{array}{c} 0 \\ 1 \\ 0 \\ 1 \\ 0 \\ 0 \end{array}$$

$$\begin{array}{l} G= \\ 6 \\ 8 \\ 0 \\ v_0= \\ 6 \\ v_1= \\ 5 \\ 0 \\ r_0^- \\ v_0= \\ 2 \\ 1 \\ v_0^- \\ v_1= \\ 1 \end{array}$$

$$\begin{array}{l} 0 \\ d\mathbf{A} \\ \mathbf{A} \\ \omega_{01} = \\ \omega_{02} = \\ 0 \\ \omega_{01} \\ \omega_{02} \\ 0 \\ 1 \\ d\mathbf{A} \\ \mathbf{A} \\ \mathbf{A}_1 \\ \omega_{02} = \\ 0 \\ 0 \\ 1 \end{array}$$

$$\omega_{01},\omega_{02}(=0)$$

$$\begin{array}{c} 2 \\ 1 \\ \mathbf{A}\mathbf{A}_1\mathbf{A}_2 \\ \mathbf{B}\mathbf{B}_1\mathbf{B}_2 \end{array}$$

$$(3) \left\{ \mathbf{A} \; = \lambda \mathbf{B}, \mathbf{A}_1 = \lambda_1 \mathbf{B}_1 + \mu \mathbf{B}, \mathbf{A}_2 = \lambda_2 \mathbf{B}_2 + \rho \mathbf{B}_1 + \sigma \mathbf{B}, \; where \lambda \lambda_1 \lambda_2 = 1. \right.$$

$$\begin{array}{c} 1 \\ \lambda \\ \lambda_1 \\ 1 \end{array}$$

$$\begin{array}{l} 1 \\ \mathbf{B}\mathbf{B}_1\mathbf{B}_2 \\ \lambda \\ \lambda_1 \\ \lambda_2 \\ d\mathbf{A}_1 \\ \mathbf{A} \\ \mathbf{A}_1 \\ \omega_{12} = \\ 0 \\ \omega_{12} \\ 1 \\ b \\ 1 \\ \omega_{12} = \\ b\omega_0 \end{array}$$

$$\begin{array}{l} 74 \\ \omega_{01} \\ \omega_{12} \\ \bar{\omega}_{01} \\ \bar{\omega}_{12} \\ \lambda \\ \lambda_1 \\ \backslash \end{array}$$